

A New Oxyprenyl Coumarin and Highly Methylated Flavones from the Exudate of *Ozothamnus lycopodioides* (Asteraceae)

Angel Rumbero^a, Fco. Javier Arriaga-Giner^b and Eckhard Wollenweber^{c,*}

^a Departamento de Química Orgánica, Universidad Autónoma de Madrid, Cantoblanco, E-28049 Madrid, Spain

^b Tabacalera S. A.; Centro I+D, c/ Embajadores 51, E-28012 Madrid, Spain

^c Institut für Botanik der Technischen Universität, Schnittspahnstrasse 3, D-64287 Darmstadt, Germany. Fax: 06151/166878. E-mail: Wollenweber@bio.tu-darmstadt.de

* Author for correspondance and reprint requests

Z. Naturforsch. **55c**, 1–4 (2000); received November 5, 1999

Ozothamnus lycopodioides, Asteraceae, Leaf and Stem Exudate, Flavones, New Oxyprenyl Coumarin

A new oxyprenyl coumarin was isolated from the lipophilic exudate of *Ozothamnus lycopodioides*. Its structure was established as 7-(3,3'-dimethylallyloxy)-5-hydroxy-6-methoxycoumarin from its UV, MS and NMR spectral data, especially two dimensional experiments. In addition to six earlier reported flavonols, we found four highly substituted flavones, including two rare methylenedioxyflavones.