

Novel Spiciferone Derivatives from the Fungus *Drechslera hawaiiensis* Isolated from the Marine Sponge *Callyspongia aerizusa*

Ru Angelie Edrada^a, Victor Wray^b, Albrecht Berg^c, Udo Gräfe^c, Sudarsono^d, Gernot Brauers^a, and Peter Proksch^{a,*}

^a Heinrich Heine Universität Düsseldorf, Institut für Pharmazeutische Biologie, Gebäude 26.23, Universitätsstr.1, D-40225 Düsseldorf, Germany.

Fax: +492118111923. E-mail: proksch@uni-duesseldorf.de

^b Gesellschaft für Biotechnologische Forschung mbH, Mascheroder Weg 1, D-38124, Braunschweig, Germany

^c Hans-Knöll Institut für Naturstoffforschung, Beutenbergstr. 11, D-07745 Jena, Germany

^d Research Center for Traditional Medicine, Faculty of Pharmacy, Gadjra Mada University, Yogyakarta, Indonesia

* Author for correspondence and reprint requests

Z. Naturforsch. **55c**, 218–221 (2000); received November 5/December 20, 1999

Sponge-Associated Fungi, *Drechslera hawaiiensis*, Spiciferone

From the marine sponge *Callyspongia aerizusa* collected from the Sea of Bali, Indonesia, fungal isolates of *Drechslera hawaiiensis* were obtained. Culture filtrates of the fungus yielded four spiciferone derivatives which include spiciferone A (**1**) and B (**2**), and two other novel derivatives including spiciferol A (**3**) which is an alcohol congener of spiciferone A (**1**) and compound **4** which is an monocyclic spiciferone congener featuring a butoxyl side chain. The structures of the novel compounds were established on the basis of NMR spectroscopic (¹H, ¹³C, COSY) and mass spectrometric (EIMS) data.