

Sulfonated Triterpenoid Saponins from *Fagonia indica*

Kamel H. Shaker^a, Mirko Bernhardt^a, M. Hani A. Elgamal^b and
Karlheinz Seifert^{a,*}

^a University of Bayreuth, Organic Chemistry I/2, NW II, D-95440 Bayreuth, Germany.
Fax: 49-921-5 553 58. E-mail: karlheinz.seifert@uni-bayreuth.de

^b National Research Centre, Laboratory of Natural Products, Dokki-Cairo, Egypt

* Author for correspondence and reprint requests

Z. Naturforsch. **55c**, 520–523 (2000); received April 5, 2000

Fagonia indica, Triterpenoid Saponins, Indicasaponin C and D

Two new triterpenoid saponins, 3-*O*-{[β -D-4-*O*-sulfonylglucopyranosyl-(1 \rightarrow 3)]- α -L-arabinopyranosyl}-ursolic acid-28-*O*-[β -D-glucopyranosyl] ester (indicasaponin C) and 3-*O*-{[β -D-4-*O*-sulfonylglucopyranosyl-(1 \rightarrow 3)]-[β -D-xylopyranosyl-(1 \rightarrow 2)]- α -L-arabinopyranosyl}-ursolic acid-28-*O*-[β -D-glucopyranosyl] ester (indicasaponin D) have been isolated from *Fagonia indica*. The structures were determined primarily by NMR spectroscopy. The assignment of NMR signals was performed by means of ¹H–¹H COSY, NOESY, ROESY, TOCSY, HMQC and HMBC experiments.