

A Pyoverdin from *Pseudomonas* sp. CFML 95–275^s

Razia Sultana, Regine Fuchs, Hans Schmickler, Karin Schlegel,
Herbert Budzikiewicz^{a,*}, Bina Shaheen Siddiqui^b, Valérie Geoffrey^c
and Jean-Marie Meyer^c

^a Institut für Organische Chemie der Universität zu Köln, Greinstr. 4, 50939 Köln, Germany.

Fax: +49–221–470–5057. E-mail: h.budzikiewicz@uni-koeln.de

^b HEJ Research Institute, University of Karachi, Karachi-75270, Pakistan

^c Laboratoire de Microbiologie et Génétique, Université Louis Pasteur,
UPRS-A 7010 du CNRS, 28 rue Goethe, 67000 Strasbourg, France

* Author for correspondence and reprint requests

Z. Naturforsch. **55c**, 857–865 (2000); received August 18, 2000

Pseudomonas, Pyoverdin, Siderophore

From *Pseudomonas* sp. CFML 95–275 a pyoverdin was isolated with a cyclopeptidic substructure. It could be shown that this pyoverdin is identical with one obtained from *Pseudomonas fluorescens* BTP 7 for which a lactone structure had been deduced from the interpretation of a FAB spectrum. The elucidation of the correct structure of the pyoverdin is described.