

11,17-Dideoxyagelorsin A and B, New Bromotyrosine Derivatives and Analogs from the Marine Sponge *Suberea aff. praetensa*

Anake Kijjoa,^{a,b} Rawiwan Watanadilok,^{b,c} Pichai Sonchaeng^c, Artur M. S. Silva^d, Graham Eaton^e and Werner Herz^{f,*}

^a Instituto de Ciências Biomédicas de Abel Salazar, Universidade do Porto, 4099-003 Porto, Portugal

^b Centro do Estudos de Química Orgânica, Fitoquímica e Farmacologia da Universidade do Porto, Rua Anibal Cunha, 4050-017, Porto, Portugal

^c Bangsae Institute of Marine Science (BIMS), Burapha University, Bangsae, Chonburi 21031, Thailand

^d Departamento de Química, Universidade de Aveiro, 3810 Aveiro, Portugal

^e Department of Chemistry, University of Leicester, University Road, Leicester, LE1 7RH, UK

^f Department of Chemistry, The Florida State University, Tallahassee, FL 32306-4390, USA. Fax: 1-850-644-8281. E-mail: jduin@chem.fsu.edu

* Author for correspondence and reprint requests;

Z. Naturforsch. **56c**, 1116-1119 (2001); received February 8/July 30, 2001

Suberea aff. praetensa, 11-Dideoxyagelorsin A and B, Bromotyrosine Derivatives

A collection of the marine sponge *Suberea aff. praetensa* from the Gulf of Thailand furnished the bromotyrosine derivatives fistularin-3, agelorsins A and B and the new dideoxyagelorsins A and B.

Introduction

All sponge genera of the order Verongida (class Demospongia, subclass Ceractinomorpha) which have so far been examined chemically contain secondary metabolites derived from bromo- or chlorotyrosine in which the side chain has been converted into a variety of nitrogenous groups while the aromatic ring has been retained or has undergone rearrangement or reduction (Bergquist and Wells, 1980). A subclass of these metabolites consist of compounds such as the fistularins (Gopichand and Schmitz, 1979) in which one or two modified tyrosine moieties are attached to a chain consisting of variously modified 3,5-dibromo-4-(γ -amino propoxy)-phenylethyl amines (Cimino *et al.*; 1983, Kernan *et al.*, 1990; Gunasekera and Cross, 1992; Ciminiello *et al.*, 1994, 1996, 1997; Compagnone *et al.*, 1999). The first and so far only bromotyrosine derivatives isolated from a non-verongid sponge were the agelorsins A (**1a**) and B (**1b**) found (Koenig and Wright, 1993) together with 11-epifistularin B in *Agelas oroides* (Demospongia, subclass Tetractinomorpha, order Axinellida, family Agelasidae). We now report isolation from a Gulf of Thailand collection of *Suberea aff. praetensa* (Demospongia, Ceractinomorpha, Ver-

ongida, family Aplysinellidae) of the agelorsins A and B, the new 11,17-dideoxyagelorsins A and B (**2a** and **2b**) and the related, fistularin-3 (Gopichand and Schmitz, 1979) as well as clionasterol.

Materials and Methods

¹H and ¹³C NMR spectra were recorded at ambient temperature on a Bruker AMC instrument operating at 300.13 and 75.47 MHz, respectively. EI mass spectra were measured on a Hitachi Perkin-Elmer RMV-6M instrument. For HRMS samples were run using +FAB ionization with Xe gas at 6KV on a KRATUS CONCEPT III, 2 sector mass spectrometer. The accelerating voltage was 8 KV. Silica gel for column chromatography was Si gel 60 (0.2-0.5 mm) Merck, for analytical and preparative TLS Si gel G-60 GF 254 Merck.

Animal material

Suberea aff. praetensa was collected from a trawl net on the seashore of Ban Phae village on the Gulf of Thailand, Rayong Province, Thailand, in March 1998. The sponge was identified by Prof. Rob van Soest, Department of Coelenterates and Porifera, Zoological Museum, University of Amsterdam. A voucher (BIMS-1954) was deposited

0939-5075/2001/1100-1116 \$ 06.00 © 2001 Verlag der Zeitschrift für Naturforschung, Tübingen · www.znaturforsch.com · D





Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

**1a** R = OH Agelorin A**2a** R = H**1b** R = OH Agelorin B**2b** R = H

at the Reference Collection Museum of the Institute of Marine Science, Burapha University, Bangsaen, Chonburi, Thailand. The collected material was immediately frozen at -20°C for one night prior to extraction.

Extraction and isolation of the constituents

The sample (164 g net weight) was thawed, homogenized with EtOH (500 ml), allowed to stand in a dark chamber for 24 hours and filtered. The residue on the filter paper was again extracted with EtOH (3×500 ml). The aqueous alcoholic extracts were combined, evaporated at reduced pressure until the final volume of the aqueous solution was ca. 100 ml and extracted with EtOAc (3×300 ml). The EtOAc extracts were combined and concentrated at reduced pressure to give a

residue (3.4 g). The latter was chromatographed over Si gel (50 g) and eluted with petrol- CHCl_3 , CHCl_3 - Me_2O and CHCl_3 -MeOH, 125 ml frs being collected as follows: frs 1–27 (petrol- CHCl_3 , 4:1), 28–46 (petrol- CHCl_3 , 3:2), 47–71 (petrol- CHCl_3 , 2:3), 72–86 (petrol- CHCl_3 , 1:4), 87–98 (CHCl_3), 99–145 (CHCl_3 - Me_2O , 9:1), 146–160 (CHCl_3 - Me_2O , 7:3), 161–177 (CHCl_3 - Me_2O , 1:1), 178–189 (CHCl_3 -MeOH, 9:1). The ratios refer to v/v.

Frs 5–8 (210 mg) were recrystallized from petrol to give clionasterol (176 mg) identified by MS and ^1H NMR spectrometry. Frs 54–59 (29 mg) were purified by TLC (Si gel, CHCl_3 -petrol-EtOAc, 8:1:1) to give 1-hexene (12 mg). Frs 70–80 (54 mg) on purification by TLC (Si gel, CHCl_3 -MeOH- HCO_2H , 85:15:0.5) gave a non-crystalline mixture of 11,17-dideoxyagelorin A and B (28 mg). Frs 105–144 (230 mg) were purified by

TLC (Si gel, CHCl₃-Me₂OH-HCO₂H, 70:30:1) to give fistularin-3 (120 mg) and a non-crystalline mixture of agelorin A and agelorin B (70 mg). Fistularin-3 was identified by HRMS, ¹H and ¹³C NMR measurements (¹H, ¹³C, NOESY, HMBC), conversion to the tetraacetate, ¹H and ¹³C NMR measurements of the latter and comparison of the data with the literature (Gopichand and Schmitz, 1979; Gunasekera and Cross, 1992; Campagnone *et al.*, 1999). No attempt was made to separate the mixture of agelorin A (**1a**) and agelorin B (**1b**) whose structures were confirmed by HRMS and extensive ¹H and ¹³C NMR measurements (¹H, ¹³C, COSY, HMBC) in DMSO solution for comparison with the mixture of new dideoxy derivatives **2a** and **2b**.

11,17-Dideoxyagelorins A and B (**2a** and **2b**).

Due to the small amount of material no attempt was made to separate the noncrystalline mixture, + FAB-HRMS (NBA) M + H⁺ 1054.68209; calcd. for ¹²C₂₉ ¹H₂₇ ¹⁴N₄ ¹⁶O₉ ⁷⁹Br₃ ⁸¹Br₃ 1054.68207. ¹H and ¹³C NMR spectra are listed in Table I, assignments being based on decoupling, COSY and HMBC correlations. Comparison with the ¹H and

¹³C NMR spectra of the agelorin A-agelorin B mixture under the same conditions established identity except for the absence of the hydroxyl groups at C-11 and C-17.

Discussion

As mentioned earlier brominated metabolites derived from tyrosine have so far been found only in sponges of the order Verongida, the exception being *Agelas oroides*. The only previous report on the chemistry of a *Suberea* species mentions the occurrence of the bromotyrosine derivatives (+) aerothoinin, homoaerothoinin, (+) aeroplysinin-1 and haloverongiaquinols in *Suberea creba* from the Coral Sea (Debitus *et al.*, 1998).

Acknowledgments

Work in Portugal was supported by Fundação para Ciências e Tecnologia (Unidade de I & D 226/94), FEDER and PRAXIS XXI. We thank Mr. Sumait Puchakarn of BIMS for collection of the sponge. Raviwan Watanadilok thanks Burapha University for a scholarship and support for his work in Thailand.

Table I. ¹H and ¹³C NMR spectra of 11,17-dideoxyagelorins A and B^a.

Position	δH(mult)A,A' ^b	δC(mult)A,A' ^c	δH(mult)B,B' ^b	δC(mult)B,B' ^c
1,1'	4.18 <i>d</i> (11,4)	73.34 <i>d</i> , 73.29 <i>d</i>	4.29 <i>brs</i>	72.01 <i>d</i>
2,2'	5.19 <i>dd</i> (11,3)	57.81 <i>d</i>	5.35 <i>m</i>	55.84 <i>d</i>
3,3'		183.81 <i>s</i>		183.39 <i>s</i>
4,4'		121.34 <i>s</i> , 121.31 <i>s</i>		122.75 <i>s</i>
5,5'	7.68 <i>s</i> , 7.65 <i>s</i>	149.06 <i>d</i> , 148.97 <i>d</i>	7.54 <i>s</i>	145.35 <i>d</i>
6,6'		90.51 <i>s</i>		89.38 <i>s</i>
7a, 7'a	3.65 <i>d</i> (18), 3.63 <i>d</i> (18)	38.03 <i>t</i>	3.65 <i>d</i> (18), 3.63 <i>m</i>	38.03 <i>t</i> , 39.79 <i>t</i>
7b, 7'b	3.13 <i>d</i> (18), 3.10 <i>d</i> (18)		3.33 <i>d</i> (18), 3.30 <i>d</i> (18)	
8,8'		153.92 <i>s</i> , 153.89 <i>s</i>		154.57 <i>s</i> , 153.92 <i>s</i>
9,9'		158.85 <i>s</i> ^d		158.78 <i>s</i> ^d
10a,b	3.42 <i>m</i>	36.31 <i>t</i>	3.42 <i>m</i>	36.31 <i>t</i>
11a,b	2.00 <i>q</i> (6)	29.62 <i>t</i>	2.00 <i>q</i> (6)	29.62 <i>t</i>
12a,b	3.95 <i>dd</i> (6,4)	71.24 <i>t</i>	3.95 <i>dd</i> (6,4)	71.24 <i>t</i>
13		150.75 <i>s</i>		150.75 <i>s</i>
14		117.38 <i>s</i>		117.38 <i>s</i>
15,15'	7.53 <i>s</i>	133.04 <i>d</i>	7.53 <i>s</i>	133.04 <i>d</i>
16		133.88 <i>s</i>		133.88 <i>s</i>
17a,b	2.76 <i>t</i> (7)	33.20 <i>t</i>	2.76 <i>t</i> (7)	33.20 <i>t</i>
18a,b	3.42 <i>m</i>	39.79 <i>t</i>	3.42 <i>m</i>	39.79 <i>t</i>
NH _a	8.67 <i>dd</i> (11.8,5.7)		8.67	
NH _b	8.64 <i>dd</i> (11.8, 5.7)		8.64	
OH	6.77 <i>br</i>		6.77 <i>br</i>	

^a From mixture of **2a** and **2b**. A and A', B and B' refer to the left and right halves of the structures of A and B. Spectra recorded in DMSO-d₆; assignments by HMBC and COSY.

^b J values in parentheses.

^c Multiplicities deduced by DEPT.

^d Assignments may be interchanged.

- Berquist P. R. and Wells R. J. (1980), Chemotaxonomy of the Porifera: The development and current status of the field. In: Marine Natural Products, Chemical and Biological Perspectives, Vol. 5, (P. J. Scheuer, ed.) Academic Press, New York and London, pp. 1–50.
- Ciminiello P., Dell'Aversano C., Fattorusso E., Magno S., Carrano L. and Pansini M. (1996), Chemistry of Verongida Sponges VII. Bromocompounds from the Caribbean sponge *Aplysina archeri*. *Tetrahedron* **52**, 9863–9868.
- Ciminiello P., Fattorusso E., Forino M., Magno S. and Pansini M. (1997), Chemistry of Verongida sponges VIII. Bromocompounds from the Mediterranean sponges *Aplysina aerophoba* and *Aplysina cavernicola*. *Tetrahedron* **53**, 6565–6572.
- Ciminiello P., Fattorusso E., Magno S. and Pansini M. (1994), Chemistry of Verongida sponges III. Constituents of a Caribbean *Verongula* sp. *J. Natural Products* **57**, 1564–1569.
- Cimino G., DeRosa S., De Stefano S., Self R. and Sodano G. (1983), The bromo-compounds of the true sponge *Verongia aerophoba*. *Tetrahedron Lett.* **24**, 3029–3032.
- Compagnone R. S., Avila R., Suárez A. I., Abrams O. V., Rangel H. R., Arvela F., Piña I. G. and Merento E. (1999), 11-Deoxyfistularin-3, a new cytotoxic metabolite from the Caribbean sponge *Aplysina fistularis insularis*. *J. Natural Products* **62**, 1443–1444.
- Debitus C., Guella C., Mancini I., Waikedre J. Guemas J.-P., Nicolas J. L. and Pietra F. (1998), Quinolones from a bacterium and tyrosine metabolites from its host sponge *Suberea creba* from the Coral Sea. *J. Marine Biotechnol.* **6**, 136–148; chemical Abstracts 129, 328033x.
- Gopichand Y. and Schmitz F. J. (1979), Marine Natural Products: Fistularin-1, 2 and -3 from the sponge *Aplysina fistularis forma fulva*. *Tetrahedron Lett.* 3921–3924.
- Gunasekera S. P. and Cross S. S. (1992), Fistularin 3 and 11-keto-fistularin 3, feline leukemia virus active bromotyrosine metabolites from the marine sponge *Aplysina archeri*.
- Kernan M. R., Cambie R. C. and Bergquist P. R. (1990), Chemistry of sponges VII. 11,19-Dideoxyfistularin 3 and 11-hydroxyaerothonin, bromotyrosine derivatives from *Pseudoceratina durissima*. *J. Natural Products* **53**, 615–622.
- König G. M. and Wright A. D. (1993), Agelorins A and B and 11-epifistularin-3, three new antibacterial fistularin-3 derivatives from the tropical marine sponge *Agelas oroides*. *Heterocycles* **36**, 1351–1359.