## The Himanimides, New Bioactive Compounds from Serpula himantoides (Fr.)Karst.

Pedro Aqueveque<sup>a,b</sup>, Timm Anke<sup>b,\*</sup> and Olov Sterner<sup>c,\*</sup>

<sup>a</sup> Universidad de Conception, Casilla 2407, Apartado 10, Conception 3, Chile
 <sup>b</sup> Lehrbereich Biotechnologie der Universität, Paul-Ehrlich-Straße 23,
 D-67663 Kaiserslautern, Germany. Fax: +49-6312052999. E-mail: anke@rhrk.uni-kl.de

S-221 00 Lund, Sweden. Fax: +46-462228209. E-mail: Olov.Sterner@orgk2.lth.se

\* Authors for correspondence and reprint requests

Department of Organic and Bioorganic Chemistry, Lund University, P. O.Box 124,

Z. Naturforsch. 57 c, 257-262 (2002); received November 7/December 3, 2001

succinimide and maleimide derivatives, of which two are N-hydroxylated.

Himanimides, Antibiotics, Basidiomycete

In a screening of basidiomycete cultures from Chile for the production of antibiotics we identified a *Serpula himantoides* strain as a producer of metabolites inhibiting the growth of bacteria and fungi. Bioactivity guided purification resulted in the isolation of four new antibiotics. Their structures were elucidated by spectroscopic methods. All four compounds are