

Flavonoids from *Ficaria verna* Huds.

Michał Tomczyk^{a,*}, Jan Gudej^a and Marek Sochacki^b

^a Department of Pharmacognosy, Faculty of Pharmacy, Medical Academy of Białystok, Mickiewicza Str. 2a, 15–230 Białystok, Poland. Fax: +48-85-7424907.

E-mail: tomczyk@amb.edu.pl

^b Centre of Molecular and Macromolecular Studies, Department of Structural Studies, Mass Spectrometry Facility, Polish Academy of Sciences, 112 Sienkiewicza Str., 90-363 Łódź, Poland

* Author for correspondence and reprint requests

Z. Naturforsch. **57c**, 440–444 (2002); received January 7/February 7, 2002

Ranunculaceae, *Ficaria verna*, Flavonoids

A phytochemical investigation of the flowers and leaves of *Ficaria verna* Huds. (*Ranunculaceae*) yielded four additional known flavonoid compounds including: kaempferol 3-O- β -D-(6''- α -L-rhamnopyranosyl)-glucopyranoside (nicotiflorin), apigenin 8-C- β -D-glucopyranoside (vitexin), luteolin 8-C- β -D-glucopyranoside (orientin) and apigenin 8-C- β -D-(2''-O- β -D-glucopyranosyl)-glucopyranoside (flavosativaside). The characterisation of these compounds was achieved by various chromatographic and spectroscopic methods (UV, ¹H NMR, ¹³C NMR and MS).