

Two Epimeric Flavalignans from *Trichilia catigua* (Meliaceae) with Antimicrobial Activity

Moacir G. Pizzolatti^{a,*}, Andreia F. Venson^a, Artur Smânia Júnior^b,
Elza de F. A. Smânia^b and Raimundo Braz-Filho^c

^a Departamento de Química, Universidade Federal de Santa Catarina,
98010-970, Florianópolis-SC, Brazil. E-mail: moacir@qmc.ufsc.br

^b Departamento de Microbiologia, Universidade Federal de Santa Catarina,
98010-970, Florianópolis-SC, Brazil

^c Setor de Química de Produtos Naturais – LCQUI – CCT,
Universidade Estadual do Norte Fluminense, 28015-620, Campos-RJ, Brazil

* Author for correspondence and reprint requests

Z. Naturforsch. **57c**, 483–488 (2002); received January 9/March 6, 2002

Trichilia catigua, Flavalignans, Antimicrobial Activity

A mixture of flavalignan cinchonains Ia and Ib was isolated from the bark of *Trichilia catigua*. The structures were established on the basis of spectroscopic data of the natural products and their methylated derivatives including 2D NMR experiments, and compared with data in the literature. These flavalignans exhibited antibacterial activity against *Bacillus cereus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*.