

Oleanane Triterpenoids from *Cedrela montana* (Meliaceae)

Leonardo Castellanos^a, Rosabel S. de Correa^a, Eduardo Martínez^a,
and José S. Calderon^{b,*}

^a Departamento de Química, Universidad Nacional de Colombia, Ciudad Universitaria,
Santa Fé de Bogotá Colombia

^b Instituto de Química, Universidad Nacional Autónoma de México, Circuito Exterior,
Ciudad Universitaria, Coyoacán 04510. México D. F. Fax: (52 55) 56 16 22 03.
E-mail: uscalder@correo.unam.mx

* Autor for correspondence and reprints requests

Z. Naturforsch. **57c**, 575–578 (2002); received February 8/March 22, 2002

Cedrela montana, Meliaceae, Oleanan-28,13 β -olides

Two new oleanane-type triterpenes, characterized as 3-oxo-11 α ,12 α -epoxy-oleanan-28,13 β -olide and 3-oxo-olean-11-en-28,13 β -olide, were isolated from the fruits and seeds of *Cedrela montana* (Meliaceae). In addition, the known compounds oleanonic acid, a mixture of β -sitosterol and stigmasterol, and the limonoid photogedunin were also isolated. The structures of the new compounds were established by spectroscopic methods, including 2D NMR.