

# **A New Triterpene Saponin from *Chenopodium ficifolium***

Ahmed A. Gohar<sup>a,\*</sup>, Galal T. Maatooq<sup>a</sup>, Masatake Niwa<sup>b</sup> and Takaya Yoshiaki<sup>b</sup>

<sup>a</sup> Pharmacognosy Department, Faculty of Pharmacy, Mansoura University, Mansoura 33516, Egypt. E-mail: gohar@mans.edu.eg

<sup>b</sup> Department of Medicinal Resources Chemistry, Faculty of Pharmacy, Meijo University, Tempaku, Nagoya 468–8503, Japan

\* Author for correspondence and reprint requests

Z. Naturforsch. **57c**, 597–602 (2002), received March 11/April 12, 2002

Chenopodiaceae, *Chenopodium ficifolium*, Triterpenes

The new triterpene saponin 3-O- $\beta$ -D-glucopyranoside, 28- $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranosiduronic acid oleanolate was isolated from the roots of *Chenopodium ficifolium*. The known compounds stigmasterol-3-O-glucoside and 3-O- $\beta$ -D-glucopyranosiduronic acid, 28- $\beta$ -D-glucopyranosyl oleanolate were also isolated. The latter compound, oleanolic acid,  $\beta$ -sitosterol and its glucoside were isolated from the aerial parts. The identity of these compounds was verified through different chemical and physico-chemical evidences including different 1D and 2D NMR experiments.