

Nuatigenin-Type Steroidal Saponins from *Veronica fuhsii* and *V. multifida*

Meltem Ozipek,^a Iclal Saracoglu,^a Yukio Ogiwara^b and İhsan Çaliş^{a,*}

^a Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University,
TR-06100 Ankara, Turkey. Fax: +90-3 12-3 11 47 77. E-mail: icalis@hacettepe.edu.tr

^b Department of Pharmacognosy, Faculty of Pharmaceutical Sciences, Nagoya
City University, Tanabe-dori, Mizuho-ku, Nagoya 467-8603, Japan

* Author for correspondence and reprint requests

Z. Naturforsch. **57c**, 603–608 (2002); received April 8/May 10, 2002

Veronica, Steroidal Glycosides, Multifidoside

A new nuatigenin-type steroidal saponin, multifidoside (**2**), was isolated from the aerial parts of *Veronica fuhsii* and *V. multifida* and its structure was identified as 3-*O*-{[α -L-rhamnopyranosyl-(1 \rightarrow 2_{glu})]-[β -D-glucopyranosyl-(1 \rightarrow 4_{rha})- α -L-rhamnopyranosyl (1 \rightarrow 4_{glu})]- β -D-glucopyranosyl}nuatigenin 26-*O*- β -D-glucopyranoside. Additionally, a known steroidal saponoside, aculeatiside A (**1**), from *V. fuhsii*, a phenylethanoid glycoside, verpectoside A (**3**), and a flavon glycoside, isoscutellarein 7-*O*-(2''-*O*-6''-*O*-acetyl- β -D-allopyranosyl- β -D-glucopyranoside) (**4**) from *V. multifida* were isolated.