

Constituents and Biological Activities from *Muehlenbeckia hastulata*

Silvia Erazo^a, Orlando Muñoz^b, Rubén García^a, Igor Lemus^a, Nadine Backhouse^a, Rosa Negrete^a, Arturo San Feliciano^c and Carla Delporte^a

^a Department of Pharmacological and Toxicological Chemistry, School of Chemical and Pharmaceutical Sciences, University of Chile, P. O. Box 233, Santiago 1, Chile. Fax: 56-2-2227900. Email: serazo@uchile.cl

^b Department of Chemistry, Faculty of Sciences, University of Chile

^c Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Salamanca, 37007 Salamanca, Spain

* Author for correspondence and reprint requests

Z. Naturforsch. **57c**, 801–804 (2002); received February 2/May 23, 2002

Muehlenbeckia hastulata, Chemical Constituents, Biological Activities

Muehlenbeckia hastulata (J. E.Sm.) Johnst. used mainly for its abortive properties was found active for oxytoxic and analgesic activities in a biological activity screening (oxytoxic, analgesic, antimicrobial, cytotoxic and xanthine oxidase assay). Epicatechin, emodin-8-glycoside and rutin were isolated from the aerial part and the root of the plant.