

A New Phenylpropanoid Ester from the Bark of *Zanthoxylum scandens* (Rutaceae)

Quang An Nguyen^a, Hanh Trinh Van-Dufat^a, Sylvie Michel^a, François Tillequin^{a,*}, Vincent Dumontet^b and Thierry Sévenet^b

^a Laboratoire de Pharmacognosie de l'Université René Descartes, U. M. R./C. N. R. S. N° 8638, Faculté des Sciences Pharmaceutiques et Biologiques, 4 Avenue de l'Observatoire, 75006 Paris, France. E-mail: tillequi@pharmacie.univ-paris5.fr

^b Institut de Chimie des Substances Naturelles du C. N. R. S., 91190 Gif-sur-Yvette, France

* Author for correspondance and reprint requests

Z. Naturforsch. **57c**, 986–989 (2002); received July 15/August 2, 2002

Zanthoxylum scandens, Alkaloids, Phenylpropanoids, (*E*)-*O*-Geranylconyferyl Alcohol (9*Z*, 12*Z*)-linoleate

The alkaloids norchelerythrine, magnoflorine and (–)(*S*)-*O*-methylbalfourodinium cation were isolated from *Zanthoxylum scandens* bark collected in Vietnam, together with the flavanone glycoside hesperidin and the phenylpropanoids (*E*)-*O*-geranylconiferyl alcohol and (*E*)-*O*-geranylconiferyl alcohol (9*Z*, 12*Z*)-linoleate. This latter is a novel compound whose structure was elucidated on the basis of its spectral data and confirmed by chemical correlation.