

Flavonols from *Scurrula ferruginea* Danser (*Loranthaceae*)

Françoise Lohézic-Le Dévéhat^{a*}, Sophie Tomasi^a,
Didier Fontanel^b and Joël Boustie^a

^a Laboratoire de Pharmacognosie et de Mycologie,
U. P. R. E. S. 2234 “Extraction et synthèse de molécules à visée thérapeutique”, 2, Avenue du P^r Léon Bernard, 35043 Rennes Cedex.

Fax: 0033(0)223234711.

E-mail: lohezic.francoise@libertysurf.fr

^b IUP “Ingénierie en Nutraceutique”, Faculté de
Pharmacie, 2, Avenue du P^r Léon Bernard,
35043 Rennes Cedex

* Author for correspondence and reprint requests

Z. Naturforsch. **57c**, 1092–1095 (2002);
received August 6/September 2, 2002

Flavonols, Cytotoxic, *Scurrula ferruginea*

Three natural flavonols compounds have been isolated from the ethyl acetate fraction of *Scurrula ferruginea* Danser (*Loranthaceae*). Besides quercetin and quercitrin, an unusual flavonol glycoside 4''-O-acetylquercitrin was isolated. Structures were determined using spectroscopic methods including UV, NMR and HRMS-EI. The incidence of 4''-O-acetylquercitrin, not previously reported in the *Loranthaceae*, is discussed. Cytotoxic evaluation on four human cancer cell lines showed quercetin to be the most active with IC₅₀ of 35 µm on U251 (human glioblastoma cells).