

Herbal Remedies Traditionally Used Against Malaria in Ghana: Bioassay-Guided Fractionation of *Microglossa pyrifolia* (Asteraceae)[§]

Inga Köhler^a, Kristina Jenett-Siems^{*a}, Carola Kraft^a, Karsten Siems^b,
Daniel Abbiw^c, Ulrich Bienzle^d and Eckart Eich^a

^a Institut für Pharmazie (Pharmazeutische Biologie), Freie Universität Berlin, Königin-Luise-Strasse 2+4, D-14195 Berlin, Germany. E-mail: kjsiems@zedat.fu-berlin.de

^b Analyticon Discovery GmbH, Potsdam, Germany

^c Department of Botany, University Legon-Accra, Accra, Ghana

^d Institut für Tropenmedizin, Medizinische Fakultät Charité, Humboldt-Universität zu Berlin, Berlin, Germany

* Author for correspondence and reprint requests

Z. Naturforsch **57c**, 1022–1027 (2002); received August 27/September 6, 2002

Plasmodium falciparum, *Microglossa pyrifolia*, Bioassay-Guided Fractionation

Different extracts from 11 West African plants traditionally used against malaria in Ghana were tested against both the chloroquine-sensitive strain PoW and the chloroquine-resistant clone Dd2 of *Plasmodium falciparum*. Due to the promising *in vitro* activity of the lipophilic extract [IC₅₀: 10.5 µg/ml (PoW); 13.1 µg/ml (Dd2)], *Microglossa pyrifolia* (Lam.) Kuntze (Asteraceae) was chosen for further phytochemical investigation. From active fractions 13 compounds were isolated; their structures were established on the basis of spectroscopic methods. 1-Acetyl-6*E*-geranylgeraniol-19-oic acid and sinapyl diangelate represent new natural compounds. The two diterpenes *E*-phytol [IC₅₀: 8.5 µM (PoW); 11.5 µM (Dd2)], and 6*E*-geranylgeraniol-19-oic acid [IC₅₀: 12.9 µM (PoW); 15.6 µM (Dd2)] proved to be the most active constituents in our test system.

Introduction

In continuation of our research on medicinal plants with antiplasmodial activity we investigated 11 plants from Ghana, where malaria causes many deaths per year, particularly in young children (Browne *et al.*, 2000). *Gomphrena celosioides* (Amaranthaceae), *Picralima nitida* (Apocynaceae), *Pergularia daemia* (Asclepiadaceae), *Emilia sonchifolia* (Asteraceae), *Microglossa pyrifolia* (Asteraceae), *Adansonia digitata* (Bombacaceae), *Euphorbia hirta*, *Phyllanthus niruroides* (Euphorbiaceae), *Tetrapleura tetraptera* (Mimosaceae), *Mitragyna inermis*, and *M. stipulosa* (Rubiaceae) are widespread in the Greater Accra Region of Ghana and are used against malaria or fever by herbalists (Bruce, 1998 and 2000). Based on our screening results (Table I) we chose *Microglossa pyrifolia* (Lam.) Kuntze (*syn.*: *Conyza pyrifolia* Lam., Asteraceae) for further investigation. *M. pyrifolia* is a shrub widespread in tropical Asia and Africa (Do-

kosi, 1998). In Ghana an aqueous leaf decoction is used against fever and malaria. In other West African countries the leaf extract is taken against abdominal pains, rheumatism, diarrhoea and many other diseases (Neuwinger, 1994). Subsequent fractionation of the lipophilic extract coupled with an antiplasmodial bioassay (Desjardins *et al.*, 1979) was performed to detect the active principles responsible for the antiprotozoal activity of *M. pyrifolia*.

Experimental

General experimental procedures

For fractionation, silica gel 60 (63–200 µm) and reversed phase material (LiChroprep[®] RP-18, 40–63 µm) were used. Preparative high performance liquid chromatography (HPLC) was performed on a Knauer Eurochrom 2000 with Knauer pumping system and a Knauer WellChrom DAD K-2700 UV-detector equipped with an Eurosphere 100 C-18 (10 µm, 22 × 250 mm) column. For preparative thin layer chromatography (TLC) aluminum sheets (20 × 20 cm) coated with silica gel 60

[§] Part 7 in the series 'Herbal remedies traditionally used against malaria', for part 6 see Onegi *et al.* (2002).

F₂₅₄ were used. Mass spectra were determined with a Finnigan MAT CH7A (220 °C, 70 eV) and FAB-MS were recorded on a VARIAN MAT CH5DF (Xenone, 7 kV). ¹H-NMR spectra, ¹H-¹H-COSY, and HMBC experiments were obtained using acetone-*d*₆ and CDCl₃ as solvents with a Bruker AVANCE DPX 400 (400 MHz, TMS as internal standard). Cultures of *P. falciparum* were harvested with an Inotech cell harvester and IC₅₀ values determined by a liquid scintillation counting.

Plant material

The plant species were collected in the Greater Accra Region from February to March 2000 and April 2001 by scientists of the Department of Botany, Legon-University Accra in Ghana. Voucher specimens of the following plants have been deposited in the herbarium of the Department of Botany, Legon-University Accra: *Gomphrena celosioides* (GC 39071), *Picralima nitida* (GC 39064), *Pergularia daemia* (GC 39070), *Emilia sonchifolia* (39072), *Microglossa pyrifolia* (GC 47681), *Euphorbia hirta* (GC 39068), *Phyllanthus niruroides* (GC 39069), *Tetrapleura tetraptera* (39065), *Mitragyna inermis* (GC 39066), *Mitragyna stipulosa* (GC 39067). *Adansonia digitata* fruits were collected in April 2001 in Navrongo (Upper East Region).

Extraction and isolation

For the screening program, the air dried plant material (20 g) was crushed and extracted three times for 2 h with 150 ml petrol ether-EtOAc (1:1, v/v) at room temperature to gain the lipophilic extracts. Afterwards the plant material was air dried and treated three times with 150 ml MeOH to afford the hydrophilic extracts. The preparation of the aqueous extracts was carried out according to (Bruce, 2000): Stem barks and roots were boiled for one hour in an aqueous solution of NaHCO₃ (1 teaspoon/300 ml). Vegetative aerial parts or fruits were extracted by an infusion with such a solution.

For further investigations of *M. pyrifolia*, vegetative aerial parts (500 g) were ground and extracted for 24 h at room temperature with petrol ether-EtOAc 1:1 (2 l). This procedure was repeated three times. Solvents were evaporated at 40 °C under reduced pressure; the residue was subjected to the antiplasmodial test. The oily residue (30 g) was purified by column chromatogra-

phy on RP-18 material and sequentially eluted with MeOH-H₂O mixtures (4:6; 5:5; 6:4; 7:3; 8:2; 9:1, v/v), MeOH, and CHCl₃ (800 ml, each). Fractions 6–10 eluting with MeOH-H₂O (8:2; 9:1, v/v, 300 ml each) and MeOH (500 ml), respectively, proved to be the most active ones in the antiplasmodial assay. Fraction 6 (residue 120 mg) was chromatographed on silica gel (7 g) with CH₂Cl₂, CH₂Cl₂-MeOH (9:1, v/v) (200 ml each) to yield compound **13** (3 mg). Further purification of the remaining fraction 6A by HPLC with MeOH-H₂O (30:70, v/v) to MeOH within 40 min lead to **11** (60 mg, R_T = 29.64 min). Fraction 7 (250 mg) was separated on silica gel (12 g) with petrol ether-EtOAc (3:1 and 1:1, v/v respectively; 170 ml each) and CH₂Cl₂-MeOH (10:1, v/v, 200 ml) and finally by preparative TLC (CH₂Cl₂-MeOH 10:1, v/v) to yield acacetin **1** (3 mg, R_f = 0.42), compound **4** (5 mg, R_f = 0.52) and **5** (10 mg, R_f = 0.48). Fraction 8 (1 g) was purified on silica gel (40 g) with petrol ether-EtOAc gradient (10:1; 3:1; 1:1, v/v, respectively; 200 ml each) and yielded the active fractions 8A and 8B. Compound **6** (12 mg, R_f = 0.49) was isolated from 8A by preparative TLC (petrol ether-EtOAc 2:1, v/v). Purification of 8B by preparative TLC (CH₂Cl₂-MeOH 10:1, v/v) lead to **7** (30 mg, R_f = 0.57). Fraction 9 (6 g) was purified on silica (60 g) with petrol ether-EtOAc and CH₂Cl₂-MeOH gradients, respectively (10:1, 5:1 and 40:1, 20:1, v/v, respectively; 250 ml each) and lead to the active fractions 9A and 9B. Compounds **8** (3 mg, R_f = 0.60), **9**, and **10** (6 mg, R_f = 0.62) were isolated from 9A by preparative TLC (CH₂Cl₂-MeOH 10:1, v/v). From fraction 9B, using preparative TLC (CH₂Cl₂-MeOH 10:1 and 20:1, v/v, respectively), we isolated compound **12** (12 mg, R_f = 0.62). Fractionation of 10 (1.5 g) on silica (20 g) lead to linoleic acid **2** (15 mg); fraction 10A was purified by preparative TLC (CHCl₃-MeOH 9:1, v/v) and yielded **3** (2 mg, R_f = 0.47). Fractionation of the inactive aqueous extract by column chromatography on RP-18 material with MeOH-H₂O mixtures of decreasing polarity (up to 90% MeOH) lead to active fractions, which contained compounds **7** and **11**.

1-Acetyl-6E-geranylgeraniol-19-oic acid (12): ¹HNMR (400 MHz), acetone-*D*₆: δ 6.77 (1H, t, *J* = 7 Hz, H-6); 5.39 (1H, tq, *J* = 7 Hz, 1 Hz, H-2); 5.19 (1H, tq, *J* = 7 Hz, 1 Hz, H-10); 5.10 (1H, tq, *J* = 7 Hz, 1 Hz, H-14); 4.59 (2H, d, *J* = 7 Hz, H-1);

2.41–2.33 (4H, m, H-8, H-9); 2.25–2.12 (4H, m, H-4, H-5); 2.05 (3H, s, OAc); 1.98 (4H, obscured by solvent, H-12, H-13); 1.72 (3H, s, H-20); 1.66 (3H, s, H-18); 1.60 (3H, s, H-16 or H-17); 1.59 (3H, s, H-17 or H-16) ppm; (-)-FAB-MS: 361 [M-H]⁻.

Sinapyl diangelate (**13**): ¹HNMR (400 MHz), acetone-D₆: δ 6.90 (2 H, s, H-2, H-6), 6.73 (1 H, brd, *J* = 16 Hz, H-7), 6.47 (1H, td, *J* = 6 Hz, 16 Hz, H-8), 6.23 (1 H, m, H-3' or H-3''), 6.12 (1H, qq, *J* = 1 Hz, 7 Hz, H-3'' or H-3'), 4.81 (2 H, dd, *J* = 1 Hz, 6 Hz, H-9), 3.84 (6 H, s, OCH₃3 and OCH₃5); 2.20 (3 H, m, H-4'' or H-5''), 2.01 (3 H, obscured by solvent, H-5'' or H-4''), 1.97 (3 H, dq, *J* = 7 Hz, 1.5 Hz, H-4'), 1.90 (3 H, quint, *J* = 1.5 Hz, H-5') ppm; EI-MS: [M]⁺ 374 (5), 292 (12), 83 (100), 55 (46); HR-MS: *m/z* = 374.17273 (calculated 374.17294 for C₂₁H₂₆O₆), 292.13099 (calculated 292.13108 for C₁₆H₂₀O₅), 83.04966 (calculated 83.04969 for C₅H₇O).

Antiplasmodial activity

The antiplasmodial assay was performed by means of the microculture radioisotope technique as described previously (Jenett-Siems *et al.*, 2000). The concentration at which growth was inhibited by 50% (IC₅₀) was estimated by interpolation. IC₅₀ values > 50 µg/ml for extracts and IC₅₀ values > 25 µg/ml for fractions, respectively, were considered inactive (O'Neill *et al.* 1985).

Results and Discussion

Lipophilic extracts of all tested plant species exhibited a moderate antiplasmodial activity with *M. pyrifolia* aerial parts as the most active ones [IC₅₀: 13.1 µg/ml (Dd2), 10.5 µg/ml (PoW)]. Bioassay-guided fractionation of the aerial parts of *M. pyrifolia* revealed the fractions eluting from a RP-18

Table I. Activity of lipophilic[#] plant extracts against *Plasmodium falciparum*.

Plant family	Species	Plant part	Traditional uses	Mean IC ₅₀ values [µg/ml ^a]	
				PoW	Dd2
Amaranthaceae	<i>Gomphrena celosioides</i> Mart.	whole plant	antimalarial (Bruce, 1998)	36.9	31.5
Apocynaceae	<i>Picralima nitida</i> (Stapf) Th. & H. Durand	stem bark	substitute for quinine (Neuwinger, 1994), antimalarial (Bruce, 1998), (François <i>et al.</i> , 1996)	> 50	14.5
Asclepiadaceae	<i>Pergularia daemia</i> (Forssk) Chiov.	stem bark	remedy for cold (Dokosi, 1998), and fever (Bruce, 2000)	> 50	> 50
Asteraceae	<i>Emilia sonchifolia</i> (L.) DC.	aerial parts	fever remedy (Abbiw, 1990), antimalarial (Azuine, 1998)	36.9	43.6
Bombacaceae	<i>Microglossa pyrifolia</i> (Lam.) Kuntze	aerial parts, root	fever remedy (Abbiw, 1990), antimalarial (Bruce, 1998)	10.5	13.1
	<i>Adansonia digitata</i> L.	fruit pulp ^b	fever remedy (Abbiw, 1990), antimalarial (Azuine, 1998)	> 50	> 50
	Euphorbiaceae	<i>Euphorbia hirta</i> L.	whole plant	antimalarial (Dokosi, 1998)	> 50
Mimosaceae	<i>Phyllanthus niruroides</i> Müll. Arg.	aerial parts	antimalarial (Bruce, 1998)	11.5	27.3
	<i>Tetrapleura tetraptera</i> (Schuhmach. & Thonn.) Taub.	fruits	antimalarial (Neuwinger, 1994)	> 50	> 50
Rubiaceae	<i>Mitragyna inermis</i> (Willd.) Kuntze	leaves, stem bark, root	analgesic properties (Abbiw, 1990), antimalarial (Bruce, 1998), (Mustofa <i>et al.</i> , 2000)	> 50 39.6 > 50	34.6 14.6 28.2
	<i>Mitragyna stipulosa</i> (DC.) Kuntze	leaves, stem bark, root	antimalarial (Bruce, 1998), antidote to poison, e.g. (Abbiw, 1990)	32.6 > 50 > 50	20.4 36.1 48.7

[#]: Petrol ether-EtOAc (1:1, v/v).
^a: Performed in duplicate.
^b: Tested as aqueous extract.

column with 80% to 100% MeOH to be most active with IC_{50} values ranging from 2.5 to 18.7 $\mu\text{g/ml}$ against a chloroquine-sensitive strain (PoW) and a chloroquine-resistant clone (Dd2) of *Plasmodium falciparum*. Further separation by several chromatographic methods yielded 13 compounds. We isolated acacetin (**1**), linoleic acid (**2**), *E*-phytol (**3**), benzyl 2,6-dimethoxybenzoate (**4**) (Lu *et al.*, 1993), 13-hydroxy-octadeca-9*Z*,11*E*,15*Z*-trien-oic acid (**5**) (Reddy *et al.*, 1994), and 1-hydroxy-calamenene as a mixture of two isomers (**6**) (El-Seedi *et al.*, 1994). Furthermore, we obtained the following furanoditerpenes: strictic acid (**7**) (Tandon and Rastogi, 1979), hardwickiic acid (**8**) (Heymann *et al.*, 1994), 10 α -nidoresedic acid (**9**) and 10 β -nidoresedic acid (**10**). The spectral data of **9** and **10** were quite similar to those of the already known methyl esters (Singh *et al.*, 1988, Bohlmann and Fritz, 1978). Several active fractions yielded 6*E*-geranylgeraniol-19-oic acid (**11**) (Zdero *et al.*, 1990). The ^1H NMR of **12** was very similar to that of **11**. An additional singlet at 2.06 ppm (3H, s, OAc) pointed to an acetate group. Due to the presence of the acetate group, the signal for the two protons at C-1 displayed a chemical shift of 4.59 ppm (2H, d, $J = 7$ Hz, H-1) versus the signal for H-1 of **11** at 4.09 ppm (2H, d, $J = 7$ Hz, H-1). The (–) FAB MS gave a parent ion $[\text{M}-\text{H}]^-$ at m/z 361 thus confirming the structure and lead to the identification of **12** as 1-acetyl-6*E*-geranylgeraniol-19-oic acid. The isolation of the 6*E*-isomer has not yet been reported, in contrast to its 6*Z*-isomer (Herz and Kulanthaivel, 1985). The EIMS spectrum of **13** showed a molecular ion peak at m/z 374. The ion fragment peak at m/z 292 indicated the loss of an angelate group. In the ^1H NMR compound **13** displayed a singlet at 6.90 ppm (2H, s, H-2, H-6) and two

singlets for two methoxyl groups at 3.84 ppm (6H, s, OMe-3 and OMe-5), belonging to a methoxylated aromatic system. Additionally a doublet at 6.73 ppm (1H, d, $J = 16$ Hz, H-7) and a triplet doublet at 6.47 ppm (1H, td, $J = 6$ Hz, 16 Hz, H-8) pointed to a *trans*-configured double bond in conjugation with the aromatic system. The chemical shift of the allylic methylene group at 4.81 ppm (2H, dd, $J = 1$ Hz, 6 Hz, H-9) indicated an esterification at this position. Additionally typical signals for two angelate moieties could be observed: 6.23 ppm (1H, m, H-3' or H-3''), 6.12 ppm (1H, qq, $J = 1$ Hz, 6 Hz, H-3'' or H-3'), 2.20 ppm (3H, m, H-4'' or H-5''), 2.01 ppm (3H, obscured by solvent, H-5'' or H-4''), 1.97 ppm (3H, dq, $J = 7$ Hz, 1.5 Hz, H-4'), 1.90 ppm (3H, quint, $J = 1.5$ Hz, H-5'). Comparison with literature data of known sinapyl alcohol derivatives (Bohlmann *et al.*, 1969) lead to the identification of **13** as sinapyl diangelate, which represents a new natural compound. This is the first report on the occurrence of **1-10**, **12**, and **13** from *M. pyrifolia*.

In our antiplasmodial test system the compounds linoleic acid (**2**), *E*-phytol (**3**), 13-hydroxy-octadeca-9*Z*,11*E*,15*Z*-trien-oic acid (**5**), and 6*E*-geranylgeraniol-19-oic acid (**11**) exhibited an activity against *P. falciparum* with IC_{50} values between 2.5 $\mu\text{g/ml}$ and 13.7 $\mu\text{g/ml}$ (Table II). From several active fractions we isolated the diterpene 6*E*-geranylgeraniol-19-oic acid (**11**). Thus, we assume that the aliphatic unsaturated compounds of *M. pyrifolia* are representing the antiprotozoal principle of this species. In our test system the methanolic and aqueous extracts of all tested species did not show any activity. However, fractionation of the aqueous extract of *M. pyrifolia* lead to the detection (^1H NMR and EIMS spectra) of **7** and **11** in active fractions. Thus, **11** is detectable in the aqueous extract despite of its rather lipo-

Compound	Mean IC_{50} values*			
	PoW		Dd2	
	[$\mu\text{g/ml}$]	[μM]	[$\mu\text{g/ml}$]	[μM]
Linoleic acid (octadeca-9,12-dienoic acid) [#] (2)	6.1	21.8	8.7	31.1
<i>E</i> -Phytol (3)	2.5	8.5	3.4	11.5
Benzyl 2,6-dimethoxybenzoate (4)	9.0	33.1	> 25	> 91.9
13-Hydroxy-octadeca-9 <i>Z</i> ,11 <i>E</i> ,15 <i>Z</i> -trien-oic acid (5)	6.7	22.8	13.7	46.6
6 <i>E</i> -Geranylgeraniol-19-oic-acid (11)	4.3	12.9	5.2	15.6
Chloroquine $\times 2$ H_3PO_4	0.008	0.015	0.073	0.14

Table II. Antiplasmodial activity of compounds isolated from *Microglossa pyrifolia* against *Plasmodium falciparum* *in vitro*.

* Performed in triplicate.

[#] Previously tested (Köhler *et al.*, 2002).




Fig. 1. Structures of compounds isolated from *Microglossa pyrifolia*.

philic character. Therefore, we conclude that 6*E*-geranylgeraniol-19-oic acid (**11**) is mainly responsible for a moderate antiplasmodial activity of a traditional plant preparation from *M. pyrifolia*.

Acknowledgements

The authors are indebted to Dr. T. B. F. Bruce, phytotherapist in Accra-North Kaneshie (Ghana),

who recommended these plants for its antimalarial potential. This study was supported by grants from Humboldt-Universität zu Berlin (N 1/01/01) and the Hans-und-Stefan-Bernbeck-Stiftung to Inga Köhler, and from Deutsche Pharmazeutische Gesellschaft to Kristina Jenett-Siems.

- Abbiw D. K. (1990), Useful plants of Ghana, Intermediate Technology Publications and The Royal Botanic Gardens Kew, London, pp. 154–157.
- Azuine M. A. (1998), Cancer and ethnobotany of Nigeria, Shaker Verlag, Aachen.
- Bohlmann F. and Zdero C. (1969), Über ein neues Coniferylalkohol-Derivat aus *Blumea lacera*. *Tetrahedron Lett.* **2**, 69–70.
- Bohlmann F. and Fritz U. (1978), Neue Diterpene und Acetylenverbindungen aus *Nidorella*-Arten. *Phytochemistry* **17**, 1769–1772.
- Browne E. N., Frimpong E., Sievertsen J., Hagen J., Hamelmann C., Dietz K., Horstmann R. D. and Burchard G. D. (2000), Malariometric update for the rainforest and savanna of Ashanti region, Ghana. *Ann. Trop. Med. Parasitol.* **94**, 15–22.
- Bruce T. B. F. (1998 and 2000), personal communications, phytotherapist, Accra-North Kaneshie (Ghana).
- Desjardins R. E., Canfield C. J., Haynes J. D. and Chulay J. D. (1979), Quantitative assessment of antimalarial activity in vitro by a semiautomated microdilution technique. *Antimicrob. Agents Chemother.* **16**, 710–718.
- Dokosi, O. B. (1998), Herbs of Ghana, Ghana Universities Press, Accra, 313–314.
- El-Seedi H., Ghia F. and Torssell K. B. G. (1994), Cadinane sesquiterpenes from *Siparuna macrotepala*. *Phytochemistry* **35**, 1495–1497.
- François G., Aké Assi L., Holenz J. and Bringmann G. (1996), Constituents of *Picralima nitida* display pronounced inhibitory activities against asexual erythrocytic forms of *Plasmodium falciparum* in vitro. *J. Ethnopharmacol.* **54**, 113–117.
- Herz W. and Kulanthaivel P. (1985), Diterpenes and sesquiterpene lactones from *Mikania congesta*. *Phytochemistry* **24**, 1761–1768.
- Heymann H., Tezuka Y., Kikuchi T. and Supriyatna S. (1994), Constituents of *Sindora sumatrana* MIQ. III.: New *trans*-clerodane diterpenoids from the dried pods. *Chem. Pharm. Bull.* **42**, 1202–1207.
- Jenett-Siems K., Siems K., Jakupovic J., Solis P. N., Gupta M. P., Mockenhaupt F. P., Bienzle, U. and Eich E. (2000), Sipandinolide: A butenolide including a novel type of carbon skeleton from *Siparuna andina*. *Planta Med.* **66**, 384–385.
- Köhler I., Jenett-Siems K., Siems K., Hernández M. A., Ibarra R. A., Berendsohn W. G., Bienzle U. and Eich E. (2002), *In vitro* antiplasmodial investigation of medicinal plants from El Salvador. *Z. Naturforsch.* **57c**, 277–281.
- Lu T., Vargas D. and Fischer N. H. (1993), Sesquiterpenes from *Brintonia discoidea*. *Phytochemistry* **34**, 737–742.
- Mustofa, Valentin A., Benoit-Vical F., Pélessier Y., Koné-Bamba D. and Mallié M. (2000), Antiplasmodial activity of plant extracts used in the west African traditional medicine. *J. Ethnopharmacol.* **73**, 145–151.
- Neuwinger, H. D. (1994), Afrikanische Arzneipflanzen und Jagdgifte, Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart.
- Onegi B., Kraft C., Köhler I., Freund M., Jenett-Siems K., Siems K., Beyer G., Melzig M. F., Bienzle U. and Eich E. (2002), Antiplasmodial activity of naphthoquinones and one anthraquinone from *Stereospermum kunthianum*. *Phytochemistry* **60**, 39–44.
- O'Neill M. J., Bray D. H., Boardmann P., Phillipson J. D. and Warhurst D. C. (1985), Plants as sources of antimalarial drugs part 1. *In vitro* test method for evaluation of crude extracts from plants. *Planta Med.* **51**, 394–398.
- Reddy K. L. N., Reddy S. P. and Sharma G. V. M. (1994), Synthesis of 13-Hydroxy-9Z, 11E, 15Z-octadecatrienoic acid. *Synth. Commun.* **24**, 1395–1401.
- Singh P., Jain S. and Jakupovic J. (1988), Clerodane derivatives from *Grangea maderaspatana*. *Phytochemistry* **27**, 1537–1539.
- Tandon S. and Rastogi R. P. (1979), Strictic Acid, a novel diterpene from *Conyza stricta*. *Phytochemistry* **18**, 494–495.
- Zdero C., Ahmed A. A., Bohlmann F. and Mungai G. M. (1990), Diterpenes and sesquiterpene xylosides from East African *Conyza* species. *Phytochemistry* **29**, 3167–3172.