

Cytotoxic Hydroazulene Diterpenes from the Brown Alga *Cystoseira myrica*

Seif-Eldin N. Ayyad^{a*}, Osama B. Abdel-Halim^b, W. Thomas Shier^c, and
Thomas R. Hoye^d

^a Department of Chemistry, Faculty of Science, Mansoura University, New Damietta, Egypt.

Fax: (2)057.-403868. E-mail: snayyad@mans.edu.eg

^b Department of Pharmacognosy, Faculty of Pharmacy, Mansoura University,
Mansoura 35511, Egypt

^c Department of Medicinal Chemistry, University of Minnesota, Minneapolis, MN, 55455,
U. S. A.

^d Department of Chemistry, University of Minnesota, Minneapolis, MN 55455, U. S. A.

* Author for correspondence and reprint requests

Z. Naturforsch. **58c**, 33–38 (2003); received June 13/September 2, 2002

Cytotoxicity-guided fractionation of the alcohol extract of the brown alga, *Cystoseira myrica*, afforded four new cytotoxic hydroazulene diterpenes, dictyone acetate (**2**), dictyol F monoacetate (**4**), isodictytriol monoacetate (**6**), and cystoseirol monoacetate (**8**), together with two known cytotoxic hydroazulene diterpenes, pachydictyol A (**1**) and dictyone (**3**). The constitution of each isolated compound has been determined on the basis of spectroscopic and chemical evidence.

Key words: *Cystoseira*, Cytotoxic, Hydroazulenes