

Chemical Composition of the Essential Oils of Two *Rhodiola* Species from Tibet

Yidong Lei^a, Peng Nan^{a,b}, Tashi Tsering^c, Zhankui Bai^a, Chunjie Tian^a, and Yang Zhong^{a,*}

^a Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, School of Life Sciences, Fudan University, Shanghai 200433, China.
Fax: 86-21-65 64 24 68. E-mail: yangzhong@fudan.edu.cn

^b Shanghai Center for Bioinformation Technology, Shanghai 201203, China

^c Department of Chemistry, Biology and Geography, Tibet University, Lhasa 850000, China

* Author for correspondence and reprint requests

Z. Naturforsch. **58c**, 161–164 (2003); received October 21/November 18, 2002

The essential oils from rhizomes of *Rhodiola crenulata* and *R. fastigiata* in eastern Tibet were analyzed by using GC-MS. The major constituents were geraniol (53.3%), *n*-octanol (13.4%), 2-methyl-3-buten-2-ol (10.8%), citronellol (5.3%), 3-methyl-2-buten-1-ol (4.0%), myteol (3.0%), and linalool (2.4%) for *R. crenulata* and geraniol (45.3%), *n*-octanol (12.3%), 2-methyl-3-buten-2-ol (8.0%), linalool (5.1%), isogeraniol (4.5%), citronellol (4.4%), and *cis*-sabinenehydrate (3.6%) for *R. fastigiata*.

Key words: *Rhodiola crenulata*, *Rhodiola fastigiata*, Essential Oil