

Iridoids from *Globularia dumulosa*

Hasan Kirmizibekmez^a, Pinar Akbay^b, Otto Sticher^b, and İhsan Çalış^{a,*}

^a Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University,
TR-06100 Ankara, Turkey. Fax: +90-3 12-3 11 47 77. E-mail: icalis@hacettepe.edu.tr

^b Department of Applied BioSciences, Institute of Pharmaceutical Sciences, Swiss Federal
Institute of Technology (ETH) Zürich, Winterthurerstr. 190, CH-8057 Zürich, Switzerland

* Author for correspondence and reprint requests

Z. Naturforsch. **58c**, 181–186 (2003); received October 15/October 31, 2002

Two new iridoids, 10-*O*-benzoylglobularigenin (**1**) and dumuloside (**2**) were isolated from the aerial parts of *Globularia dumulosa* together with seven known iridoid glucosides, davisioside (**3**), aucubin (**4**), melampyroside (**5**), catalpol (**6**), 10-*O*-benzoylcatalpol (**7**), alpinoside (**8**) and deacetylalpinoside (**9**). Three phenylethanoid glycosides, verbascoside, decaffeoylverbascoside, leucosceptoside A and three flavone glucosides, pectolinarigenin 7-*O*-β-D-glucopyranoside, nepetin 7-*O*-β-D-glucopyranoside, demethoxycentaureidin 7-*O*-β-D-glucopyranoside were also isolated and characterized. The structure elucidation of the isolated compounds was performed by spectroscopic (UV, IR, HR-MALDIMS, 1D- and 2D NMR) methods.

Key words: *Globularia dumulosa*, Iridoids, Phenylethanoid Glycosides, Flavone Glycosides