

Sesquiterpenes from *Achillea pannonica* Scheele[§]

Ingrid Werner^{a*}, Sabine Glasl^a, Armin Presser^b, Ernst Haslinger^b, and Johann Jurenitsch^a

^a Institute of Pharmacognosy, University of Vienna, PharmaCenter, Althanstrasse 14, A-1090 Vienna, Austria. Fax: +431 42 77 95 52. E-mail: ingrid.werner@univie.ac.at

^b Institute of Pharmaceutical Chemistry, Karl-Franzens-University Graz, Universitätsplatz 1, A-8010 Graz, Austria

* Author for correspondence and reprint requests

Z. Naturforsch. **58c**, 303–307 (2003); received November 29/December 23, 2002

From dichloromethane extracts of flowerheads of *Achillea pannonica* SCHEELLE three sesquiterpenes were isolated and identified: 11,13-dehydrodesacetylmaticarin, (6E)-5-tigloxy-9-hydroxynerylol and α -longipin-2-en-1-one. The structures were determined by MS, IR and NMR spectroscopic analyses. (6E)-5-Tigloxy-9-hydroxynerylol is reported here for the first time. Additionally spathulenol, a compound of the essential oil was identified using GC-MS and GC-FTIR.

Key words: *Achillea pannonica*, Sesquiterpenes