

Iridoid and Bisiridoid Glycosides from *Globularia cordifolia*

Hasan Kirmizibekmez^a, Ihsan Çalış^{a,*}, Pinar Akbay^b, and Otto Sticher^b

^a Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University,
TR-06100 Ankara, Turkey. Fax: +90-3 12-3 11 47 77. E-mail: icalis@hacettepe.edu.tr

^b Department of Applied BioSciences, Institute of Pharmaceutical Sciences, Swiss Federal
Institute of Technology (ETH) Zurich, Winterthurerstr. 190, CH-8057 Zürich, Switzerland

* Author for correspondence and reprint requests

Z. Naturforsch. **58c**, 337–341 (2003); received December 19, 2002

From the methanolic extract of the underground parts of *Globularia cordifolia*, a new iridoid glycoside, 5-hydroxydavisioside (**1**) and a new bisiridoid glycoside, globuloside C (**2**) were isolated along with six known iridoid glycosides, aucubin, melampyroside, monomelittoside, globularifolin, alpinoside and asperuloside. The structures of the isolates were established by 1D and 2D NMR spectroscopy in combination with IR, UV and MS analyses.

Key words: *Globularia cordifolia*, Iridoid and Bisiridoid Glycosides