

# Iridoid Glucosides from *Veronica pectinata* var. *glandulosa*

U. Sebnem Harput<sup>a</sup>, Akito Nagatsu<sup>b</sup>, Yukio Ogiwara<sup>b,c</sup>, and Iclal Saracoglu<sup>a,\*</sup>

<sup>a</sup> Hacettepe University, Faculty of Pharmacy, Department of Pharmacognosy,  
TR-06100 Ankara, Turkey. Fax: +90-312-311 4777. E-mail: isaracog@hacettepe.edu.tr

<sup>b</sup> Nagoya City University, Graduate School of Pharmaceutical Sciences, Tanabe-dori 3-1,  
Mizuho-ku, Nagoya 467-8603, Japan

<sup>c</sup> *Present address:* Meijo University, Faculty of Pharmacy, 150 Yagotoyama, Tenpaku-ku,  
Nagoya 468-8503, Japan

\* Author for correspondence and reprints requests

Z. Naturforsch. **58c**, 481–484 (2003); received January 20/March 10, 2003

A new highly oxygenated iridoid glucoside, urphoside B (**1**) was isolated from the *Veronica pectinata* var. *glandulosa* together with seven known iridoid glucosides, aucubin, catalpol, veronicoside, catalposide, verproside, amphicoside and 6-*O*-veratroyl catalpol. The planar as well as the stereo structures of the isolated compounds were determined by means of extensive 1D- and 2D-NMR spectroscopy and confirmed by HR-Mass.

*Key words:* Iridoid Glucoside, Chlorinated Iridoid Glucoside, Urphoside B