

# Triterpenoid Saponins from *Atriplex semibaccata*

Kamel H. Shaker<sup>b</sup>, Katja Dockendorff<sup>a</sup>, and Karlheinz Seifert<sup>a,\*</sup>

<sup>a</sup> University of Bayreuth, Organic Chemistry, NW II, D-95440 Bayreuth, Germany.

Fax: 49-921-55 53 58. E-mail: karlheinz.seifert@uni-bayreuth.de

<sup>b</sup> National Research Centre, Laboratory of Natural Products, Dokki-Cairo, Egypt

\* Author for correspondence and reprint requests

Z. Naturforsch. **58c**, 485–489 (2003); received February 24/March 13, 2003

Four new triterpenoid saponins, 3-*O*-{[ $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)]- $\beta$ -D-galactopyranosyl}-11 $\alpha$ -methoxy-23-hydroxylongispinogenin, 3-*O*-{[ $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)]- $\beta$ -D-galactopyranosyl}-11 $\alpha$ -methoxy-23,29-dihydroxylongispinogenin, 3-*O*-{[ $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)]- $\beta$ -D-galactopyranosyl}-29-hydroxysaikogenin F and 3-*O*-{[ $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)]- $\beta$ -D-galactopyranosyl}-saikogenin F, have been isolated from *Atriplex semibaccata*. The structures were determined primarily by NMR spectroscopy. The assignment of NMR signals was performed by means of <sup>1</sup>H-<sup>1</sup>H COSY, ROESY, HMQC and HMBC experiments.

*Key words:* *Atriplex semibaccata*, Triterpenoid Saponins, Chenopodiaceae