

A New Polyisoprenylated Phloroglucinol Derivative from *Hypericum perforatum* (Clusiaceae)

Naòma Benkiki^a, Zahia Kabouche^b, François Tillequin^{c,*}, Philippe Vérité^d,
Elizabeth Chosson^d, and Elisabeth Seguin^d

^a Département de Chimie, Faculté des Sciences, Université Colonel El-Hadj Lakhdar,
05000 Batna, Algeria

^b Laboratoire d'Obtention de Substances Thérapeutiques (LOST), Faculté des Sciences,
Université Mentouri – Constantine, Campus Chaabet Ersas, 25000 Constantine, Algeria

^c Laboratoire de Pharmacognosie de l'Université René Descartes, U. M. R./C. N. R. S.
N° 8638, Faculté des Sciences Pharmaceutiques et Biologiques, 4 Avenue de l'Observatoire,
F-75006 Paris, France. E-mail: tillequi@pharmacie.univ-paris5.fr

^d Equipe Pharmacomodulation d'antitumoraux d'origine naturelle et synthétique,
Laboratoire de Pharmacognosie, Faculté de Pharmacie, 22, Boulevard Gambetta,
F-76183 Rouen Cedex 1, France

* Author for correspondence and reprint requests

Z. Naturforsch. **58c**, 655–658 (2003); received March 17, 2003

Hyperfoliatin, a new polyisoprenylated phloroglucinol derivative was isolated from the aerial parts of *Hypericum perforatum* (Clusiaceae) collected in Algeria. The structure of hyperfoliatin was elucidated on the basis of its spectral data, mainly MS and multiple-pulse NMR.

Key words: *Hypericum perforatum*, Polyisoprenylated Phloroglucinol