First Evidence for the Presence of Weddellite Crystallites in *Opuntia ficus* indica Parenchyma

Mohamed E. Malainine^{a,b}, Alain Dufresne^b, Danielle Dupeyre^b, Michel R. Vignon^b, and Mostafa Mahrouz^{a*}

* Author for correspondence and reprint requests

- Unité de Chimie Agroalimentaire, Laboratoire de Chimie Organique Appliquée, Département de Chimie, Faculté des Sciences Semlalia, Université Cadi Ayvad. Marrakech, Maroc. Fax: +21244431093. E-mail: mahrouz@ucam.ac.ma
- b Centre de Recherche sur les Macromolécules Végétales, Grenoble, France
- Z. Naturforsch. **58c**, 812–816 (2003); received February 10/April 4, 2003

Calcium oxalate crystallites occur very often in the plants tissues and their role is still poorly known. We report here the experimental protocol leading to the isolation of two forms of calcium oxalate crystallites differing in their hydration level in the parenchymal tissues of Opuntia ficus indica (Miller). Whereas the whewellite crystallites are habitual in all Opuntia species, the weddellite form has never been isolated from these species before, which is probably due to their small size (about $1 \mu m$). We have identified these forms using X-ray diffraction and scanning electron microscopy.

Key words: Opuntia ficus indica, Weddellite, Whewellite