

Phenylethanoid Glycosides from *Phlomis integrifolia* Hub.-Mor.

Iclal Saracoglu^{a,*}, Mehtap Varel^a, Junko Hada^b, Noriyasu Hada^b,
Tadahiro Takeda^b, Ali A. Donmez^c, and Ihsan Calis^a

^a Hacettepe University, Faculty of Pharmacy, Department of Pharmacognosy,
TR-06100 Ankara, Turkey. Fax: +90-3 12-3 11 47 77. E-mail: isaracog@hacettepe.edu.tr

^b Kyoritsu College of Pharmacy, Department of Natural Medicines, Shibakoen 1-5-30,
Minato-ku, Tokyo 105-8512, Japan

^c Hacettepe University, Faculty of Science, Department of Biology, TR-06532, Ankara,
Turkey

* Author for correspondence and reprint requests

Z. Naturforsch. **58c**, 820–825 (2003); received March 27/May 21, 2003

Two new phenylethanoid glycosides integrifoliosides A (**2**) and B (**3**), along with a known phenylethanoid glycoside alyssonoside (**1**) and a flavone glucoside chrysoeriol-7-*O*- β -D-glucopyranoside (**4**) were isolated from the aerial parts of *Phlomis integrifolia*. The structures of the new compounds were identified as 3,4-dihydroxy- β -phenylethoxy-*O*- β -D-apiofuranosyl-(1 \rightarrow 4)- α -L-rhamnopyranosyl-(1 \rightarrow 3)-4-*O*-feruloyl- β -D-glucopyranoside (**2**) and 3-hydroxy-4-methoxy- β -phenylethoxy-*O*- β -D-apiofuranosyl-(1 \rightarrow 4)- α -L-rhamnopyranosyl-(1 \rightarrow 3)-4-*O*-feruloyl- β -D-glucopyranoside (**3**), on the basis of spectroscopic (UV, IR, 1D- and 2D-NMR, and HR-FABMS) methods.

Key words: *Phlomis integrifolia* Hub.-Mor., Phenylethanoid Glycosides, Integrifoliosides A and B