In vitro Anti-inflammatory Effect of Carthamus lanatus L.

Saima Jalil^a, Bozhanka Mikhova^{b*}, Rilka Taskova^c, Maya Mitova^b, Helmut Duddeck^d, Muhammad Igbal Choudhary^a, and Atta-ur-Rahman^a

^a H. E. J. Research Institute of Chemistry, International Center for Sciences, University of

Karachi, Karachi-75270, Pakistan

Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria. Fax: ++3592/8700225. E-mail: bozhana@orgchm.bas.bg

c Institute of Botany, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria
d Institute of Organic Chemistry, University of Hannover, 30167 Hannover, Germany

* Author for correspondence and reprint requests

Z. Naturforsch. **58c**, 830–832 (2003); received May 30/July 17, 2003

The anti-inflammatory activity of four total extracts, their fractions and two main constituents (α-bisabolol β-D-fucopyranoside and luteolin 7-O-glucoside) of Carthamus lanatus L. aerial parts, were assessed *in vitro* by determining the inhibitory effects on induced human

neutrophils. The dichloromethane extract and its water-alcoholic part exhibited the most significant inhibitory effects.

Key words: Carthamus lanatus, Anti-inflammatory Activity, Neutrophils