

Sesquiterpenes and a Phenylpropanoid from *Cordia trichotoma*

Jane Eire S. A. de Menezes^a, Francisca Elane A. Machado^a, Telma Leda G. Lemos^a,
Edilberto R. Silveira^a, Raimundo Braz Filho^b, and Otília Deusdênia L. Pessoa^{a,*}

^a Curso de Pós-Graduação em Química Orgânica, Departamento de Química Orgânica e Inorgânica, Centro de Ciências, Universidade Federal do Ceará, 12.200, Fortaleza – CE, 60021-970, Brazil. E-mail: otilia@dqoi.ufc.br

^b Setor de Química de Produtos Naturais – LCQUI-CCT, Universidade Estadual do Norte Fluminense, Campos – RJ, 28015-620, Brazil

* Author for correspondence and reprint requests

Z. Naturforsch. **59c**, 19–22 (2004); received March 11/June 5, 2003

Two new secondary metabolites, the phenylpropanoid 3-(2',4',5'-trimethoxyphenyl)propanoic acid (**1**) and the sesquiterpene (+)-1 β ,4 β ,6 α -trihydroxyeudesmane (**2**) were isolated from the heartwood of *Cordia trichotoma* Vell., along with the known sesquiterpenes (–)-1 β ,4 β ,7 α -trihydroxyeudesmane (**3**) and (+)-1 β ,4 β ,11-trihydroxyoppositane (**4**). Their structures were elucidated by means of spectroscopic data interpretation, mainly 1D and 2D NMR and mass spectrometry.

Key words: *Cordia trichotoma*, Sesquiterpenes, Phenylpropanoid