Antimicrobial Activity of Fractions and Compounds from Calophyllum brasiliense (Clusiaceae/Guttiferae) Juliana B. Pretto, Valdir Cechinel-Filho, Vânia F. Noldin, Mara R. K. Sartori,

Daniela E. B. Isaias, and Alexandre Bella Cruz*

flowers and fruits) and studied. The following compounds were isolated and tested against

Programa de Mestrado em Ciências Farmacêuticas (PMCF) e Núcleo de Investigações Químico Farmacêuticas (NIQFAR) / CCS, Universidade do Vale do Itajaí (UNIVALI),

88.302-202, Itajaí, SC, Brazil. Fax. +55021473417601. E-mail: abcruz@ccs.univali.br

* Author for correspondence and reprint requests

Z. Naturforsch. **59 c**, 657–662 (2004); received March 17/April 27, 2004

Calophyllum brasiliense (Clusiaceae/Guttiferae) is a native Brazilian medicinal plant traditionally used against several diseases, including infectious pathologies. Crude methanolic extracts (CME) and two fractions, denoted non-polar (soluble in chloroform) and polar (non-soluble in chloroform), were prepared from different parts of the plant (roots, stems, leaves,

pathogenic bacteria and yeasts by determination of the minimal inhibitory concentration (MIC): brasiliensic acid (1), gallic acid (2), epicatechin (3), protocatechuic acid (4), friedelin (5) and 1,5-dihydroxyxanthone (6). The results indicated that all the parts of the plant exhibited antimicrobial activity against Gram-positive bacteria, which are selectively inhibited by components of *C. brasiliense*. No activity was observed against Gram-negative bacteria and yeasts tested. Regarding the isolated compounds, substance 4 showed antimicrobial activity against all the tested microorganisms, whereas compound 6 exhibited antimicrobial activity only against Gram-positive bacteria. The results from the current study confirm and justify the popular use of this plant to treat infectious processes.

Key words: Calophyllum brasiliense, Antimicrobial Activity, Phenolic Compounds