

Chemical Constituents of *Lavatera trimestris* L. – Antioxidant and Antimicrobial Activities

Krystyna Skalicka-Woźniak^{a,b,*}, Eleni Melliou^b, Olga Gortzi^c, Kazimierz Glowinski^a, and Ioanna B. Chinou^b

^a Department of Pharmacognosy with Medicinal Plant Laboratory,
Skubiszewski Medicinal University of Lublin, 1 Chodzki Str., 20-993 Lublin, Poland.
Fax: +4 88 17 41 03 51. E-mail: kskalicka@pharmacognosy.org

^b Department of Pharmacognosy and Chemistry of Natural Products, School of Pharmacy,
University of Athens, University Campus of Zografou,
157 71 Zografou, Athens, Greece

^c Department of Food Technology, Technological Educational Institution (T.E.I.)
of Athens, Egaleo, Athens, Greece

* Author for correspondence and reprint requests

Z. Naturforsch. **62c**, 797–800 (2007); received May 10/June 20, 2007

Nine phenolic compounds, such as *cis*-/*trans*-*p*-coumaric acid, *cis*-/*trans*-*p*-coumaric acid methyl ester, glucose ester of *cis*-/*trans*-*p*-coumaric acid, caffeic acid methyl ester, kaempferol 7-*O*- β -D-glucoside and kaempferol 3-*O*- β -D-glucoside, were isolated from *Lavatera trimestris* flowers by chromatographic techniques and their structures were elucidated by spectral means (NMR). All compounds were tested for their antioxidant activity, while the methanolic extract was tested also for its antimicrobial activity. Also several non-polar constituents have been identified using GC and GC/MS methods. This is the first time that phenolic esters and non-polar constituents were identified in the flowers of *L. trimestris* L.

Key words: *Lavatera trimestris* L., Flavonoids, Phenolic Acids