

# New Naphthoquinone Derivatives from the Ascomycete IBWF79B-90A

Anja Schüffler<sup>a</sup>, Johannes C. Liermann<sup>b</sup>, Heinz Kolshorn<sup>c</sup>, Till Opatz<sup>b</sup>,  
and Heidrun Anke<sup>d,\*</sup>

<sup>a</sup> University of Kaiserslautern, Paul-Ehrlich-Str. 23, D-67663 Kaiserslautern, Germany

<sup>b</sup> Institute of Organic Chemistry, University of Hamburg, Martin-Luther-King-Platz 6,  
D-20146 Hamburg, Germany

<sup>c</sup> Institute of Organic Chemistry, University of Mainz, Duesbergweg 10–14, D-55128  
Mainz, Germany

<sup>d</sup> Institute of Biotechnology and Drug Research, Erwin-Schrödinger-Str. 56, D-67663  
Kaiserslautern, Germany. E-mail: anke@ibwf.de

\* Author for correspondence and reprint requests

Z. Naturforsch. **64c**, 25–31 (2009); received September 23/October 24, 2008

Bioactivity-guided fractionation of extracts from the fungus IBWF79B-90A resulted in the isolation of three known naphthoquinones, herbarin, dehydroherbarin, and *O*-methylherbarin and the azaanthraquinone scorpinone as well as three structurally related derivatives, *O*-phenethylherbarin and herbaridines A and B. All seven compounds exhibited cytotoxic activities against several cell lines.

*Key words:* Naphthoquinone, Herbaridine, Cytotoxic Activity