Xylomexicanins A and B, New ^{14,15}-Mexicanolides from Seeds of the Chinese Mangrove *Xylocarpus granatum*

Li-Ru Shen^a, Mei Dong^b, Dong Guo^c, Bao-Wei Yin^a, Man-Li Zhang^a, Qing-Wen Shi^a, Chang-Hong Huo^{a,*}, Hiromasa Kiyota^{d,*}, Nobuo Suzuki^c, and Bin Cong^{b,*}

- ^a Department of Medicinal Natural Product Chemistry, School of Pharmaceutical Sciences, Hebei Medical University, 361 Zhongshan East Road, Shijiazhuang, 050017, Hebei Province, P. R. China. E-mail: rainbowhuo@hebmu.edu.cn
- b College of Basic Medicine, Hebei Medical University, 050017, Shijiazhuang, P. R. China.
- E-mail: qingwenshi@hebmu.edu.cn

 ^c North China Pharmaceutical Group Corporation, New Drug R&D Co., Ltd., 198
 Huanghe Road, High Technology Industrial Zone, 050035, Shijiazhuang, Hebei Province,
 P. R. China
- Department of Bioscience and Biotechnology for Future Bioindustry, Graduate School of Agricultural Science, Tohoku University, 1–1 Tsutsumidori-Amamiya, Aoba-ku, Sendai, 981–8555, Japan. Fax: +8 12 27 17 87 85. E-mail: kiyota@biochem.tohoku.ac.jp
 Department of Environmental Biochemistry, Graduate School of Medicine, Chiba
- * Authors for correspondence and reprint requests
- Z. Naturforsch. **64c**, 37–42 (2009); received May 13/July 28, 2008

University, Inohana 1–8–1, Chuoku, Chiba, 260–8670, Japan

Two new mexicanolide-type limonoids, named xylomexicanin A (1) and xylomexicanin B (2), were isolated from seeds of the Chinese mangrove *Xylocarpus granatum*. Their structures were elucidated on the basis of spectroscopic methods. Compound 1 exhibited antiproliferative activity against human breast carcinoma cells (KT), while 2 did not show inhibitory effects on eleven human tumour cell lines tested.

Key words: Xylocarpus granatum, Limonoids, Antiproliferative Activity