

New Caloporoside Derivatives and their Inhibition of Fungal Spore Germination

Anja Schüffler^a, Johannes C. Liermann^b, Heinz Kolshorn^c, Till Opatz^b,
and Timm Anke^{d,*}

^a University of Kaiserslautern, Paul-Ehrlich-Str. 23, D-67663 Kaiserslautern, Germany

^b Institute of Organic Chemistry, University of Hamburg, Martin-Luther-King-Platz 6,
D-20146 Hamburg, Germany

^c Institute of Organic Chemistry, University of Mainz, Duesbergweg 10–14,
D-55128 Mainz, Germany

^d Institute of Biotechnology and Drug Research, Erwin-Schrödinger-Str. 56,
D-67663 Kaiserslautern, Germany. E-mail: anke@rhrk.uni-kl.de

* Author for correspondence and reprint requests

Z. Naturforsch. **64c**, 521–525 (2009); received May 19, 2009

In our ongoing screening culture fluid extracts of *Gloeoporus* (*Caloporus*) *dichrous* strain 83065 inhibited the germination of *Magnaporthe grisea* and *Fusarium graminearum* spores. While isolating the active metabolites two new caloporosides, caloporoside G and caloporoside H, in addition to the known caloporoside derivatives F-16438G, caloporoside A, and 2-hydroxy-6-(16-hydroxyheptadecyl)benzoic acid were obtained.

Key words: Caloporoside, Spore Germination Inhibition, *Gloeoporus dichrous*