

Seasonal Variation in the Essential Oil Composition of *Origanum majorana* L. Cultivated in Egypt

Fathy M. Soliman, Miriam F. Yousif*, Soumaya S. Zaghloul, and Mona M. Okba

Pharmacognosy Department, Faculty of Pharmacy, Cairo University, Kasr-El-Ainy St.,
Cairo 11562, Egypt. Fax: 2 02 37 42 68 07. E-mail: miriamyousif@hotmail.com

* Author for correspondence and reprint requests

Z. Naturforsch. **64c**, 611–614 (2009); received March 1/30, 2009

The hydrodistilled essential oil content of *Origanum majorana* L. (Lamiaceae) ranged from 2.5–3% with the maximal value (3%) in spring. Analysis of the oil by GC/MS resulted in the identification of 15, 15, 11, and 14 components in the oils prepared in spring, summer, autumn, and winter, respectively. The composition of the essential oils differed quantitatively and qualitatively according to the time of collection. Thymol (38.4%) and *cis*-sabinene hydrate (25.3%) were the major components in spring plants. Terpinen-4-ol (37.4%, 20.5%, 16.3%) was a major component in the summer, autumn and winter oils, respectively. *cis*-Sabinene hydrate (54.4%) was major in winter plants while terpinolene (43.1%) was the main component in autumn plants. Other components detected in lower amounts in all oil samples were sabinene and *p*-cymene (up to 7.4% and 13.9% in autumn), and α -terpinene (up to 13.3% in summer).

Key words: *Origanum majorana*, Essential Oil, Seasonal Variation