

# Anticancer Activity of Diterpenoids from *Amoora ouangliensis* and *Amoora stellato-squamosa*

Shu-Min Yang<sup>a,\*</sup>, Da-Gang Wu<sup>b</sup>, and Xi-Kui Liu<sup>b,\*</sup>

<sup>a</sup> School of Chemical Biology and Pharmaceutical Sciences, Capital Medical University, Beijing 100069, China. Fax: +86 01 83 91 15 33. E-mail: xiaoyang618@yahoo.com.cn

<sup>b</sup> State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, the Chinese Academy of Sciences, Kunming, Yunnan 650204, China. Fax: +86 87 15 15 02 27. E-mail: liuxikui@mail.kib.ac.cn

\* Authors for correspondence and reprint requests

Z. Naturforsch. **65 c**, 39–42 (2010); received August 7, 2009

A new *ent*-halimane-type diterpene, named 5(10),14-*ent*-halimadien-3 ,13*S*-diol (**1**), was isolated from the bark of *Amoora ouangliensis* and its chemical structure determined on the basis of spectroscopic analysis. Additionally, ten other diterpenoids were obtained from *A. ouangliensis* and *A. stellato-squamosa*. The bioactive experiments of all compounds against AGZY 83-a (human lung cancer cells) and SMMC-7721 (human liver cancer cells) cells were documented.

**Key words:** 5(10),14-*ent*-Halimadien-3 ,13*S*-diol, *Amoora ouangliensis*, *Amoora stellato-squamosa*