

# Steroidal Alkaloids of *Veratrum lobelianum* Bernh. and *Veratrum nigrum* L.

Veselin Christov<sup>a,\*</sup>, Bozhanka Mikhova<sup>a</sup>, Antoaneta Ivanova<sup>a</sup>, Julianna Serly<sup>b</sup>, Joseph Molnar<sup>b</sup>, Dangaa Selenge<sup>c</sup>, Amgalan Solongo<sup>c</sup>, Nadezhda Kostova<sup>a</sup>, Yadamsuren Gerelt-Od<sup>c</sup>, and Dimitar Dimitrov<sup>d</sup>

<sup>a</sup> Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria. E-mail: christov@orgchm.bas.bg

<sup>b</sup> Department of Medical Microbiology and Immunobiology, University of Szeged, H-6720 Szeged, Hungary

<sup>c</sup> Institute of Chemistry and Chemical Technology, Mongolian Academy of Sciences, 211051 Ulaanbaatar-51, Mongolia

<sup>d</sup> National Historical Museum, Bulgarian Academy of Sciences, 1000 Sofia, Bulgaria

\* Author for correspondence and reprint requests

Z. Naturforsch. **65c**, 195–200 (2010); received December 10, 2009/January 22, 2010

Twelve steroidal alkaloids were isolated from four populations of *Veratrum lobelianum* Bernh. and *Veratrum nigrum* L. Full NMR data for veralosinine (**1**), and extensive <sup>1</sup>H NMR data for veralosine (**3**) and teinemine (**5**) are presented here for the first time. (±)-15-*O*-(2-Methylbutyroyl)germine (**10**) is undescribed up to now. The antiproliferative activities of veranigrine, veralosinine, and neogermitrine have shown that they are a perspective for further studies.

**Key words:** *Veratrum lobelianum* Bernh., *Veratrum nigrum* L., Steroidal Alkaloids, Antiproliferative Activity