

New Guaianolide-Type Sesquiterpene Lactones from *Inula verbascifolia*

Abou El-Hamd H. Mohamed^{a,*}, Hosam-Eldin H. Mahmoud^b,
Fathy F. Abdellatif^c, Yousif S. Mohamed^c, the late Ahmed A. Ahmed^c,
and Shinji Ohta^d

^a Chemistry Department, Faculty of Science, Aswan University, Aswan, Egypt.
Fax: +973480450. E-mail: abouhassan68@yahoo.com

^b Medical Chemistry Department, Faculty of Medicine, Jazan University,
Jazan, Saudi Arabia

^c Chemistry Department, Faculty of Science, El-Minia University, El-Minia, Egypt

^d Nagahama Institute of Bio-Science and Technology, Nagahama, Shiga 526-0829, Japan

* Author for correspondence and reprint requests

Z. Naturforsch. **68c**, 175–180 (2013); received December 12, 2012/May 8, 2013

The aerial parts of *Inula verbascifolia* afforded two new guaianolide-type sesquiterpene lactones. Their structures were determined by spectroscopic methods (IR, MS, ¹H NMR, ¹³C NMR, DEPT, ¹H-¹H COSY, HMQC, and HMBC).

Key words: Asteraceae, *Inula*, Sesquiterpene Lactones