

Exact Periodic Solitary Wave and Double Periodic Wave Solutions for the (2+1)-Dimensional Korteweg-de Vries Equation*

Changfu Liu^a and Zhengde Dai^b

^a Department of Mathematics and Physics, Wenshan University, Wenshan 663000, China

^b School of Mathematics and Statistics, Yunnan University, Kunming 650091, China

Reprint requests to C. .L.; E-mail: chfuliu@yahoo.com.cn

Z. Naturforsch. **64a**, 609 – 614 (2009); received October 27, 2008 / revised December 29, 2008

A new technique, the extended ansatz function method, is proposed to seek periodic solitary wave solutions of integrable systems. Exact periodic solitary wave solutions for the (2+1)-dimensional Korteweg-de Vries (KdV) equation are obtained by using this technique. By using the trial function method, Jacobi elliptic function double periodic solutions are also constructed for this equation. This result shows that there exist periodic solitary waves in the different directions for the (2+1)-dimensional KdV equation.

Key words: Two-Dimensional KdV Equation; Extended Ansatz Function Method; Periodic Solitary Wave Solution; Double Periodic Solution.

1. Introduction

Boiti et al. derived the (2+1)-dimensional KdV equation [1]

$$u_t + 3(uv)_x + u_{xxx} = 0, \quad u_x - v_y = 0 \quad (1)$$

by using the idea of the weak Lax pair. (1) is also called Boiti-Leon-Manna-Pempinelli equation [2]. For $v = u$ and $y = x$, (1) reads as the ubiquitous KdV equation in dimensionless variables

$$u_t + 6uu_x + u_{xxx} = 0. \quad (2)$$

Thus, (1) is a generalisation of the KdV equation. The rich dromion structures and localized structures are revealed by Lou et al. for (1) [3,4] and Wazwaz [5]. In the recent years, many authors have applied the Auto-Bäcklund transformation method [6] and the F-expansion method [7] and found abundant exact solutions for many nonlinear partial differential equations. But these obtained structures do not include some periodic solitary wave solutions which is a new type of soliton solutions.

Recently, Dai et al. introduced a new technique, the extended homoclinic test technique, which is used

for seeking periodic solitary wave solutions of integrable equations and obtained periodic solitary wave solutions for integrable equations [8–12]. In this paper, we extend the technique mentioned above, introduce a new ansatz function $f(x, y, t) = b_1 e^{(Px+Qy+Wt)} + b_2 \cos(Kx + Ly + Vt) + b_3 e^{-(Px+Qy+Wt)}$, which is different from the function $f(x, y, t) = 1 + a_1 p(x, t) + a_2 q(y, t) + Ap(x, t)q(y, t)$ in [4], and consider (1). Then, we use the trial function method [13] to look for the double periodic wave solutions of (1). Exact periodic solitary wave and double periodic wave solutions are obtained for (1). This result shows that there exists periodic solitary waves in the different directions for (1). To our knowledge, some periodic solitary wave and double periodic wave solutions of the (2+1)-dimensional KdV equation are new solutions up to now.

The rest of this paper is organized as follows: in Section 2, a procedure for solving the (2+1)-dimensional KdV equation is given. Finally, some conclusions follow.

2. Exact Periodic Solitary Wave and Double Periodic Wave Solutions for the (2+1)-dimensional KdV Equation

Method 1.

We transform (1) into the bilinear form

$$D_y(D_t + D_x^3)f \cdot f = 0 \quad (3)$$

* Project supported by Chinese Natural Science Foundation (No. 10361007, 10661002), Yunnan Natural Science Foundation (No. 2004A0001M) and Yunnan Educational Science Foundation (No. 06Y041A).

by the transformation

$$u = 2 \ln(fx, y, t)_{xy}, v = 2 \ln(f(x, y, t))_{xx}, \quad (4)$$

where the bilinear operator $D_x^m D_t^n$ is defined as

$$D_x^m D_t^n a \cdot b = \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^m \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^n \cdot a(x, t) b(x', t') \Big|_{x=x', t=t'}. \quad (5)$$

Then (3) can be rewritten as

$$2(f_{yt}f - f_y f_t + f_{xxx}f + 3f_{xx}f_{xy} - 3f_{xy}f_x - f_{xx}f_y) = 0. \quad (6)$$

We express the function f in the form

$$f(x, y, t) = b_1 e^{(Px+Qy+Wt)} + b_2 \cos(Kx+Ly+Vt) + b_3 e^{-(Px+Qy+Wt)}. \quad (7)$$

Substituting (7) into (6), we obtain an algebraic equation

for $e^{j(Px+Qy+Wt)}$, $\sin(Kx+Ly+Vt)$, and $\cos(Kx+Ly+Vt)$. Equating all therein coefficients to zero ($j = -1, 0, 1$) yields a set of algebraic equations for $b_1, b_2, b_3, P, Q, W, K, L$ and V :

$$\begin{aligned} 16b_1 Q b_3 P^3 + 4b_1 Q b_3 W - b_2^2 LV + 4b_2^2 LK^3 &= 0, \\ -b_2 L b_3 P^3 + b_3 Q b_2 K^3 - b_3 Q b_2 V - b_2 L b_3 W \\ + 3b_3 P b_2 LK^2 - 3b_2 K b_3 P^2 Q &= 0, \\ -b_2 b_3 QW - b_3 b_2 VL + b_3 P^3 Q b_2 + b_2 LK^3 b_3 \\ - 3b_2 K^2 b_3 PQ - 3b_3 P^2 b_2 LK &= 0, \\ -b_1 Q b_2 K^3 + b_2 L b_1 P^3 + b_2 L b_1 W + b_1 Q b_2 V \\ - 3b_1 P b_2 LK^2 + 3b_2 K b_1 P^2 Q &= 0, \\ b_1 P^3 Q b_2 + b_2 LK^3 b_1 + b_2 b_1 QW - b_1 b_2 VL \\ - 3b_2 K^2 b_1 PQ - 3b_1 P^2 b_2 LK &= 0. \end{aligned} \quad (8)$$

Solving this system of algebraic equations with the aid of Maple, we can obtain the following results.

Case 1.

$$\begin{aligned} P &= -\frac{1}{4} \frac{KLb_2^2}{b_1 Q b_3}, \quad W = -\frac{1}{64} \frac{b_2^2 LK^3 (48b_1^2 Q^2 b_3^2 - L^2 b_2^4)}{b_1^3 Q^3 b_3^3}, \quad V = \frac{1}{16} \frac{K^3 (16b_1^2 Q^2 b_3^2 - 3L^2 b_2^4)}{b_1^2 Q^2 b_3^2}, \\ K &= K, \quad L = L, \quad Q = Q, \quad b_1 = b_1, \quad b_2 = b_2, \quad b_3 = b_3, \end{aligned} \quad (9)$$

where K, L, Q, b_1, b_2, b_3 are free parameters. Setting $b_3 = \pm b_1$, we have

$$\begin{aligned} f_1(x, y, t) &= 2b_1 \cosh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt), \\ f_2(x, y, t) &= 2b_1 \sinh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt). \end{aligned} \quad (10)$$

Substituting (10) into (4) yields periodic soliton solutions of (1) as follows:

$$\begin{aligned} u_1(x, y, t) &= 2 \left\{ \frac{4b_1^2 PQ - b_2^2 LK + 2b_1 b_2 (PQ - KL) \cosh(Px + Qy + Wt) \cos(Kx + Ly + Vt)}{(2b_1 \cosh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt))^2} \right. \\ &\quad \left. + \frac{2b_1 b_2 (PL + KQ) \sinh(Px + Qy + Wt) \sin(Kx + Ly + Vt)}{(2b_1 \cosh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt))^2} \right\}, \\ v_1(x, y, t) &= 2 \left\{ \frac{4b_1^2 P^2 - b_2^2 K^2 + 2b_1 b_2 (P^2 - K^2) \cosh(Px + Qy + Wt) \cos(Kx + Ly + Vt)}{(2b_1 \cosh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt))^2} \right. \\ &\quad \left. + \frac{4b_1 b_2 PK \sinh(Px + Qy + Wt) \sin(Kx + Ly + Vt)}{(2b_1 \cosh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt))^2} \right\}, \end{aligned} \quad (11)$$

and

$$\begin{aligned} u_2(x, y, t) &= 2 \left\{ \frac{-4b_1^2 PQ - b_2^2 LK + 2b_1 b_2 (PQ - KL) \sinh(Px + Qy + Wt) \cos(Kx + Ly + Vt)}{(2b_1 \sinh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt))^2} \right. \\ &\quad \left. + \frac{2b_1 b_2 (PL + KQ) \cosh(Px + Qy + Wt) \sin(Kx + Ly + Vt)}{(2b_1 \sinh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt))^2} \right\}, \end{aligned}$$

$$v_2(x, y, t) = 2 \left\{ \frac{-4b_1^2 P^2 - b_2^2 K^2 + 2b_1 b_2 (P^2 - K^2) \sinh(Px + Qy + Wt) \cos(Kx + Ly + Vt)}{(2b_1 \sinh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt))^2} + \frac{4b_1 b_2 P K \cosh(Px + Qy + Wt) \sin(Kx + Ly + Vt)}{(2b_1 \sinh(Px + Qy + Wt) + b_2 \cos(Kx + Ly + Vt))^2} \right\}, \quad (12)$$

where all parameters are defined by (9).

Especially, when $L = 0$, (11) and (12) become

$$\begin{aligned} u_1(x, y, t) &= \frac{4b_1 b_2 K Q \sin(Kx + K^3 t) \sinh(Qy)}{(2b_1 \cosh(Qy) + b_2 \cos(Kx + K^3 t))^2}, \\ v_1(x, y, t) &= \frac{-2b_2 K^2 (2b_1 \cos(Kx + K^3 t) \cosh(Qy) + b_2)}{(2b_1 \cosh(Qy) + b_2 \cos(Kx + K^3 t))^2}, \end{aligned} \quad (13)$$

and

$$\begin{aligned} u_2(x, y, t) &= \frac{4b_1 b_2 K Q \sin(Kx + K^3 t) \cosh(Qy)}{(2b_1 \sinh(Qy) + b_2 \cos(Kx + K^3 t))^2}, \\ v_2(x, y, t) &= \frac{-2b_2 K^2 (2b_1 \cos(Kx + K^3 t) \sinh(Qy) + b_2)}{(2b_1 \sinh(Qy) + b_2 \cos(Kx + K^3 t))^2}, \end{aligned} \quad (14)$$

where b_1 , b_2 , K and Q are free parameters.

Case 2.

$$\begin{aligned} L &= L, \quad P = P, \quad W = -P^3, \quad V = 0, \quad Q = 0, \\ K &= 0, \quad b_1 = b_1, \quad b_2 = b_2, \quad b_3 = b_3. \end{aligned} \quad (15)$$

Similar as previous case, we set $b_3 = \pm b_1$ and obtain the following periodic soliton solutions of (1):

$$\begin{aligned} u_1(x, y, t) &= \frac{4b_1 b_2 P L \sinh(Px - P^3 t) \sin(Ly)}{(2b_1 \cosh(Px - P^3 t) + b_2 \cos(Ly))^2}, \\ v_1(x, y, t) &= \frac{4b_1 P^2 (2b_1 + b_2 \cosh(Px - P^3 t) \cos(Ly))}{(2b_1 \cosh(Px - P^3 t) + b_2 \cos(Ly))^2}, \end{aligned} \quad (16)$$

and

$$\begin{aligned} u_2(x, y, t) &= \frac{4b_1 b_2 P L \cosh(Px - P^3 t) \sin(Ly)}{(2b_1 \sinh(Px - P^3 t) + b_2 \cos(Ly))^2}, \\ v_2(x, y, t) &= \frac{-4b_1 P^2 (2b_1 - b_2 \cosh(Px - P^3 t) \cos(Ly))}{(2b_1 \sinh(Px - P^3 t) + b_2 \cos(Ly))^2}, \end{aligned} \quad (17)$$

where b_1 , b_2 , P and L are free parameters.




Fig. 1. Periodic soliton solution of (13) with $b_1 = 1$, $b_2 = 2$, $K = 2$, $Q = 2$; (a): $t = 1$; (b): $y = 1$, $t = 1$; (c): $x = 0$, $t = 1$.

Case 3.

$$\begin{aligned} K &= \sqrt{3}P, \quad w = 8P^3, \quad L = L, \quad P = P, \\ Q &= -\frac{\sqrt{3}L}{4b_3 b_1} b_2^2, \quad v = 0, \quad b_1 = b_1, \\ b_2 &= b_2, \quad b_3 = b_3. \end{aligned} \quad (18)$$




Fig. 2. Periodic soliton solution of (13) with $b_1 = 1$, $b_2 = 2$, $K = 2$, $Q = 2$; (a): $t = 1$; (b): $y = 1$, $t = 1$; (c): $x = 0$, $t = 1$.

Setting $b_3 = \pm b_1 = b_2$, we get

$$\begin{aligned} f_1(x,y,t) &= b_2 \left[2 \cosh \left(Px - \frac{1}{4} L \sqrt{3} y + 8P^3 t \right) \right. \\ &\quad \left. + \cos(\sqrt{3} Px + Ly) \right], \\ f_2(x,y,t) &= b_2 \left[2 \sinh \left(Px + \frac{1}{4} L \sqrt{3} y + 8P^3 t \right) \right. \\ &\quad \left. + \cos(\sqrt{3} Px + Ly) \right]. \end{aligned} \quad (19)$$




Fig. 3. Periodic soliton solution of (17) with $b_1 = 1$, $b_2 = 2$, $P = 2$, $L = 2$, and $t = 0$.




Fig. 4. Periodic soliton solution of (20) with $P = 2$, $L = 2$, and $t = 0$.

Substituting (19) into (4) yields periodic soliton solutions of (1) as follows:

$$\begin{aligned} u_1(x, y, t) &= -PL \left\{ \frac{4\sqrt{3} + 5\sqrt{3} \cosh(Px - \frac{1}{4}L\sqrt{3}y + 8P^3t) \cos(\sqrt{3}Px + Ly)}{(2 \cosh(Px - \frac{1}{4}L\sqrt{3}y + 8P^3t) + \cos(\sqrt{3}Px + Ly))^2} \right. \\ &\quad \left. - \frac{\sinh(Px - \frac{1}{4}L\sqrt{3}y + 8P^3t) \sin(\sqrt{3}Px + Ly)}{(2 \cosh(Px - \frac{1}{4}L\sqrt{3}y + 8P^3t) + \cos(\sqrt{3}Px + Ly))^2} \right\}, \\ v_1(x, y, t) &= 2P^2 \left\{ \frac{1 - 4 \cosh(Px - \frac{1}{4}L\sqrt{3}y + 8P^3t) \cos(\sqrt{3}Px + Ly)}{(2 \cosh(Px - \frac{1}{4}L\sqrt{3}y + 8P^3t) + \cos(\sqrt{3}Px + Ly))^2} \right. \\ &\quad \left. + \frac{4\sqrt{3} \sinh(Px - \frac{1}{4}L\sqrt{3}y + 8P^3t) \sin(\sqrt{3}Px + Ly)}{(2 \cosh(Px - \frac{1}{4}L\sqrt{3}y + 8P^3t) + \cos(\sqrt{3}Px + Ly))^2} \right\}, \end{aligned} \quad (20)$$

and

$$\begin{aligned} u_2(x, y, t) &= -PL \left\{ \frac{4\sqrt{3} + 3\sqrt{3} \sinh(Px + \frac{1}{4}L\sqrt{3}y + 8P^3t) \cos(\sqrt{3}Px + Ly)}{(2 \sinh(Px + \frac{1}{4}L\sqrt{3}y + 8P^3t) + \cos(\sqrt{3}Px + Ly))^2} \right. \\ &\quad \left. - \frac{7 \cosh(Px + \frac{1}{4}L\sqrt{3}y + 8P^3t) \sin(\sqrt{3}Px + Ly)}{(2 \sinh(Px + \frac{1}{4}L\sqrt{3}y + 8P^3t) + \cos(\sqrt{3}Px + Ly))^2} \right\}, \\ v_2(x, y, t) &= 2P^2 \left\{ \frac{-7 - 4 \sinh(Px + \frac{1}{4}L\sqrt{3}y + 8P^3t) \cos(\sqrt{3}Px + Ly)}{(2 \sinh(Px + \frac{1}{4}L\sqrt{3}y + 8P^3t) + \cos(\sqrt{3}Px + Ly))^2} \right. \\ &\quad \left. + \frac{4\sqrt{3} \cosh(Px + \frac{1}{4}L\sqrt{3}y + 8P^3t) \sin(\sqrt{3}Px + Ly)}{(2 \sinh(Px + \frac{1}{4}L\sqrt{3}y + 8P^3t) + \cos(\sqrt{3}Px + Ly))^2} \right\}. \end{aligned} \quad (21)$$

As shown in Figure 4, the solutions $u_1(x, y, t)$ and $v_1(x, y, t)$ in (20) (similarly, other periodic soliton solutions) are obviously two periodic solitary waves which are periodic waves along the X -direction: $X = \sqrt{3}Px + Ly$ with period 2π , and are solitary waves along Y -direction: $Y = Px - \frac{1}{4}L\sqrt{3}y + 8P^3t$. We note that here waves are two left-propagation waves along Y -direction: $Y = Px - \frac{1}{4}L\sqrt{3}y + 8P^3t$. This is a interesting phenomenon to the evolution of shallow water waves. One problem, which we will study, is whether there exist a similar phenomenon to other equations of shallow water waves or not.

Method 2.

Suppose that (1) has the Jacobi elliptic function double periodic solution as follows:

$$u = \frac{A + B \operatorname{cn}^2(\xi, m)}{C + \operatorname{cn}^2(\xi, m)}, \quad v = \frac{E + F \operatorname{cn}^2(\xi, m)}{G + \operatorname{cn}^2(\xi, m)}, \quad (22)$$

where $\xi = Px + Qy + Wt$ and $A, B, C, E, F, G, P, Q, W$ are constants to be determined, m is the modulus of the Jacobi elliptic function ($0 \leq m \leq 1$). With the aid of Maple, we substitute (22) into (1) and balance the coefficients of $\operatorname{cn}(\xi, m)^j$ ($j = 0, 2, 4, 6, 8$) to yield a set of algebraic equations. By solving this set of equations, we can determine A, B, C, E, F, G, P, Q and W . Then, from (1) and (22) we obtain the double periodic function solutions of (1), which read

$$u_1 = -\frac{Q(3PF - 4P^3 + W + 8m^2P^3) \operatorname{cn}^2(Px + Qy + Wt, m) - 3EP}{P^2 \operatorname{cn}^2(Px + Qy + Wt, m)}, \quad v_1 = \frac{E + F \operatorname{cn}^2(Px + Qy + Wt, m)}{\operatorname{cn}^2(Px + Qy + Wt, m)}, \quad (23)$$

$$u_2 = \frac{Bcn^2(Px + Qy + Wt, m) - B + 2QP}{-1 + cn^2(Px + Qy + Wt, m)},$$

$$v_2 = \frac{1}{3} \left[\frac{(4P^3Q - WQ - 3P^2B + 4P^3m^2Q)cn^2(Px + Qy + Wt, m)}{QP(-1 + cn^2(Px + Qy + Wt, m))} + \frac{2P^3Q - 4P^3m^2Q + WQ + 3P^2B}{QP(-1 + cn^2(Px + Qy + Wt, m))} \right], \quad (24)$$

and

$$u_3 = \frac{m^2Bcn^2(Px + Qy + Wt, m) + B - Bm^2 - 2Pm^2Q + 2PQ}{1 - m^2 + m^2cn^2(Px + Qy + Wt, m)},$$

$$v_3 = \frac{1}{3} \left[\frac{(4m^4P^3Q - m^2WQ - 3m^2P^2B - 8P^3m^2Q)cn^2(Px + Qy + Wt, m)}{QP(1 - m^2 + m^2cn^2(Px + Qy + Wt, m))} + \frac{WQm^2 + 6P^3m^2Q - 2P^3Q - 3P^2B + 3P^2Bm^2 - 4P^3m^4Q - WQ}{QP(1 - m^2 + m^2cn^2(Px + Qy + Wt, m))} \right], \quad (25)$$

where B, E, F, P, Q and W are free parameters.

3. Conclusion

In this paper, the (2+1)-dimensional KdV equation is investigated by the extended ansatz function method and trial function method. Some periodic solitary wave solutions and double periodic solutions are obtained for this equation. The result shows that there exists periodic solitary waves in the different directions for

(2+1)-dimensional KdV equation. These methods are effective for seeking periodic solitary wave solutions and double periodic solutions of some integrable equations.

Acknowledgement

The authors would like to thank the referees for valuable suggestion and help.

- [1] M. Boiti, J. Leon, and F. Pempinelli, *Inverse Probl.* **2**, 271 (1985).
- [2] D. S. Wang and H. Li, *J. Math. Anal. Appl.* **343**, 273 (2008).
- [3] S. Y. Lou, *J. Phys. A: Math. Gen.* **28**, 7227 (1995).
- [4] S. Y. Lou and H. Y. Ruan, *J. Phys. A: Math. Gen.* **34**, 305 (2001).
- [5] A. M. Wazwaz, *Appl. Math. Comput.* **204**, 20 (2008).
- [6] D. S. Wang and H. Q. Zhang, *Intern. J. Modern Phys. C* **16**, 393 (2005).
- [7] D. S. Wang and H. Q. Zhang, *Chaos, Solitons and Fractals* **25**, 601 (2005).
- [8] Z. D. Dai, Z. T. Li, Z. J. Liu, and D. L. Li, *Phys. Lett. A* **372**, 3010 (2008).
- [9] Z. D. Dai, Z. J. Liu, and D. L. Li, *Chin. Phys. Lett. A* **25**, 1531 (2008).
- [10] Z. D. Dai, S. L. Li, Q. Y. Dai, and J. Huang, *Chaos, Solitons and Fractals* **34**, 1148 (2007).
- [11] Z. D. Dai, M. R. Jiang, Q. Y. Dai, and S. L. Li, *Chin. Phys. Lett.* **23**, 1065 (2006).
- [12] Z. D. Dai, J. Huang, M. R. Jiang, and S. H. Wang, *Chaos Solitons and Fractals* **26**, 1189 (2005).
- [13] F. D. Xie and Z. Y. Yan, *Chaos, Solitons and Fractals* **36**, 1108 (2008).