

Chemical and Biological Investigation of *Araucaria heterophylla* Salisb. Resin

Essam Abdel-Sattar^{a,*}, Azza R. Abdel Monem^b, Shahira M. Ezzat^b,
Ali M. El-Halawany^{b,c}, and Samar M. Mouneir^d

^a Department of Natural Products, Faculty of Pharmacy, King Abdulaziz University, Jeddah 21589, Kingdom of Saudi Arabia. Fax: +9 66-2-6 95 16 96.
E-mail: abdelstarr@yahoo.com

^b Department of Pharmacognosy, Faculty of Pharmacy, Cairo University, Cairo 11562, Egypt

^c Department of Metabolic Engineering, Institute of Natural Medicine, University of Toyama, Sugitani 2630, Japan

^d Department of Pharmacology, Faculty of Veterinary Medicine, Cairo University, Cairo 12211, Egypt

* Author for correspondence and reprint requests

Z. Naturforsch. **64c**, 819–823 (2009); received May 1/June 8, 2009

Three labdane diterpenes, namely labda-8(17),14-diene, 13-epicupressic acid, and 13-*O*-acetyl-13-epicupressic acid, were isolated from the resin collected from stem exudates of *Araucaria heterophylla* Salisb. (Araucariaceae). The isolated compounds were identified using different spectroscopic methods (¹H NMR, ¹³C NMR, HMQC, HMBC and COSY). The resin extract showed antiulcerogenic activity against ethanol-induced stomach ulcers in Sprague Dawley rats using ranitidine as standard. In addition, the resin and the isolated compounds showed variable cytotoxic activities against breast (MCF7) and colon (HCT116) cancer cell lines.

Key words: *Araucaria heterophylla*, Labdane Diterpenes, Antiulcerogenic and Cytotoxic Activities

Introduction

Araucariaceae (Coniferae) is a small family; it comprises two genera, *Araucaria* and *Agathis*, and 38 species of trees (Trease and Evans, 1978). *Araucaria heterophylla* Salisb. Franco (syn. *A. excelsa*) is a popular columnar tree used as Christmas tree (Schans *et al.*, 2004). Diterpenes (Caputo and Mangoni, 1974; Caputo *et al.*, 1974a, b; Cox *et al.*, 2007) and lignans (Ohashi *et al.*, 1992; Fonseca *et al.*, 1978, 1979) were isolated from resin exudates of different *Araucaria* species including *A. heterophylla* Salisb. (Caputo *et al.*, 1972). The resins of this genus were reported to possess cytotoxic and gastroprotective activities (Lee and Cheng, 2001; Schmeda-Hirschmann *et al.*, 2005a, b). In the present study, the resin exudates from stems of *A. heterophylla* Salisb. growing in Egypt as ornamental plant were subjected to chemical and biological investigations.

Material and Methods

Plant material

The resin exudates from the stems of *A. heterophylla* Salisb. were collected from El-Muntaza Palace Garden, Alexandria, Egypt in August 2007. The plant was identified by Mrs. T. Labib, taxonomist of Orman Botanical Garden, Giza, Egypt. A specimen was deposited in the herbarium of Orman Botanical Garden, Giza, Egypt.

General

¹H NMR (400 MHz), ¹³C NMR (100 MHz), and 2D NMR spectra were measured on a JHA-LAA 400 WB-FT Jeol spectrophotometer. TLC was performed on precoated silica gel plates using the solvent system *n*-hexane/EtOAc (4:1); the chromatograms were visualized by spraying with anisaldehyde/sulfuric acid spray reagent followed by heating at 110 °C for 5 min.

Extraction and fractionation

The resin of *A. heterophylla* Salisb. (20 g) was extracted with CHCl_3 , then with $\text{CHCl}_3/\text{MeOH}$ (1:1, 3×100 ml for each solvent). The combined extracts were evaporated under reduced pressure ($\leq 60^\circ\text{C}$) to give 15 g of a sticky yellow residue.

The residue (10 g) was chromatographed on a silica gel G 60 column (7×20 cm, VLC) eluted with *n*-hexane, *n*-hexane/ CHCl_3 and $\text{CHCl}_3/\text{MeOH}$ mixtures. Fractions of 200 ml each were collected and monitored by TLC. Similar fractions were pooled together. Fractions 8–10 (2.8 g), eluted with *n*-hexane/ CHCl_3 3:7 to 1:9, showed three major spots. The combined fractions were rechromatographed on successive silica gel columns using *n*-hexane/ethyl acetate mixtures to afford compounds **1** (12 mg), **2** (20 mg), and **3** (50 mg). The three compounds were isolated as yellow resins.

Antiulcerogenic effect

Sprague Dawley rats (150–200 g body wt.) were kept under standard hygienic conditions before use. They were fed and watered *ad libitum*. Rats were randomly divided into 4 equal groups of 5 rats each. Animals were starved for 48 h before use to ensure an empty stomach and were kept in cages with raised floors of wide wire mesh to prevent corporophagy (Garg *et al.*, 1993). To prevent excessive dehydration during the fasting period, rats were supplied with 0.2% sucrose (BHD) which was removed 1 h before the experiment (Glavin and Mikhaeil, 1976). The control group was given an equal volume of distilled water instead of the resin extract, but received ethanol in the same dose and route (orally) as the other groups. In addition, the second group was given ranitidine as a standard in a dose of 100 mg/kg body wt. orally by a stomach tube. The third and fourth groups were orally given the resin extract in doses of 100 and 200 mg/kg body wt., respectively. On the first day, rats of the third and fourth groups were orally given two doses of the resin extract with 6 h apart; a third dose was given on the second day 1.5 h before oral administration of ethanol (Merck), 50% (v/v in distilled water), in a dose of 10 ml/kg. 1 h after ethanol administration, all rats were euthanized by an overdose of chloroform, their abdomens were opened and their stomachs rapidly removed, opened along their greater curvature and gently raised under

running water. Lesions in the glandular part of the stomach were measured under an illuminated magnifying microscope (10 \times). Long lesions were counted and measured along their greater length. Petechial lesions were taken as 1 mm ulcer (Cho and Ogle, 1978). The sum of the total length of long ulcers and petechial lesions in each group of rats was divided by its number to calculate the ulcer index (in mm). The curative ratio was determined by the following formula:

$$\text{curative ratio} = (\text{control ulcer index} - \text{test ulcer index}) / (\text{control ulcer index}) \cdot 100.$$

Statistical analysis was carried out by using one-way ANOVA followed by Dubcan test and SPSS Version 14.0. The difference of means at $P < 0.05$ is considered significant (Snedecor and Cochran, 1969).

Cytotoxic activity

The resin extract of *A. heterophylla* Salisb. as well as the isolated compounds were tested *in vitro* for their cytotoxic activity using the sulphorhodamine B assay (Skehan *et al.*, 1990) against colon (HCT116) and breast (MCF7) human cancer cell lines. The results are shown in Table III.

Results and Discussion

Three labdane diterpenes, **1–3** (Fig. 1), were isolated from the resin of *A. heterophylla* Salisb. by chromatographic fractionation on series of silica gel columns. ^1H NMR and ^{13}C NMR spectra showed a labdane diterpene skeleton with two double bonds at C-8(17) and C-14 (Wang *et al.*, 2008).

The ^1H NMR spectrum of **1** (Table I) showed three methyl singlets at δ_{H} 0.64 (CH_3 -18), 1.22 (CH_3 -19), and 1.23 (CH_3 -20). A methyl doublet




Fig. 1. Chemical structures of the compounds isolated from *A. heterophylla*.

Table I. ^1H NMR spectral data of compounds **1–3** (δ_{H} in ppm).

H	1	2	3
14	5.85 (1H, <i>m</i>)	5.88 (1H, <i>dd</i> , $J = 17.2, 10.8$ Hz)	5.81 (1H, <i>dd</i> , $J = 17.2, 10.8$ Hz)
15 α	5.10 (1H, <i>d</i> , $J = 14$ Hz)	5.02 (1H, <i>d</i> , $J = 10.8$ Hz)	4.98 (1H, <i>d</i> , $J = 10.8$ Hz)
β	4.93 (1H, <i>d</i> , $J = 14$ Hz)	5.03 (1H, <i>d</i> , $J = 10.8$ Hz)	5.13 (1H, <i>d</i> , $J = 10.8$ Hz)
16	0.82 (3H, <i>d</i> , $J = 8$ Hz)	1.45 (3H, <i>s</i>)	1.22 (3H, <i>s</i>)
17 α	4.55 (1H, <i>s</i>)	4.43 (1H, <i>br s</i>)	4.48 (1H, <i>s</i>)
β	4.83 (1H, <i>s</i>)	4.76 (1H, <i>br s</i>)	4.79 (1H, <i>s</i>)
18	0.64 (3H, <i>s</i>)	0.52 (3H, <i>s</i>)	0.54 (3H, <i>s</i>)
19	1.22 (3H, <i>s</i>)	1.16 (3H, <i>s</i>)	1.18 (3H, <i>s</i>)
20	1.23 (3H, <i>s</i>)	–	–
COCH ₃	–	1.94 (3H, <i>s</i>)	–

at δ_{H} 0.82 (3H, $J = 8$ Hz) was assigned to CH₃-16. Two doublets at δ_{H} 4.93 and 5.10 (each 1H, $J = 14$ Hz) were assigned to the two olefinic protons at C-15, while a multiplet at δ_{H} 5.85 (1H, *m*) was assigned to the olefinic proton at C-14. Inspection of the ^{13}C NMR spectrum of **1** (Table II) indicated the assignment of signals at δ_{C} 111.9 and 146.6 to C-15 and C-14, respectively. The two exomethylene protons (H-17) appeared at δ_{H} 4.55 (H α) and 4.83 (H β), corresponding to a carbon signal at δ_{C} 107.4 (C-17), while the signal at 150.0 was assigned to C-8. Compound **1** was identified as labda-8(17)14-diene by comparison of its ^1H and ^{13}C NMR spectral data with those of closely related labdane diterpenes (Su *et al.*, 1994; Wang *et al.*, 2008). This compound was only identified in the essential oil of *Cistus monspeliensis* by GC-MS (Angelopoulou *et al.*, 2002) and for the first time in the family Araucariaceae.

The ^1H NMR spectrum of **3** showed three methyl singlets at δ_{H} 1.22, 0.54 and 1.18 assigned to C-16, C-18 and C-19, respectively. The two singlets resonating at δ_{H} 4.48 and 4.79 (each 1H) were assigned to α - and β -exomethylene protons at C-17 and confirmed by HMQC relation to the olefinic carbon signal at δ_{C} 106.6 (C-17) and the presence of a quaternary carbon signal at δ_{C} 147.8 (C-8). The two doublets at δ_{H} 4.98 and 5.13 (each 1H, d , $J = 10.8$ Hz) were assigned to the two olefinic protons at C-15, while the doublet of doublet at δ_{H} 5.81 (1H, $J = 17.2$ and 10.8 Hz) was assigned to the olefinic proton at C-14. This was further confirmed by the correlation to olefinic carbon signals at δ_{C} 111.6 (C-15) and 144.9 (C-14) in the HMQC spectrum. The presence of a tertiary OH group at C-13 was indicated by the quaternary signal at δ_{C} 73.7, which showed a long-range correlation to the downfield shifted proton of CH₃-16 at δ_{H} 1.22. A quaternary carbon sig-

nal at δ_{C} 183.3 was assigned to COOH (C-19) at C-4. Complete assignments of proton and carbon signals were confirmed by the analysis of ^1H - ^1H COSY, HMQC and HMQC spectra. From the previous data and by comparison with reported spectral data (Caputo *et al.*, 1974b; Su *et al.*, 1994; Wang *et al.*, 2008), compound **3** was identified as 13-epicupressic acid. This is the first report on the isolation of 13-epicupressic acid from *A. heterophylla* Salisb.

Spectral data of **2** were heavily similar to those of **3**, except for the presence of an additional downfield shifted methyl signal at δ_{H} 1.94 (3H, *s*, OCOCH₃) and a quaternary carbon signal at

Table II. ^{13}C NMR spectral data of compounds **1–3** (δ_{C} in ppm).

C	1	2	3
1	39.0	38.9	39.0
2	18.8	19.8	19.8
3	42.5	37.8	37.9
4	32.9	44.1	44.1
5	54.8	56.3	56.3
6	23.4	26.0	25.9
7	37.6	38.6	38.7
8	150.0	147.8	147.8
9	57.1	56.5	56.5
10	40.8	39.2	40.6
11	20.9	17.5	17.8
12	30.0	40.6	41.3
13	39.8	83.3	73.7
14	146.6	141.8	144.9
15	111.9	113.0	111.6
16	19.2	23.4	28.9
17	107.4	106.4	106.6
18	28.4	28.9	27.4
19	27.4	184.0	183.3
20	13.4	12.7	12.9
COCH ₃	–	21.0	–
COCH ₃	–	171.1	–

δ_C 171.1, both indicating the acylation of the OH group at C-13 with an acetyl moiety. Acylation of OH at C-13 was further confirmed by the down-field shift of C-13 (δ_C 83.3) relative to the same carbon atom in **3** (δ_C 73.7). The HMBC spectrum showed a long correlation between the signal at δ_C 83.3 (C-13) and proton signals at δ_H 5.88 (H-14) and 1.45 (H-16), which confirmed the presence of an acetyl group at C-13. Thus compound **2** was identified as 13-*O*-acetyl-13-epicupressic acid, a new natural product.

The resin extract of *A. heterophylla* Salisb. showed dose-dependent antiulcerogenic activity against ethanol-induced stomach ulcers in Sprague Dawley rats. A curative ratio of 55.87 compared to 61.64 for ranitidine at the same dose (100 mg/kg body wt.) was recorded, the curative ratio of the resin extract raised to 63.26 at a dose of 200 mg/kg body wt. The activity of the resin extract as a gastroprotective may be attributed to its diterpene constituents. In this concern several terpenes or their derivatives have been shown to possess gastroprotective activity in different models of induced gastric lesions in animals (Wada *et al.*, 1985; Giorando *et al.*, 1990; Lewis and Hanson, 1991; Farina *et al.*, 1998; Mastuda *et al.*, 1998). The gastroprotective activity of the diterpenes and their derivatives have been explained by mechanisms that include stimulation of prostaglandin synthesis, increase of mucus production, and suppression of gastric acid secretion (Hiruma-Lima *et al.*, 1999, 2002).

Table III. Results of the *in vitro* cytotoxicity test of the resin extract and isolated compounds of *A. heterophylla* Salisb.

Compound	IC ₅₀ [μ g/ml] ^a	
	MCF7 ^b	HCT116 ^b
Doxorubicin®	0.7	0.69
Resin extract	0.54	0.94
1	3.88	4.59
2	2.33	8.04
3	9.77	–

^a IC₅₀, the concentration that causes 50% death of the cancer cells.

^b MCF7, breast cancer cell line; HCT116, colon cancer cell line; –, no effect.

The resin extract of *A. heterophylla* Salisb. exhibited also a strong cytotoxic activity against breast (MCF7) and colon (HCT116) cancer cell lines; the recorded IC₅₀ values were 0.54 and 0.94 μ g/ml, respectively. The isolated compounds **1–3** were subjected to an *in vitro* cytotoxicity test. Only compounds **1** and **2** showed moderate cytotoxic activity against both tested cell lines with IC₅₀ values lower than that of the resin extract (IC₅₀ 2.33–8.04 μ g/ml), see Table III. Compound **3** showed only weak activity against the breast (MCF7) cancer cell line (IC₅₀ 9.77 μ g/ml). It is worth to mention that the *in vitro* cytotoxic effect of the resin extract was comparable to that of the reference drug Doxorubicin®. This effect may be a synergism of the effect of the diterpene content of the resin.

Angelopoulou D., Demetzos C., and Perdetzoglou D. (2002), Diurnal and seasonal variation of the essential oil labdanes and clerodanes from *Cistus monspeliensis* L. leaves. *Biochem. Syst. Ecol.* **30**, 189–203.

Caputo R. and Mangoni L. (1974), Diterpenes from *Araucaria bidwillii*. *Phytochemistry* **13**, 467–470.

Caputo R., Mangoni L., and Monaco P. (1972), Diterpenes of *Araucaria excels.* *Phytochemistry* **11**, 839–840.

Caputo R., Dovinola V., and Mangoni L. (1974a), New diterpenes from *Araucaria cunninghami*. *Phytochemistry* **13**, 475–478.

Caputo R., Mangoni L., Monaco P., and Previtera L. (1974b), New labdane diterpenes from *Araucaria cookii*. *Phytochemistry* **13**, 471–474.

Cho C. H. and Ogle C. W. A. (1978), Correlative study of the antiulcer effect of zinc sulphate in stressed rats. *Eur. J. Pharmacol.* **48**, 97–105.

Cox R. E., Yamamoto S., Otto A., and Simoneit B. R. T. (2007), Oxygenated di- and tricyclic diterpenoids of

southern hemisphere conifers. *Biochem. Syst. Ecol.* **35**, 342–362.

Farina C., Pinza M., and Pifferi G. (1998), Synthesis and antiulcer activity of new derivatives of glycyrrhetic, oleanolic and urosolic acids. *Farmacol.* **53**, 22–32.

Fonseca S. F., Campello J. D. P., Barata L. E. S., and Ruveda E. A. (1978), ¹³C NMR spectral analysis of lignans from *Araucaria angustifolia*. *Phytochemistry* **17**, 499–502.

Fonseca S. F., Nielsen L. T., and Ruveda E. A. (1979), Lignans of *Araucaria angustifolia* and ¹³C NMR analysis of some phenyltetralin lignans. *Phytochemistry* **18**, 1703–1708.

Garg G. P., Nigam S. K., and Ogle C. W. (1993), The gastric antiulcer effect of the leaves of Neem tree. *Planta Med.* **59**, 215–217.

Giorando O. S., Guerreiro E., Pestchanker M. J., Guzman J., Pastor D., and Guardia T. (1990), The gastric cytoprotective effect of several sesquiterpene lactones. *J. Nat. Prod.* **53**, 803–809.

- Glavin G. B. and Mikhaeil A. A. (1976), Stress and ulcer etiology in the rat. *Physiol. Behav.* **16**, 135–139.
- Hiruma-Lima C., Bratisch R., Kassisse D., and Souza-Brito A. R. (1999), Antiulcerogenic mechanisms of dehydrocrotonin, a diterpene lactone obtained from *Croton cajucara*. *Planta Med.* **65**, 325–330.
- Hiruma-Lima C., Toma W., Gracioso J. S., Almeida A. B. A., Batista L. M., Magri L., Paula A. C. B., Soares F. R., Nunes D. S., and Souza-Brito A. R. M. (2002), Natural trans-crotonin: the antiulcerogenic effect of another diterpene isolated from the bark of *Croton cajucara* Benth. *Biol. Pharm. Bull.* **25**, 452–456.
- Lee C. K. and Cheng Y. S. (2001), Diterpenoids from the leaves of *Juniperus chinensis* var. *kaizuka*. *J. Nat. Prod.* **64**, 511–514.
- Lewis D. A. and Hanson P. J. (1991), Antiulcer drugs of plant origin. *Prog. Med. Chem.* **28**, 201–231.
- Mastuda H., Li Y., Murakami T., Yamohara J., and Yoshikawa M. (1998), Protective effect of oleanolic acid oligoglycosides on ethanol or indomethacin-induced gastric mucosal lesions in rats. *Life Sci.* **63**, 245–250.
- Ohashi H., Kawai S., Sakurai Y., and Yasue M. (1992), Norlignan from the knot resin of *Araucaria angustifolia*. *Phytochemistry* **31**, 1371–1373.
- Schans A. V. D., Schokman L. M., and Cloudy J. B. (2004), *Tropical and Subtropical Trees*. Thames & Hudson Ltd, Holborn, London, p. 30.
- Schmeda-Hirschmann G., Astudillo L., Rodriguez J. A., Theoduloz C., and Yanez T. (2005a), Gastroprotective effect of the mapuche crude drug *Araucaria araucana* resin and its main constituents. *J. Ethnopharmacol.* **101**, 271–276.
- Schmeda-Hirschmann G., Astudillo L., Sepulveda B., Rodriguez J. A., Theoduloz C., Yanez T., and Palenzuela J. A. (2005b), Gastroprotective effect and cytotoxicity of natural and semisynthetic labdane diterpenes from *Araucaria araucana* resin. *Z. Naturforsch.* **60c**, 511–522.
- Skehan P., Storeng R., Scudiero D., Monks A., Mahon J. M., Vistica D., Warren J. T., Bokesch H., Kenney S., and Boyd M. R. (1990), Cytotoxic assay for anti-cancer drug screening. *J. Natl. Cancer Inst.* **82**, 1107–1112.
- Snedecor G. W. and Cochran W. G. (1969), *Statistical Methods*, 4th ed. Iowa State University Press, Ames, Iowa, USA, p. 91.
- Su W. C., Fang J. M., and Cheng Y. S. (1994), Labdanes from *Cryptomeria japonica*. *Phytochemistry* **37**, 1109–1114.
- Trease G. E. and Evans W. C. (1978), *Pharmacognosy*, 11th ed. Macmillan Publishing Co. Inc., New York, London, p. 82.
- Wada H., Kodato S., Kawamori M., Morikawa T., Nakai H., Takeda M., Saito S., Onoda Y., and Tamaki H. (1985), Antiulcer activity of dehydroabietic acid derivatives. *Chem. Pharm. Bull.* **33**, 1472–1487.
- Wang Y. Z., Tang C. P., Ke C. Q., Weiss H. C., Gesing E. R., and Ye Y. (2008), Diterpenoids from the pericarp of *Platycladus orientalis*. *Phytochemistry* **69**, 518–526.