

Parallel Solution-phase Synthesis of (2*S*,4*E*)-4-(Arylaminomethylidene)pyroglutamic Acids

Jurij Svete^a, Uroš Grošelj^a, Jernej Baškovč^a, Georg Dahmann^b, and Branko Stanovnik^a

^a Faculty of Chemistry and Chemical Technology, University of Ljubljana, Aškerčeva 5, P. O. Box 537, 1000 Ljubljana, Slovenia

^b Boehringer Ingelheim Pharma GmbH & Co. KG, Dept. Chemical Research, 88397 Biberach, Germany

Reprint requests to Prof. Jurij Svete. Fax: +386 1 2419 220. E-mail: jurij.svete@fkkt.uni-lj.si

Z. Naturforsch. **2010**, *65b*, 811–820; received January 26, 2010

A library of twelve *N*(4′)-substituted di-*tert*-butyl (2*S*,4*E*)-4-arylaminomethylidene-5-oxopyrrolidine-1,2-dicarboxylates **6/6′a–l** were prepared in 47–90 % yield by parallel acid-catalysed treatment of di-*tert*-butyl (2*S*,4*E*)-4-[(dimethylamino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (**4**) with anilines **5a–j**, ethyl glycinate (**5k**), and ethyl β-alaninate (**3l**). Acidolytic deprotection of compounds **6a–c**, **e–j** afforded the corresponding (2*S*,4*E*)-4-arylaminomethylidene-5-oxopyrrolidine-2-carboxylic acids **7a–c**, **e–j** in 39–99 % yield. The configuration around the C=C double bond in the enaminones **6** and **7** was determined by NMR spectroscopy.

Key words: Pyroglutamic Acid, Enaminones, Amines, Combinatorial Synthesis, Pyrrolidinone

Introduction

(*S*)-Pyroglutamic acid (**1**) is a naturally occurring heterocyclic α-amino acid which is abundant in peptides and proteins, and its structure can also be found in a variety of other biologically important compounds. On the other hand, **1** is also a useful chiral building block, which is frequently used as a commercially available starting material in chiral-pool syntheses of peptidomimetics, natural products, and their analogues. Therefore it is not surprising, that several reviews on the chemistry of (*S*)-pyroglutamic acid (**1**) have recently been published [1].

2-Substituted alkyl 3-(dimethylamino)prop-2-enates and related enaminones are a group of enamino-masked alkyl α-formylacetates, which are easily available and versatile reagents in heterocyclic synthesis [2]. In addition to their extensive use in the synthesis of various heterocyclic systems, recent applications of enaminones have mostly focused on the synthesis of functionalised heterocyclic compounds including natural product analogues [2–4], and on combinatorial syntheses of functionalised heterocycles [5]. Within this context, various functionalised enaminones have been prepared and used as key intermediates in the synthesis of 3-heteroarylalanine derivatives [6], histamine analogues [5b, 7], and het-




Fig. 1. α-Enamino (*S*)-pyroglutamic acid **7** as a conformationally constrained heterocyclic analogue of Δ-β-Ala-AlaOH.


erocyclic analogues of dipeptides containing the (*S*)-pyroglutamic acid structural motif [5c, 8]. Previously, we reported the synthesis of a series of *N*(4′)-substituted methyl (2*S*,4*E*)-1-acyl-4-(aminomethylidene)-5-oxopyrrolidine-2-carboxylates as stable intermediates in the ‘ring switching’ synthesis of 3-heteroarylalanines [9]. Recently, this type of compounds attracted our attention again, since such α-enamino pyroglutamic acids are conformationally constrained heterocyclic dipeptides comprising α,β-dehydro-β-alanine (Δ-β-Ala) and (*S*)-alanine (L-Ala) structural units (Fig. 1).

Therefore, we were intrigued to study the synthesis of γ-enamino pyroglutamic acids **7**, which might be interesting and useful building blocks for further derivatisation. Herein, we report the result of this study – a simple parallel solution-phase

synthesis of di-*tert*-butyl (*S*)-4-arylamino-methylidene-5-oxopyrrolidine-1,2-dicarboxylates **6/6'a–l** and (*S*)-4-arylamino-methylidene-5-oxopyrrolidine-1,2-dicarboxylates **7a–c, e–j**.

Results and Discussion

The key intermediate, di-*tert*-butyl (2*S*,4*E*)-4-[(dimethylamino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (**4**), was prepared in three steps from (*S*)-



Scheme 1.

Table 1. Selected experimental data of the di-*tert*-butyl (2*S*,4*E*)-4-arylamino-methylidene-5-oxopyrrolidine-1,2-dicarboxylates **6a–l** and **7a–c, e–j**.

Compound	Ar	Yield (%)	<i>E</i> : <i>Z</i> ^a	Purity (%) ^b
6a/6'a	phenyl	90	56:44 ^{c,d}	> 95
6/6'b	3-methylphenyl	65	21:79	> 95
6/6'c	4-methylphenyl	83	17:83	> 95
6d	3-hydroxyphenyl	62	100:0	> 95
6e	4-hydroxyphenyl	69	100:0	> 95
6/6'f	3-methoxyphenyl	73	16:84	> 95
6/6'g	3-bromophenyl	73	15:85	> 95
6/6'h	4-bromophenyl	79	15:85	> 95
6'i	3-nitrophenyl	65	100:0	> 95
6j	4-nitrophenyl	81	100:0	> 95
6/6'k	CH ₂ COOEt	47	85:15	> 95
6/6'l	CH ₂ CH ₂ COOEt	68	88:12	> 95
7a	phenyl	39	100:0	> 95
7b	3-methylphenyl	96	100:0	> 95
7c	4-methylphenyl	90	100:0	> 95
7e	4-hydroxyphenyl	37	100:0	> 95
7f	3-methoxyphenyl	99	100:0	> 95
7g	3-bromophenyl	87	100:0	> 95
7h	4-bromophenyl	86	100:0 ^d	> 95
7i	3-nitrophenyl	90	100:0 ^{c,d}	> 95
7j	4-nitrophenyl	88	100:0 ^d	> 95

^a Determined by ¹H-NMR spectroscopy; ^b determined by CHN-analyses and ¹H-NMR spectroscopy. The values found for C, H, and N were within ±0.4% with respect to the calculated values; ^c the configuration around the C=C double bond was determined by HMBC NMR spectroscopy; ^d the configuration around the C=C double bond was determined by NOESY spectroscopy.

pyroglutamic acid (**1**) following the literature procedures [5c, 10, 11]. Thus, acid-catalysed esterification of (*S*)-pyroglutamic acid (**1**) with *tert*-butyl acetate gave *tert*-butyl pyroglutamate (**2**) [10], which was *N*-acylated with Boc₂O in acetonitrile in the presence of catalytic amounts of 4-dimethylaminopyridine (DMAP) to afford di-*tert*-butyl (2*S*)-5-oxopyrrolidine-1,2-dicarboxylate (**3**) [11]. Finally, heating of **3** with one equivalent of *tert*-butoxy-bis(dimethylamino)methane (Bredereck's reagent, TBDMAM) furnished di-*tert*-butyl (2*S*,4*E*)-4-[(dimethylamino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (**4**) in 70% yield over three steps [5c] (Scheme 1).

Further parallel treatment of the enaminolactam **4** with 1.2 equivalents of amine hydrochlorides **5a–l** in 50% aqueous ethanol at r.t. resulted in the formation of the dimethylamine substitution products, *N*-(4')-substituted di-*tert*-butyl (2*S*,4*EZ*)-4-amino-methylidene-5-oxopyrrolidine-1,2-dicarboxylates **6/6'a–l**, which precipitated from the reaction mixtures and were isolated by filtration. Upon washing with water and thorough drying *in vacuo* over P₄O₁₀, a library of twelve analytically pure esters **6/6'a–l** was




Fig. 2. Structure determination by HMBC and NOESY spectroscopy.

obtained in 47–90% yield. In the final step, parallel acidolytic deprotection of compounds **6/6'a–l** was carried out. Stirring of esters **6/6'** with 2 M HCl/EtOAc at r. t. for 12 h resulted in the initial formation of clear colourless or yellow solutions followed by formation of precipitates, which were then collected by filtration, washed with EtOAc, and dried *in vacuo* over P_4O_{10} . In this manner, analytically pure carboxylic acids **7a–c**, **e–j** were prepared in 37–99% yield. Acidolysis of compounds **6d**, **k**, **l** did not produce precipitates. Attempts to isolate the final products **7d**, **k**, **l** by evaporation of the reaction mixtures followed by crystallisation and/or chromatographic workup failed. Since we were particularly interested in a simple and practical procedure using just a filtration work-up, no further attempts to isolate compounds **7d**, **k**, **l** were made (Scheme 1, Table 1).

The structures of the novel compounds **6/6'a–l** and **7a–c**, **e–j** were determined by spectroscopic (NMR, IR, MS) methods and by elemental analyses. The configuration around the exocyclic C=C bond in compounds **6a**, **6'a**, and **7i** was determined by HMBC NMR spectroscopy on the basis of the long-range coupling constant ($^3J_{C-H}$) between the methyldene proton ($H-C(4')$) and the carbonyl carbon atom ($O=C(5)$),

measured from the antiphase splitting of cross peaks. Generally, the magnitude of this coupling constant, $^3J_{C-H}$, for nuclei with *cis*-configuration around the C=C double bond is smaller (2–6 Hz) than that for *trans*-oriented nuclei (8–12 Hz) [2, 12]. In compounds **6a** and **7i**, the magnitude of the coupling constants, $^3J_{C(1)-H(4')} = 4 \text{ Hz (cis)}$ and 5 Hz (cis) , respectively, confirmed the (*E*)-configuration around the exocyclic C=C double bond. In the minor isomer **6'a**, on the other hand, a large coupling constant, $^3J_{C(1)-H(4')} = 10 \text{ Hz (trans)}$, confirmed the (*Z*)-configuration around the C=C double bond (Fig. 2). Additionally, the configuration around the exocyclic C=C double bond in compounds **6a**, **6'a**, and **7h–j** was confirmed by NOESY spectroscopy. A NOE between the NH and the CH_2 group in compounds **6a** and **7h–j** was in agreement with the (*E*)-configuration around the exocyclic C=C double bond, whilst a NOE between the $4'$ -H and the CH_2 group in the minor isomer **6'a** supported the (*Z*)-configuration (Fig. 2).

Finally, the configurations around the C=C double bond in compounds **6** and **7** were confirmed by correlation of chemical shifts for $4'$ -H and NH and vicinal coupling constants, $^3J_{H-H}$. The major and most characteristic difference was observed in chemical shifts δ

Compound	Solvent	δ (ppm)		3–4'	$^3J_{\text{H-H}}$ (Hz) 3a–3b	CHNH
		4'-H	4'-NH			
(2 <i>S</i> ,4 <i>E</i>)-Isomers 6 and 7						
6a	CDCl ₃	7.79	6.09	2.2	15.8	13.7
6b	CDCl ₃	7.79	6.14	^a	^a	13.5
6c	CDCl ₃	7.76	6.09	^a	^a	13.6
6d	[D ₆]DMSO	7.50	8.89	1.9	16.4	13.1
6e	[D ₆]DMSO	7.47	8.77	1.8	16.1	13.4
6f	CDCl ₃	7.76	6.09	^a	^a	13.5
6g	CDCl ₃	7.70	6.24	^a	^a	13.3
6h	CDCl ₃	7.68	6.42	2.1	16.1	13.4
6i	CDCl ₃	7.74	7.39	2.3	16.4	13.0
6j	CDCl ₃	7.74	7.18	2.2	16.5	13.2
6j	[D ₆]DMSO	7.79	9.62	2.4	17.0	^b
6k	CDCl ₃	7.11	~ 4.5 ^c	2.0	15.4	13.2
6l	CDCl ₃	7.17	4.65	1.9	15.3	13.6
7a	[D ₆]DMSO	7.37	8.59	2.1	16.7	11.6
7b	[D ₆]DMSO	7.37	8.53	2.1	16.7	12.0
7c	[D ₆]DMSO	7.33	8.47	2.1	16.7	10.4
7e	[D ₆]DMSO	7.29	8.34	2.0	16.7	^b
7f	[D ₆]DMSO	7.33	8.53	2.2	16.7	12.3
7g	[D ₆]DMSO	7.27	8.65	2.1	16.9	12.3
7h	[D ₆]DMSO	7.28	8.65	2.2	16.7	12.7
7i	[D ₆]DMSO	7.37	9.04	2.2	17.0	12.3
7j	[D ₆]DMSO	7.43	9.44	2.4	16.7	12.2
(2 <i>S</i> ,4 <i>Z</i>)-Isomers 6'						
6'a	CDCl ₃	7.17	9.95	1.3	15.2	12.3
6'b	CDCl ₃	7.17	9.89	1.2	15.2	12.2
6'c	CDCl ₃	7.13	9.90	1.2	15.1	12.3
6'f	CDCl ₃	7.14	9.92	1.3	15.3	12.4
6'g	CDCl ₃	~ 7.1 ^a	9.94	1.4	15.4	12.0
6'h	CDCl ₃	7.09	9.94	1.4	15.4	12.1
6'k	CDCl ₃	6.46	7.96	^a	^a	12.6
6'l	CDCl ₃	6.56	~ 7.9 ^c	^a	^a	12.6

Table 2. Correlation between the chemical shifts for 4'-H and (4')N-H and coupling constants, $^3J_{\text{H3-H4}}$ and $^3J_{\text{CH-NH}}$, in compounds **6a-l**, **6'a-c**, **f-h**, **k**, **l**, and **7a-c**, **e-j**.

^a Overlapped by other signals; ^b the signals for the CHNH fragment appeared as two broad singlets; ^c broad signal.

for the aminomethylidene -NH-CH= protons. In the $^1\text{H-NMR}$ spectra of the *E/Z*-mixtures **6/6'a-c**, **f-h**, **k**, **l** taken in CDCl₃, the NH protons of the (*E*)-isomers **6a-c**, **f-h**, **k**, **l** had lower δ values (~ 6.2 ppm) than the 4'-H protons (~ 7.7 ppm), while in the (*Z*)-isomers **6'a-c**, **f-h**, **k**, **l** the NH protons exhibited higher δ values (~ 9.9 ppm) than the 4'-H protons (~ 7.2 ppm). Such a difference in chemical shifts of the NH protons in CDCl₃ solution could be explained by intramolecular $\text{N-H}\cdots\text{O=C}$ hydrogen bonding, which is possible only in the (*Z*)-isomers **6'a-c**, **f-h**, **k**, **l** and not in the (*E*)-isomers **6a-c**, **f-h**, **k**, **l**. Accordingly, in [D₆]DMSO as a hydrogen bond acceptor, the NH protons of the (*E*)-isomers **6d**, **e**, **j** and **7a-c**, **e-j** appear at ~ 8.5 ppm. Similarly, small yet characteristic differences between typical coupling constant values, $^2J_{\text{H3aH3b}}$, $^3J_{\text{NHCH}}$, and $^4J_{\text{H3H4'}}$, were also observed; typical values were larger in case of the (*E*)-isomers: $^2J_{\text{H3aH3b}} \sim 16 \text{ Hz}$ (*E*) $>$ $^2J_{\text{H3aH3b}} \sim 15 \text{ Hz}$ (*Z*), $^3J_{\text{NHCH}} \sim 13.5 \text{ Hz}$ (*E*) $>$ $^3J_{\text{NHCH}} \sim 12.5 \text{ Hz}$

(*Z*), and $^4J_{\text{H3H4'}} \sim 2 \text{ Hz}$ (*E*) $>$ $^4J_{\text{H3H4'}} \sim 1.3 \text{ Hz}$ (*Z*). These characteristic values are also in agreement with the literature data for related aminomethylidene compounds [2–4, 13] (Table 2, Fig. 2).

Conclusion

Di-*tert*-butyl(2*S*,4*E*)-4-[(dimethylamino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (**4**) is an easily available reagent, which is suitable for two-step parallel solution-phase synthesis of 4-arylaminomethylidene-substituted (*S*)-pyroglutamic acids **7** as conformationally constrained analogues of *N*-[*N*-(aryl)- α,β -didehydro- β -alanyl]-(*S*)-alanine (*c.f.* Fig. 1). The synthesis comprises acid-catalysed substitution of the dimethylamino group in the enamino lactam **4** with aromatic and aliphatic primary amines **5** to give the substitution products **6/6'**, followed by acidolytic deprotection to furnish the title compounds **7** in good yields over two steps. Furthermore, all intermediates

6a–l and final products **7a–c**, **e–j** were obtained in analytical purity following a simple parallel filtration work-up protocol in both synthetic steps. In conclusion, this synthetic method offers an easy access to diversity-oriented libraries of (*S*)-4-[(substituted amino)methylidene]pyroglutamic acid derivatives in search for novel bioactive compounds.

Experimental Section

Melting points were determined on a Stanford Research Systems MPA100 OptiMelt automated melting point system. The NMR spectra were obtained on a Bruker Avance DPX 300 spectrometer at 300 MHz for ^1H and 75.5 MHz for ^{13}C , using CDCl_3 and $[\text{D}_6]\text{DMSO}$ (with TMS as the internal standard) as solvents. Mass spectra were recorded on a Q-TOF Premier spectrometer, IR spectra on a Perkin-Elmer Spectrum BX FTIR spectrophotometer. Microanalyses were performed on a Perkin-Elmer CHN analyser 2400 II.

(*S*)-Pyroglutamic acid (**1**), di-*tert*-butoxy-bis(dimethylamino)methane (Bredereck's reagent), and amines **5a–l** are commercially available (Sigma Aldrich). Di-*tert*-butyl (2*S*,4*E*)-4-[(dimethylamino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (**4**) was prepared from **1** following the literature procedures [5c, 10, 11].

Parallel stirring and filtrations were carried out on a Mettler-Toledo Bohdan MiniBlockTM Compact Shaking and Washing Station and Vacuum Collection Base (12 positions, Vortex stirring, 400 r. p. m. in all cases).

Parallel solution-phase synthesis of *N*(4')-substituted di-*tert*-butyl (2*S*)-4-aminomethylidene-5-oxopyrrolidine-1,2-dicarboxylates **6/6'a–l**

The MiniBlockTM was assembled with 12 fritted vessels, and the frits were wetted with ethanol (0.5 mL each). A stock solution of enaminone **4** (0.25 M in ethanol, 12 × 4 mL, 12 × 1 mmol) was added followed by addition of aqueous solutions of amines **5a–l** hydrochlorides* (0.25 M in water, 12 × 5 mL, 12 × 1.2 mmol). The MiniBlockTM was closed and the reaction mixtures were stirred at 20 °C for 24 h. The precipitates were collected by filtration, washed with water (12 × 3 mL), and dried *in vacuo* at r. t. over P_4O_{10} for 24 h to give **6/6'a–l**.

The following compounds were prepared in this manner:

Di-*tert*-butyl (2*S*,4*E*)-4-anilinoethylidene-5-oxopyrrolidine-1,2-dicarboxylate (**6a**) and its minor (4*Z*)-isomer **6'a**

Compound **6a** was prepared from **4** and aniline hydrochloride (**5a**). Yield: 365 mg (90 %) of a pale-beige

solid. – M. p. 177–180 °C; **6a**: **6'a** = 56:44. – $[\alpha]_{589}^{20} +19.5$ ($c = 0.33$, CH_2Cl_2). – IR (KBr): $\nu = 3237, 3120, 3040, 2978, 1757, 1710, 1686, 1639, 1594, 1495, 1445, 1369, 1313, 1231, 1151, 1000, 959, 866, 806, 775, 753, 692\text{ cm}^{-1}$. – $^1\text{H-NMR}$ (CDCl_3), *major* (*E*)-isomer **6a**: $\delta = 1.48$ and 1.53 (18H, 2s, 1:1, $2 \times \text{}^t\text{-Bu}$), 2.54 (1H, ddd, $J = 2.2, 3.6, 15.8\text{ Hz}$, 3-Ha), 2.98 (1H, ddd, $J = 2.2, 10.7, 15.8\text{ Hz}$, 3-Hb), 4.56 (1H, dd, $J = 3.6, 10.7\text{ Hz}$, 2-H), 6.09 (1H, d, $J = 13.7\text{ Hz}$, NH), 6.90–7.05 (3H, m, *o*, *p*- C_6H_5), 7.24–7.35 (2H, m, *m*- C_6H_5), 7.79 (1H, dt, $J = 1.9, 13.7\text{ Hz}$, 4'-H); *minor* (*Z*)-isomer **6'a**: $\delta = 1.48$ and 1.54 (18H, 2s, 1:1, $2 \times \text{}^t\text{-Bu}$), 2.61 (1H, ddd, $J = 1.3, 3.5, 15.2\text{ Hz}$, 3-Ha), 3.06 (1H, ddd, $J = 1.3, 10.6, 15.2\text{ Hz}$, 3-Hb), 4.51 (1H, dd, $J = 3.7, 10.6\text{ Hz}$, 2-H), 6.90–7.05 (3H, m, *o*, *p*- C_6H_5), 7.17 (1H, br d, $J = 12.3\text{ Hz}$, 4'-H), 7.24–7.35 (2H, m, *m*- C_6H_5), 9.95 (1H, d, $J = 12.3\text{ Hz}$, NH). – $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_5$ (388.5): calcd. C 64.93, H 7.27, N 7.21; found C 65.11, H 7.48, N 7.44.

Di-*tert*-butyl (2*S*,4*Z*)-4-[(3-methylanilino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (**6'b**) and its minor (4*E*)-isomer **6b**

Compound **6'b** was prepared from **4** and 3-methylaniline hydrochloride (**5b**). Yield: 260 mg (65 %) of a pale-beige solid. – M. p. 155–157 °C; **6b**: **6'b** = 21:79. – $[\alpha]_{589}^{20} +17.4$ ($c = 0.48$, CH_2Cl_2). – IR (KBr): $\nu = 3464, 3318, 2979, 2933, 1756, 1686, 1632, 1599, 1368, 1321, 1246, 1231, 1155, 1001, 1155, 781\text{ cm}^{-1}$. – $^1\text{H-NMR}$ (CDCl_3), *major* (*Z*)-isomer **6'b**: $\delta = 1.48$ and 1.54 (18H, 2s, 1:1, $2 \times \text{}^t\text{-Bu}$), 2.31 (3H, s, Me), 2.60 (1H, ddd, $J = 1.2, 3.6, 15.2\text{ Hz}$, 3-Ha), 3.06 (1H, ddd, $J = 1.2, 10.6, 15.2\text{ Hz}$, 3-Hb), 4.50 (1H, dd, $J = 3.6, 10.6\text{ Hz}$, 2-H), 6.69–6.85 (3H, m, *o*, *p*- C_6H_4), 7.12–7.20 (1H, m, *m*- C_6H_4), 7.17 (1H, br d, $J = 12.2\text{ Hz}$, 4'-H), 9.89 (1H, d, $J = 12.2\text{ Hz}$, NH); *minor* (*E*)-isomer **6b**: $\delta = 2.32$ (3H, s, Me), 4.55 (1H, dd, $J = 3.4, 10.7\text{ Hz}$, 2-H), 6.14 (1H, broad signal, NH), 7.79 (1H, br d, $J = 13.5\text{ Hz}$, 4'-H). – $\text{C}_{22}\text{H}_{30}\text{N}_2\text{O}_5$ (402.5): calcd. C 65.65, H 7.51, N 6.96; found C 65.81, H 7.72, N 7.17.

Di-*tert*-butyl (2*S*,4*Z*)-4-[(4-methylanilino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (**6'c**) and its minor (4*E*)-isomer **6c**

Compound **6'c** was prepared from **4** and 4-methylaniline hydrochloride (**5c**). Yield: 334 mg (83 %) of a pale-beige solid. – M. p. 163–170 °C; **6c**: **6'c** = 17:83. – $[\alpha]_{589}^{20} +21.5$ ($c = 0.40$, CH_2Cl_2). – IR (KBr): $\nu = 3454, 3285, 2980, 2930, 1761, 1710, 1684, 1641, 1524, 1454, 1366, 1307, 1233, 1153, 999, 962, 870, 810, 774\text{ cm}^{-1}$. – $^1\text{H-NMR}$ (CDCl_3), *major* (*Z*)-isomer **6'c**: $\delta = 1.48$ and 1.54 (18H, 2s, 1:1, $2 \times \text{}^t\text{-Bu}$), 2.28 (3H, s, Me), 2.60 (1H, ddd, $J = 1.2, 3.7, 15.1\text{ Hz}$, 3-Ha), 3.05 (1H, ddd, $J = 1.2, 10.6, 15.1\text{ Hz}$, 3-Hb), 4.50 (1H, dd, $J = 3.7, 10.6\text{ Hz}$, 2-H), 6.83 (2H, br d, $J = 8.3\text{ Hz}$, *m*- C_6H_4), 7.08 (2H, br d, $J = 8.3\text{ Hz}$, *o*- C_6H_4), 7.13 (1H, br

*In the case of anilines **5i**, **j**, the solid-free nitroanilines **5i** and **5j** were added to ethanolic solutions of **4**, followed by addition of 0.25 M aq. HCl.

d, $J = 12.3$ Hz, 4'-H), 9.90 (1H, d, $J = 12.3$ Hz, NH); *minor* (*E*)-isomer **6c**: $\delta = 2.29$ (3H, s, Me), 4.55 (1H, dd, $J = 3.6$, 10.6 Hz, 2-H), 6.09 (1H, broad signal, NH), 6.83 (2H, br d, $J = 8.3$ Hz, *m*-C₆H₄), 7.76 (1H, br d, $J = 13.6$ Hz, 4'-H). – C₂₂H₃₀N₂O₅ (402.5): calcd. C 65.65, H 7.51, N 6.96; found C 65.73, H 7.72, N 7.28.

Di-tert-butyl (2S,4E)-4-[(3-hydroxyanilino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (6d)

Compound **6d** was prepared from **4** and 3-hydroxyaniline hydrochloride (**5d**). Yield: 251 mg (62 %) of a pale-grey solid. – M. p. 180–181 °C; **6d**: **6'd** = 100:0. – $[\alpha]_{589}^{20} +2.0$ ($c = 0.49$, EtOH). – IR (KBr): $\nu = 3383$, 3307, 2978, 1757, 1709, 1688, 1634, 1612, 1500, 1460, 1392, 1369, 1346, 1305, 1240, 1156, 979, 959, 779 cm⁻¹. – ¹H-NMR ([D₆]DMSO): $\delta = 1.42$ and 1.44 (18H, 2s, 1:1, 2 × ^{*t*}-Bu), 2.53 (1H, ddd, $J = 1.9$, 3.4, 16.4 Hz, 3-Ha), 2.99 (1H, ddd, $J = 1.9$, 10.7, 16.4 Hz, 3-Hb), 4.52 (1H, dd, $J = 3.4$, 10.7 Hz, 2-H), 6.36 (1H, dd, $J = 1.6$, 7.9 Hz, *o*-C₆H₄), 6.49–6.57 (2H, m, *o,m*-C₆H₄), 7.06 (1H, t, $J = 8.0$ Hz, *m*-C₆H₄), 7.50 (1H, br d, $J = 13.1$ Hz, 4'-H), 8.89 (1H, d, $J = 13.1$ Hz, NH), OH exchanged. – C₂₁H₂₈N₂O₆ (404.5): calcd. C 62.36, H 6.98, N 6.93; found C 62.62, H 7.22, N 6.92.

Di-tert-butyl (2S,4E)-4-[(4-hydroxyanilino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (6e)

Compound **6e** was prepared from **4** and 4-hydroxyaniline hydrochloride (**5e**). Yield: 279 mg (69 %) of a pale-grey solid. – M. p. 167–168 °C; **6e**: **6'e** = 100:0. – $[\alpha]_{589}^{20} -2.1$ ($c = 0.34$, EtOH). – IR (KBr): $\nu = 3402$, 3321, 2978, 2935, 1724, 1695, 1634, 1519, 1433, 1391, 1369, 1321, 1225, 1150, 1005, 962, 822, 791, 745, 673 cm⁻¹. – ¹H-NMR ([D₆]DMSO): $\delta = 1.42$ and 1.44 (18H, 2s, 1:1, 2 × ^{*t*}-Bu), 2.48 (1H, ddd, $J = 1.8$, 3.5, 16.1 Hz, 3-Ha), 2.96 (1H, ddd, $J = 1.8$, 10.7, 16.1 Hz, 3-Hb), 4.51 (1H, dd, $J = 3.5$, 10.8 Hz, 2-H), 6.70 (2H, d, $J = 8.8$ Hz, *o*-C₆H₄), 6.95 (2H, d, $J = 8.8$ Hz, *m*-C₆H₄), 7.47 (1H, br d, $J = 13.4$ Hz, 4'-H), 8.77 (1H, d, $J = 13.4$ Hz, NH), OH exchanged. – C₂₁H₂₈N₂O₆ (404.5): calcd. C 62.36, H 6.98, N 6.93; found C 62.66, H 7.21, N 7.09.

Di-tert-butyl (2S,4Z)-4-[(3-methoxyanilino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (6f) and its minor (4E)-isomer 6f'

Compound **6f** was prepared from **4** and 3-methoxyaniline hydrochloride (**5f**). Yield: 304 mg (73 %) of a grey solid. – M. p. 151–156 °C; **6f**: **6'f** = 16:84. – $[\alpha]_{589}^{20} +19.5$ ($c = 1.00$, CH₂Cl₂). – IR (KBr): $\nu = 3453$, 3251, 3127, 2978, 2934, 1761, 1733, 1682, 1637, 1593, 1537, 1497, 1481, 1456, 1391, 1368, 1310, 1286, 1244, 1223, 1196, 1155, 1146, 1047, 997, 964, 929, 842, 768, 688 cm⁻¹. – ¹H-NMR (CDCl₃), *major* (*Z*)-isomer **6f**: $\delta = 1.48$ and 1.54 (18H,

2s, 1:1, 2 × ^{*t*}-Bu), 1.58 (0.7 H, s, 1/3H₂O), 2.61 (1H, ddd, $J = 1.3$, 3.6, 15.3 Hz, 3-Ha), 3.06 (1H, ddd, $J = 1.3$, 10.6, 15.3 Hz, 3-Hb), 3.79 (3H, s, OMe), 4.51 (1H, dd, $J = 3.6$, 10.6 Hz, 2-H), 6.46 (1H, br t, $J = 2.2$ Hz, *o*-C₆H₄), 6.52 (1H, d, $J = 8.1$ Hz, *p*-C₆H₄), 6.53 (1H, d, $J = 8.1$ Hz, *o*-C₆H₄), 7.14 (1H, dt, $J = 1.3$, 12.3 Hz, 4'-H), 7.17 (1H, t, $J = 8.1$ Hz, *m*-C₆H₄), 9.92 (1H, d, $J = 12.3$ Hz, NH); *minor* (*E*)-isomer **6f'**: $\delta = 3.80$ (3H, s, OMe), 6.09 (1H, br d, $J = 13.5$ Hz, NH), 7.76 (1H, br d, $J = 13.5$ Hz, 4'-H). – C₂₂H₃₀N₂O₆ · 1/3H₂O (424.5): calcd. C 62.25, H 7.28, N 6.60; found C 62.35, H 7.34, N 6.58.

Di-tert-butyl (2S,4Z)-4-[(3-bromoanilino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (6'g) and its minor (4E)-isomer 6g

Compound **6'g** was prepared from **4** and 3-bromoaniline hydrochloride (**5g**). Yield: 341 mg (73 %) of a pale-grey solid. – M. p. 163–166 °C; **6g**: **6'g** = 15:85. – $[\alpha]_{589}^{20} +29.0$ ($c = 1.00$, CH₂Cl₂). – IR (KBr): $\nu = 3447$, 3295, 3260, 2979, 2933, 1763, 1734, 1716, 1686, 1638, 1594, 1475, 1368, 1312, 1236, 1223, 1154, 964, 899, 964, 773, 680 cm⁻¹. – ¹H-NMR (CDCl₃), *major* (*Z*)-isomer **6'g**: $\delta = 1.48$ and 1.54 (18H, 2s, 1:1, 2 × ^{*t*}-Bu), 2.61 (1H, ddd, $J = 1.4$, 3.5, 15.4 Hz, 3-Ha), 3.06 (1H, ddd, $J = 1.4$, 10.5, 15.4 Hz, 3-Hb), 4.51 (1H, dd, $J = 3.5$, 10.5 Hz, 2-H), 6.82 (1H, br dt, $J = 1.8$, 7.4 Hz, *o*-C₆H₄), 7.05–7.16 (4H, m, 3H of C₆H₄ and 4'-H), 9.94 (1H, d, $J = 12.0$ Hz, NH); *minor* (*E*)-isomer **6g**: $\delta = 6.24$ (1H, br d, $J = 13.3$ Hz, NH), 7.70 (1H, br d, $J = 13.3$ Hz, 4'-H). – C₂₁H₂₇BrN₂O₅ (467.4): calcd. C 53.97, H 5.82, N 5.99; found C 54.01, H 5.95, N 6.01.

Di-tert-butyl (2S,4E)-4-[(4-bromoanilino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (6h) and its minor (4Z)-isomer 6'h

Compound **6'h** was prepared from **4** and 4-bromoaniline hydrochloride (**5h**). Yield: 369 mg (79 %) of a pale-grey solid. – M. p. 175–177 °C; **6h**: **6'h** = 85:15. – $[\alpha]_{589}^{20} -17.4$ ($c = 0.35$, CH₂Cl₂). – IR (KBr): $\nu = 3264$, 2980, 2934, 1759, 1725, 1687, 1638, 1588, 1485, 1454, 1368, 1312, 1231, 1152, 999, 960, 820, 775 cm⁻¹. – ¹H-NMR (CDCl₃), *major* (*E*)-isomer **6h**: $\delta = 1.47$ and 1.52 (18H, 2s, 1:1, 2 × ^{*t*}-Bu), 2.55 (1H, ddd, $J = 2.1$, 3.5, 16.1 Hz, 3-Ha), 2.99 (1H, ddd, $J = 2.1$, 10.6, 16.1 Hz, 3-Hb), 4.53 (1H, dd, $J = 3.5$, 10.6 Hz, 2-H), 6.42 (1H, d, $J = 13.4$ Hz, NH), 6.87 (2H, d, $J = 8.8$ Hz, *o*-C₆H₄), 7.39 (2H, d, $J = 8.8$ Hz, *m*-C₆H₄), 7.68 (1H, dt, $J = 2.1$, 13.4 Hz, 4'-H); *minor* (*Z*)-isomer **6'h**: $\delta = 1.53$ (9H, s, ^{*t*}-Bu), 2.60 (1H, ddd, $J = 1.4$, 3.5, 15.4 Hz, 3-Ha), 3.05 (1H, ddd, $J = 1.4$, 10.5, 15.4 Hz, 3-Hb), 4.51 (1H, dd, $J = 3.5$, 10.5 Hz, 2-H), 6.80 (2H, br d, $J = 8.8$ Hz, *o*-C₆H₄), 7.09 (1H, br d, $J = 12.2$ Hz, 4'-H), 7.37 (2H, br d, $J = 8.8$ Hz, *m*-C₆H₄), 9.94 (1H, br d, $J = 12.2$ Hz, NH). – C₂₁H₂₇BrN₂O₅ (467.4): calcd. C 53.97, H 5.82, N 5.99; found C 54.18, H 5.96, N 6.00.

Di-tert-butyl (2S,4E)-4-[(3-nitroanilino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (6i)

Compound **6i** was prepared from **4** and 3-nitroaniline (**5i**) in the presence of one equivalent of hydrochloric acid. Yield: 282 mg (65 %) of a yellow solid. – M. p. 149–152 °C; **6i**: **6i** = 100:0. – $[\alpha]_{589}^{20}$ –16.0 (c = 0.50, CH₂Cl₂). – IR (KBr): ν = 3458, 3285, 3090, 2981, 2935, 1761, 1719, 1688, 1648, 1616, 1588, 1535, 1749, 1369, 1352, 1317, 1238, 1224, 1155, 999, 966, 845, 814, 777, 737, 674 cm^{–1}. – ¹H-NMR (CDCl₃): δ = 1.48 and 1.51 (18H, 2s, 1:1, 2 × *t*-Bu), 2.67 (1H, ddd, J = 2.3, 2.9, 16.4 Hz, 3-Ha), 3.09 (1H, ddd, J = 2.3, 10.5, 16.4 Hz, 3-Hb), 4.56 (1H, dd, J = 3.3, 10.5 Hz, 2-H), 7.07 (2H, d, J = 9.2 Hz, *o*-C₆H₄), 7.39 (1H, br d, J = 13.0 Hz, NH), 7.71 (1H, br dt, J = 2.3, 13.0 Hz, 4'-H), 8.17 (2H, d, J = 9.2 Hz, *m*-C₆H₄). – C₂₁H₂₇N₃O₇ (433.5): calcd. C 58.19, H 6.28, N 9.69; found C 58.34, H 6.44, N 9.73.

Di-tert-butyl (2S,4E)-4-[(4-nitroanilino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (6j)

Compound **6j** was prepared from **4** and 4-nitroaniline (**5j**) in the presence of one equivalent of hydrochloric acid. Yield: 351 mg (81 %) of a pale-grey solid. – M. p. 169–170 °C; **6j**: **6j** = 100:0. – $[\alpha]_{589}^{20}$ –2.1 (c = 0.34, EtOH). – IR (KBr): ν = 3480, 3379, 3260, 3224, 3188, 2977, 1760, 1726, 1686, 1648, 1589, 1506, 1492, 1371, 1315, 1285, 1236, 1224, 1153, 1109, 999, 965, 876, 844, 775, 753, 693 cm^{–1}. – ¹H-NMR (CDCl₃): δ = 1.48 and 1.52 (18H, 2s, 1:1, 2 × *t*-Bu), 2.65 (1H, ddd, J = 2.2, 3.4, 16.5 Hz, 3-Ha), 3.08 (1H, ddd, J = 2.2, 10.5, 16.5 Hz, 3-Hb), 4.56 (1H, dd, J = 3.4, 10.5 Hz, 2-H), 7.08 (2H, dt, J = 2.6, 9.1 Hz, *o*-C₆H₄), 7.18 (1H, br d, J = 13.2 Hz, NH), 7.74 (1H, br dt, J = 2.2, 13.2 Hz, 4'-H), 8.18 (2H, dt, J = 2.6, 9.1 Hz, *m*-C₆H₄). – ¹H-NMR ([D₆]DMSO): δ = 1.54 and 1.55 (18H, 2s, 1:1, 2 × *t*-Bu), 2.72 (1H, br dt, J = 2.4, 17.0 Hz, 3-Ha), 3.18 (1H, ddd, J = 2.4, 10.6, 17.0 Hz, 3-Hb), 4.68 (1H, dd, J = 3.3, 10.6 Hz, 2-H), 7.44 (2H, br d, J = 9.2 Hz, *o*-C₆H₄), 7.79 (1H, br s, 4'-H), 8.26 (2H, br d, J = 9.2 Hz, *m*-C₆H₄), 9.62 (1H, br s, NH). – C₂₁H₂₇N₃O₇ (433.5): calcd. C 58.19, H 6.28, N 9.69; found C 58.12, H 6.45, N 9.62.

Di-tert-butyl (2S,4E)-4-[(2-ethoxy-2-oxoethylamino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (6k) and its minor (4Z)-isomer 6'k

Compound **6'k** was prepared from **4** and ethyl glycinate hydrochloride (**5k**). Yield: 188 mg (47 %) of a colourless solid. – M. p. 128–132 °C; **6k**: **6'k** = 85:15. – $[\alpha]_{589}^{20}$ –20.2 (c = 1.00, CH₂Cl₂). – IR (KBr): ν = 3286, 2979, 2936, 1755, 1728, 1686, 1622, 1458, 1370, 1314, 1257, 1194, 1152, 1094, 1025, 989, 945, 861, 846, 778, 764, 739, 716 cm^{–1}. – ¹H-NMR (CDCl₃), *major (E)isomer 6k*: δ = 1.29 (3H, t, J = 7.1 Hz, CH₃CH₂), 1.47 and 1.51 (18H, 2s, 1:1, 2 × *t*-Bu),

2.42 (1H, ddd, J = 2.0, 3.6, 15.4 Hz, 3-Ha), 2.87 (1H, ddd, J = 2.0, 10.6, 15.4 Hz, 3-Hb), 3.96 (2H, d, J = 5.5 Hz, NHCH₂), 4.23 (2H, q, J = 7.1 Hz, CH₂CH₃), 4.40–4.52 (1H, broad signal, NH), 4.49 (1H, dd, J = 3.6, 10.6 Hz, 2-H), 7.11 (1H, dt, J = 2.0, 13.2 Hz, 4'-H); *minor (Z)isomer 6'k*: δ = 3.85 (1H, dd, J = 1.1, 6.3 Hz, NHCH₂), 4.20 (2H, q, J = 7.2 Hz, CH₂CH₃), 4.44 (1H, dd, J = 4.0, 10.7 Hz, 2-H), 6.46 (1H, br d, J = 12.6 Hz, 4'-H), 7.96 (1H, br dt, J = 6.3, 12.6 Hz, NH). – C₁₉H₃₀N₂O₇ (398.5): calcd. C 57.27, H 7.59, N 7.03; found C 57.54, H 7.78, N 7.33.

Di-tert-butyl (2S,4E)-4-[(3-ethoxy-3-oxopropylamino)methylidene]-5-oxopyrrolidine-1,2-dicarboxylate (6l) and its minor (4Z)-isomer 6'l

Compound **6l** was prepared from **4** and ethyl β-alaninate hydrochloride (**5l**). Yield: 279 mg (68 %) of a pale-yellow solid. – M. p. 85–87 °C; **6l**: **6'l** = 88:12. – $[\alpha]_{589}^{20}$ –20.2 (c = 0.50, CH₂Cl₂). – IR (KBr): ν = 3447, 3291, 3259, 2981, 2935, 1750, 1678, 1626, 1458, 1369, 1317, 1253, 1155, 1081, 1024, 1001, 952, 848, 773 cm^{–1}. – ¹H-NMR (CDCl₃), *major (E)isomer 6l*: δ = 1.27 (3H, t, J = 7.1 Hz, CH₃CH₂), 1.47 and 1.51 (18H, 2s, 1:1, 2 × *t*-Bu), 2.33 (1H, ddd, J = 1.9, 3.7, 15.3 Hz, 3-Ha), 2.56 (2H, t, J = 5.9 Hz, CH₂COOEt), 2.79 (1H, ddd, J = 1.9, 10.7, 15.3 Hz, 3-Hb), 3.46 (2H, deg q, J = 6.0 Hz, NHCH₂), 4.17 (2H, q, J = 7.1 Hz, CH₂CH₃), 4.46 (1H, dd, J = 3.8, 10.7 Hz, 2-H), 4.65 (1H, deg. quintet, J = 6.4 Hz, NH), 7.17 (1H, dt, J = 1.7, 13.6 Hz, 4'-H); *minor (Z)isomer 6'l*: δ = 2.51 (2H, t, J = 6.4 Hz, CH₂COOEt), 4.15 (2H, q, J = 7.1 Hz, CH₂CH₃), 4.41 (1H, dd, J = 4.3, 10.9 Hz, 2-H), 6.56 (1H, br d, J = 12.6 Hz, 4'-H), 7.84–7.96 (1H, broad signal, NH). – C₂₀H₃₂N₂O₇ (412.5): calcd. C 58.24, H 7.82, N 6.79; found C 58.35, H 8.13, N 6.71.

Parallel solution-phase synthesis of (2S,4E)-4-(anilinomethylidene)pyroglutamic acids 7a–c, e–j

The MiniBlock™ was assembled with 12 fritted vessels and charged with compounds **6/6'a–l** (12 × 0.5 mmol) and 2M HCl-EtOAc (12 × 5 mL). The MiniBlock™ was closed and the reaction mixtures were stirred at 20 °C for 12 h. The precipitates were collected by filtration, washed with EtOAc (4 × 3 mL), and dried *in vacuo* at r. t. with P₄O₁₀ for 24 h to give **7a–c, e–j**. Compounds **7d, k, l**, which did not precipitate from the reaction mixtures, were not isolated.

The following compounds were prepared in this manner:

(2S,4E)-4-(Anilinomethylidene)pyroglutamic acid hydrochloride (7a)

Compound **7a** was prepared from **6/6'a**. Yield: 49 mg (39 %) of a light-yellow solid. – M. p. 159–163 °C (partial decomposition above 125 °C). – $[\alpha]_{589}^{20}$ +116.3 (c = 0.52,

MeOH). – IR (KBr): $\nu = 3269, 3236, 3086, 2543, 2473, 1732, 1666, 1595, 1575, 1500, 1323, 1247, 1203, 1086, 924, 804, 754, 719, 687 \text{ cm}^{-1}$. – $^1\text{H-NMR}$ ($[\text{D}_6]$ DMSO): $\delta = 2.77$ (1H, ddd, $J = 2.1, 3.7, 16.8 \text{ Hz}$, 3-Ha), 3.06 (1H, ddd, $J = 2.1, 9.9, 16.7 \text{ Hz}$, 3-Hb), 4.19 (1H, dd, $J = 3.8, 9.9 \text{ Hz}$, 2-H), 6.85 (1H, br t, $J = 7.3 \text{ Hz}$, *p*-C₆H₅), 7.05 (2H, br d, $J = 7.8 \text{ Hz}$, *o*-C₆H₅), 7.05 (2H, br dd, $J = 7.3, 7.8 \text{ Hz}$, *m*-C₆H₅), 7.37 (1H, br d, $J = 11.5 \text{ Hz}$, 4'-H), 8.59 (1H, d, $J = 11.7 \text{ Hz}$, NH), 1-NH₂⁺ and COOH exchanged. – C₁₂H₁₂N₂O₃ · HCl (268.7): calcd. C 53.64, H 4.88, N 10.43; found C 53.52, H 4.83, N 10.15.

(2S,4E)-4-(3-Methylanilinomethylidene)pyroglutamic acid hydrochloride (7b)

Compound **7b** was prepared from **6/6'b**. Yield: 136 mg (96 %) of a colourless solid. – M.p. 169–173 °C (partial decomposition above 140 °C). – $[\alpha]_{589}^{20} +104.8$ ($c = 0.65$, MeOH). – IR (KBr): $\nu = 3269, 2920, 1738, 1665, 1616, 1508, 1309, 1264, 1219, 1171, 1108, 1091, 827, 785, 706 \text{ cm}^{-1}$. – $^1\text{H-NMR}$ ($[\text{D}_6]$ DMSO): $\delta = 2.26$ (3H, s, Me), 2.76 (1H, ddd, $J = 2.1, 3.7, 16.7 \text{ Hz}$, 3-Ha), 3.05 (1H, ddd, $J = 2.1, 9.9, 16.7 \text{ Hz}$, 3-Hb), 4.19 (1H, dd, $J = 3.7, 9.9 \text{ Hz}$, 2-H), 6.68 (1H, br d, $J = 7.4 \text{ Hz}$, *o*-C₆H₄), 6.84 (1H, br t, $J = 8.1 \text{ Hz}$, *p*-C₆H₄), 6.88 (1H, br s, *o*-C₆H₄), 7.12 (1H, t, $J = 7.7 \text{ Hz}$, *m*-C₆H₄), 7.37 (1H, br d, $J = 8.2 \text{ Hz}$, 4'-H), 7.66 (2H, broad signal, 1-NH₂⁺), 8.53 (1H, d, $J = 12.0 \text{ Hz}$, NH), COOH exchanged. – C₁₃H₁₄N₂O₃ · 11/8HCl (287.3): calcd. C 54.35, H 5.31, N 9.75; found C 54.15, H 5.39, N 9.51.

(2S,4E)-4-(4-Methylanilinomethylidene)pyroglutamic acid hydrochloride (7c)

Compound **7c** was prepared from **6/6'c**. Yield: 127 mg (90 %) of a light-yellow solid. – M.p. 162–167 °C (partial decomposition above 130 °C). – $[\alpha]_{589}^{20} +81.6$ ($c = 0.91$, MeOH). – IR (KBr): $\nu = 3268, 3233, 3084, 3037, 2546, 1732, 1665, 1597, 1514, 1326, 1246, 1204, 1087, 984, 812, 718, 661 \text{ cm}^{-1}$. – $^1\text{H-NMR}$ ($[\text{D}_6]$ DMSO): $\delta = 2.21$ (3H, s, Me), 2.74 (1H, ddd, $J = 2.1, 3.8, 16.7 \text{ Hz}$, 3-Ha), 3.04 (1H, ddd, $J = 2.1, 9.9, 16.7 \text{ Hz}$, 3-Hb), 4.17 (1H, dd, $J = 3.8, 9.9 \text{ Hz}$, 2-H), 6.94 (2H, d, $J = 8.4 \text{ Hz}$, *o*-C₆H₄), 7.05 (2H, br d, $J = 8.4 \text{ Hz}$, *m*-C₆H₄), 7.33 (1H, br d, $J = 10.4 \text{ Hz}$, 4'-H), 8.47 (1H, d, $J = 10.4 \text{ Hz}$, NH), 10.32 (1H, br s, COOH), 1-NH₂⁺ exchanged. – C₁₃H₁₄N₂O₃ · 11/6HCl (288.8): calcd. C 54.06, H 5.29, N 9.70; found C 54.15, H 5.39, N 9.51.

(2S,4E)-4-(4-Hydroxyanilinomethylidene)pyroglutamic acid hydrochloride (7e)

Compound **7e** was prepared from **6e**. Yield: 53 mg (37 %) of a colourless solid. – M.p. 153–156 °C (partial decomposition above 130 °C). – $[\alpha]_{589}^{20} +93.0$ ($c = 0.54$, MeOH). – IR (KBr): $\nu = 3263, 2926, 2566, 1740, 1665, 1616, 1605,$

1508, 1321, 1310, 1265, 1234, 1209, 1171, 1091, 827, 785, 691 cm^{-1} . – $^1\text{H-NMR}$ ($[\text{D}_6]$ DMSO): $\delta = 2.70$ (1H, ddd, $J = 2.0, 3.9, 16.5 \text{ Hz}$, 3-Ha), 3.05 (1H, ddd, $J = 2.0, 10.0, 16.5 \text{ Hz}$, 3-Hb), 4.16 (1H, dd, $J = 3.9, 10.0 \text{ Hz}$, 2-H), 6.66 (2H, br d, $J = 8.8 \text{ Hz}$, *o*-C₆H₄), 6.86 (2H, br d, $J = 8.8 \text{ Hz}$, *m*-C₆H₄), 7.29 (1H, br s, 4'-H), 8.34 (1H, br s, NH), 1-NH₂⁺, OH, and COOH exchanged. – C₁₂H₁₂N₂O₄ · HCl (284.7): calcd. C 50.63, H 4.60, N 9.84; found C 50.24, H 4.67, N 9.54.

(2S,4E)-4-(3-Methoxyanilinomethylidene)pyroglutamic acid hydrochloride (7f)

Compound **7f** was prepared from **6/6'f**. Yield: 153 mg (99 %) of a yellowish solid. – M.p. 153–155 °C (partial decomposition above 130 °C). – $[\alpha]_{589}^{20} +106.6$ ($c = 0.65$, MeOH). – IR (KBr): $\nu = 3225, 3068, 2619, 2471, 1731, 1669, 1603, 1513, 1495, 1485, 1322, 1244, 1208, 1197, 1159, 1088, 1050, 985, 804, 766, 720, 681 \text{ cm}^{-1}$. – $^1\text{H-NMR}$ ($[\text{D}_6]$ DMSO): $\delta = 2.75$ (1H, ddd, $J = 2.2, 3.8, 16.7 \text{ Hz}$, 3-Ha), 3.05 (1H, ddd, $J = 2.2, 9.9, 16.7 \text{ Hz}$, 3-Hb), 3.73 (3H, s, OMe), 4.18 (1H, dd, $J = 3.8, 9.9 \text{ Hz}$, 2-H), 6.44 (1H, dd, $J = 2.1, 8.1 \text{ Hz}$, *o*-C₆H₄), 6.59 (1H, t, $J = 2.1 \text{ Hz}$, *o*-C₆H₄), 6.63 (1H, dd, $J = 2.1, 8.1 \text{ Hz}$, *p*-C₆H₄), 7.14 (1H, t, $J = 8.1 \text{ Hz}$, *m*-C₆H₄), 7.33 (1H, br d, $J = 12.3 \text{ Hz}$, 4'-H), 7.72 (2H, br s, 1-NH₂⁺), 8.53 (1H, d, $J = 12.3 \text{ Hz}$, NH), COOH exchanged. – C₁₃H₁₄N₂O₄ · 11/4HCl (307.8): calcd. C 50.72, H 4.99, N 9.10; found C 50.77, H 5.04, N 8.81.

(2S,4E)-4-(3-Bromoanilinomethylidene)pyroglutamic acid (7g)

Compound **7g** was prepared from **6g**. Yield: 152 mg (87 %) of a colourless solid. – M.p. 188–190 °C (partial decomposition above 140 °C). – $[\alpha]_{589}^{20} +85.6$ ($c = 1.03$, MeOH). – IR (KBr): $\nu = 3267, 3233, 2707, 2635, 2471, 1731, 1666, 1592, 1479, 1326, 1246, 1206, 1088, 985, 923, 888, 766, 717, 674 \text{ cm}^{-1}$. – $^1\text{H-NMR}$ ($[\text{D}_6]$ DMSO): $\delta = 2.73$ (1H, ddd, $J = 2.2, 3.7, 16.9 \text{ Hz}$, 3-Ha), 3.03 (1H, ddd, $J = 2.2, 9.8, 16.9 \text{ Hz}$, 3-Hb), 4.15 (1H, dd, $J = 3.7, 9.8 \text{ Hz}$, 2-H), 6.98 (1H, br d, $J = 7.8 \text{ Hz}$, *o*-C₆H₄), 7.04 (1H, br d, $J = 8.0 \text{ Hz}$, *p*-C₆H₄), 7.16 (1H, t, $J = 8.0 \text{ Hz}$, *m*-C₆H₄), 7.22 (1H, br s, *o*-C₆H₄), 7.27 (1H, br d, $J = 12.3 \text{ Hz}$, 4'-H), 7.75 (2H, br s, 1-NH₂⁺), 8.65 (1H, d, $J = 12.3 \text{ Hz}$, NH), COOH exchanged. – C₁₂H₁₁BrN₂O₃ · 9/10HCl (343.9): calcd. C 41.90, H 3.49, N 8.14; found C 41.94, H 3.27, N 8.11.

(2S,4E)-4-(4-Bromoanilinomethylidene)pyroglutamic acid (7h)

Compound **7h** was prepared from **6h**. Yield: 149 mg (86 %) of a light-yellow solid. – M.p. 170–174 °C (partial decomposition above 114 °C). – $[\alpha]_{589}^{20} -23.6$ ($c = 0.45$, MeOH). – IR (KBr): $\nu = 3269, 3231, 3073, 2538,$

1730, 1665, 1587, 1489, 1325, 1244, 1204, 1086, 1075, 984, 819, 718, 652 cm⁻¹. – ¹H-NMR ([D₆]DMSO): δ = 2.75 (1H, ddd, *J* = 2.2, 3.8, 16.9 Hz, 3-Ha), 3.04 (1H, ddd, *J* = 2.2, 9.9, 16.9 Hz, 3-Hb), 4.17 (1H, dd, *J* = 3.8, 9.9 Hz, 2-H), 7.02 (2H, d, *J* = 8.9 Hz, *o*-C₆H₄), 7.28 (1H, br d, *J* = 12.7 Hz, 4'-H), 7.38 (2H, br d, *J* = 8.9 Hz, *m*-C₆H₄), 8.65 (1H, d, *J* = 12.7 Hz, NH), 1-NH₂⁺ and COOH exchanged. – C₁₂H₁₁BrN₂O₃ · HCl (347.6): calcd. C 41.46, H 3.48, N 8.06; found C 41.50, H 3.32, N 7.95.

(2*S*,4*E*)-4-(3-Nitroanilinomethylidene)pyroglutamic acid (7i)

Compound **7i** was prepared from **6i**. Yield: 141 mg (90 %) of a yellow solid. – M. p. 190–194 °C (partial decomposition above 150 °C). – [α]_D²⁰₅₈₉ +52.3 (*c* = 0.78, MeOH). – IR (KBr): ν = 3233, 3072, 2615, 2524, 2445, 1728, 1669, 1621, 1602, 1532, 1483, 1447, 1347, 1328, 1244, 1206, 1087, 986, 930, 814, 796, 739, 720, 663 cm⁻¹. – ¹H-NMR ([D₆]DMSO): δ = 2.79 (1H, ddd, *J* = 2.2, 3.7, 17.0 Hz, 3-Ha), 3.09 (1H, ddd, *J* = 2.2, 9.8, 17.0 Hz, 3-Hb), 4.20 (1H, dd, *J* = 3.7, 9.8 Hz, 2-H), 7.37 (1H, br d, *J* = 12.3 Hz, 4'-H), 7.44–7.55 (2H, m, 2H of C₆H₄), 7.62–7.69 (1H, m, 1H of C₆H₄), 7.84–7.87 (1H, m, 1H of C₆H₄), 9.04 (1H, d, *J* = 12.3 Hz, NH), 1-NH₂⁺ and COOH exchanged. – C₁₂H₁₁N₃O₅ · HCl (313.7):

calcd. C 45.95, H 3.86, N 13.40; found C 45.95, H 3.66, N 13.32.

(2*S*,4*E*)-4-(4-Nitroanilinomethylidene)pyroglutamic acid (7j)

Compound **7j** was prepared from **6j**. Yield: 137 mg (88 %) of a yellow solid. – M. p. 177–180 °C (partial decomposition above 140 °C). – [α]_D²⁰₅₈₉ +90.6 (*c* = 0.63, MeOH). – IR (KBr): ν = 3296, 2920, 2533, 2440, 1725, 1669, 1593, 1556, 1500, 1317, 1242, 1218, 1111, 840, 750, 686 cm⁻¹. – ¹H-NMR ([D₆]DMSO): δ = 2.89 (1H, ddd, *J* = 2.4, 3.5, 17.3 Hz, 3-Ha), 3.17 (1H, ddd, *J* = 2.4, 9.7, 17.3 Hz, 3-Hb), 4.26 (1H, dd, *J* = 3.5, 9.7 Hz, 2-H), 7.29 (2H, br d, *J* = 9.2 Hz, *o*-C₆H₄), 7.43 (1H, br d, *J* = 12.2 Hz, 4'-H), 8.01 (2H, br s, 1-NH₂⁺), 8.17 (2H, br d, *J* = 9.2 Hz, *m*-C₆H₄), 9.44 (1H, d, *J* = 12.2 Hz, NH), COOH exchanged. – C₁₂H₁₁N₃O₅ · HCl (313.7): calcd. C 45.95, H 3.86, N 13.40; found C 46.26, H 4.12, N 13.31.

Acknowledgements

We thank Boehringer Ingelheim Pharma GmbH & Co. KG (Biberach, Germany) for financial support. The financial support from the Slovenian Research Agency through grant P1-0179 is gratefully acknowledged.

- [1] a) S.K. Panday, J. Prasad, D.K. Dikshit, *Tetrahedron: Asymmetry* **2009**, 20, 1581–1632; b) C. Najera, M. Yus, *Tetrahedron: Asymmetry* **1999**, 10, 2245–2303; c) M.B. Smith, *Alkaloids: Chemical and Biological Perspectives* **1998**, 12, 229–287; d) B. Rigo, P. Cauliez, D. Fasseur, F. Sauvage, *Trends in Heterocyclic Chemistry* **1991**, 2, 155–204.
- [2] a) B. Stanovnik, J. Svete, *Chem. Rev.* **2004**, 104, 2433–2480; b) B. Stanovnik, J. Svete, *Synlett* **2000**, 1077–1091; c) B. Stanovnik, J. Svete, *Targets in Heterocyclic Systems* **2000**, 4, 105–137.
- [3] a) J. Svete, *Monatsh. Chem.* **2004**, 135, 629–641; b) J. Svete, *J. Heterocycl. Chem.* **2002**, 39, 437–454; c) J. Svete, *Arkivoc* **2006**, vii, 35–46.
- [4] a) B. Stanovnik, J. Svete, *Mini-Reviews Org. Chem.* **2005**, 2, 211–224; b) S. Pirc, D. Bevk, R. Jakše, S. Rečnik, L. Golič, A. Golobič, A. Meden, B. Stanovnik, J. Svete, *Synthesis* **2005**, 2969–2988; c) Z. Časar, D. Bevk, J. Svete, B. Stanovnik, *Tetrahedron* **2005**, 61, 7508–7519; d) J. Waggener, A. Meden, U. Grošelj, J. Svete, B. Stanovnik, *Tetrahedron* **2008**, 64, 2801–2815; e) J. Waggener, J. Svete, B. Stanovnik, *Synthesis* **2008**, 1436–1442; f) U. Grošelj, A. Meden, B. Stanovnik, J. Svete, *Tetrahedron: Asymmetry* **2008**, 19, 330–342.
- [5] a) J. Baškovč, D. Bevk, B. Stanovnik, J. Svete, *J. Comb. Chem.* **2009**, 11, 500–507; b) D. Kralj, A. Novak, G. Dahmann, U. Grošelj, A. Meden, J. Svete, *J. Comb. Chem.* **2008**, 10, 664–670; c) Č. Malavašič, B. Brulc, P. Čebašek, G. Dahmann, N. Heine, D. Bevk, U. Grošelj, A. Meden, B. Stanovnik, J. Svete, *J. Comb. Chem.* **2007**, 9, 219–229; d) P. Čebašek, D. Bevk, S. Pirc, B. Stanovnik, J. Svete, *J. Comb. Chem.* **2006**, 8, 95–102; e) P. Čebašek, J. Waggener, D. Bevk, R. Jakše, J. Svete, B. Stanovnik, *J. Comb. Chem.* **2004**, 6, 356–362; f) S. Pirc, D. Bevk, S. Golič Grdadolnik, J. Svete, *Arkivoc* **2003**, xiv, 37–48; g) J. Westman, R. Lundin, *Synthesis* **2003**, 1025–1030.
- [6] a) A.N. Bowler, A. Dinsmore, P.M. Doyle, D.W. Young, *J. Chem. Soc., Perkin Trans.* **1997**, 1, 1297–1306; b) A. Dinsmore, P.M. Doyle, D.W. Young, *Tetrahedron Lett.* **1995**, 36, 7503–7506; c) A.N. Bowler, P.M. Doyle, D.W. Young, *J. Chem. Soc., Chem. Comm.* **1991**, 314–316; d) M. Škof, J. Svete, B. Stanovnik, *Heterocycles* **2000**, 53, 339–346; e) U. Uršič, D. Bevk, S. Pirc, L. Pezdirc, B. Stanovnik, J. Svete, *Synthesis* **2006**, 2376–2384.
- [7] a) S. Pirc, D. Bevk, A. Golobič, B. Stanovnik, J. Svete, *Helv. Chim. Acta* **2006**, 89, 30–44; b) D. Kralj, U. Grošelj, A. Meden, G. Dahmann, B. Stanovnik, J. Svete, *Tetrahedron* **2007**, 63, 11213–11222; c) D. Kralj, M. Friedrich, U. Grošelj, S. Kiraly-Potpara, A. Meden, J. Waggener, G. Dahmann, B. Stanovnik, J. Svete, *Tetrahedron* **2009**, 65, 7151–7162.

- [8] a) M. Škof, J. Svete, B. Stanovnik, L. Golič, S. Golič-Grdadolnik, L. Selič, *Helv. Chim. Acta* **1998**, *81*, 2332–2340; b) M. Škof, J. Svete, M. Kmetič, S. Golič-Grdadolnik, B. Stanovnik, *Eur. J. Org. Chem.* **1999**, 1581–1584; c) S. Pirc, S. Rečnik, M. Škof, J. Svete, L. Golič, B. Stanovnik, *J. Heterocycl. Chem.* **2002**, *39*, 411–416; d) M. Škof, S. Pirc, S. Rečnik, J. Svete, B. Stanovnik, L. Golič, L. Selič, *J. Heterocycl. Chem.* **2002**, *39*, 957–963.
- [9] M. Škof, J. Svete, B. Stanovnik, *Acta Chim. Slov.* **1999**, *46*, 567–585.
- [10] M. Besson, F. Delbecq, P. Gallezot, C. Pinel, *Chem. Eur. J.* **2000**, *6*, 949–958.
- [11] R. Jain, *Org. Prep. Proc. Int.* **2001**, *33*, 405–409.
- [12] a) J. J. Titman, J. Foote, J. Jarvis, J. Keeler, D. Neuhaus, *J. Chem. Soc., Chem. Comm.* **1991**, 419–421; b) T. Ando, N. Koseki, R. F. Toia, J. E. Casida, *Magn. Res. Chem.* **1993**, *31*, 90–93; c) P. Fischer, E. Schweizer, J. Langner, U. Schmidt, *Magn. Res. Chem.* **1994**, *32*, 567–568.
- [13] a) U. Grošelj, D. Bevk, R. Jakše, A. Meden, S. Pirc, S. Rečnik, B. Stanovnik, J. Svete, *Tetrahedron: Asymmetry* **2004**, *15*, 2367–2383; b) U. Grošelj, D. Bevk, R. Jakše, A. Meden, B. Stanovnik, J. Svete, *Tetrahedron: Asymmetry* **2005**, *16*, 2927–2945.