


ECONOMETRIC INSTITUTE

ON THE SNAPPER, LIEBLER - VITALE,  
LAM THEOREM ON PERMUTATION  
REPRESENTATIONS OF THE SYMMETRIC GROUP

M. HAZEWINKEL - T. VORST

A handwritten signature in black ink that reads "Erasmus". The signature is written in a cursive style with a large, sweeping initial 'E'.

REPORT 8022/M

ON THE SNAPPER, LIEBLER-VITALE, LAM THEOREM ON PERMUTATION  
REPRESENTATIONS OF THE SYMMETRIC GROUP.

Michiel Hazewinkel and Ton Vorst

ABSTRACT. Let  $\kappa = (\kappa_1, \dots, \kappa_m)$  be a descending partition of  $n$  and  $S_\kappa = S_{\kappa_1} \times S_{\kappa_2} \times \dots \times S_{\kappa_m}$  be the corresponding Young subgroup of  $S_n$ . Denote by  $\rho(\kappa)$  the representation of  $S_n$  which one gets by inducing the trivial representation of  $S_\kappa$ . If  $\lambda = (\lambda_1, \dots, \lambda_m)$  is another partition of  $n$  with  $\sum_{i=1}^r \kappa_i \geq \sum_{i=1}^r \lambda_i$  ( $\forall 1 \leq r \leq m$ ) then  $\rho(\kappa)$  is a subrepresentation of  $\rho(\lambda)$ . In this note we give an elementary complete direct proof of this fact.

Contents

1. Introduction
  2. The Snapper, Liebler-Vitale, Lam Theorem
- References

ON THE SNAPPER, LIEBLER-VITALE, LAM THEOREM ON  
 PERMUTATION REPRESENTATIONS OF THE SYMMETRIC GROUP

Michiel Hazewinkel and Ton Vorst

1. INTRODUCTION.

1. Let  $\kappa = (\kappa_1, \dots, \kappa_m)$ ,  $\kappa_i \in \mathbb{N} \cup \{0\}$ ,  $\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_m \geq 0$  be a descending partition of  $n$ . We identify partitions which differ only by the addition of some additional zero's. An ordering, which we call the specialization order, is defined on the set of all partitions by

$$(1.1) \quad \kappa > \lambda \iff \sum_{i=1}^r \kappa_i < \sum_{i=1}^r \lambda_i, \quad r = 1, 2, \dots$$

The reverse order has been called the dominance order. It occurs in many, seemingly unrelated parts of mathematics [1,2,3], and one of the central occurrences is in the representation theory of the symmetric groups in characteristic zero.

Let  $S_\kappa = S_{\kappa_1} \times \dots \times S_{\kappa_m}$  be the Young subgroup of  $S_n$  ( $S_{\kappa_i}$  is viewed as the permutation subgroup of  $S_n$  permuting the letters  $\kappa_1 + \dots + \kappa_{i-1} + 1, \dots, \kappa_1 + \dots + \kappa_i$ ) corresponding to the partition  $\kappa$  and let  $\rho(\kappa)$  be the representation of  $S_n$  obtained by inducing the trivial representation of  $S_\kappa$  up to  $S_n$ . Also let  $[\kappa]$  be the irreducible representation of  $S_n$  (in characteristic zero) associated to the partition  $\kappa$ . Snapper [5] proved that  $[\kappa]$  occurs in  $\rho(\lambda)$  implies  $\kappa < \lambda$  (this also follows readily from Young's rule) and conjectured the reverse, which he proved for  $m = 2$ . Proofs of the conjecture were given by Liebler-Vitale [4] and Lam [3]. Liebler and Vitale proved more precisely that  $\kappa < \lambda$  implies that  $\rho(\kappa)$  is a subrepresentation of  $\rho(\lambda)$  (which obviously implies, the conjecture because  $[\kappa]$  occurs in  $\rho(\kappa)$ ).

In this note we give a completely elementary direct proof of the Liebler-Vitale result which requires no representation theory at all (beyond the definition of the permutation representations  $\rho(\kappa)$ ) by constructing explicit homomorphisms of representations.

## 2. THE SNAPPER, LIEBLER-VITALE, LAM THEOREM.

### 2.1. Description of the permutation representation $\rho(\kappa)$ .

Let  $W(\kappa)$  be the set of all words of length  $n$  in the symbols  $a_1, \dots, a_m$  such that each  $a_i$  occurs exactly  $\kappa_i$  times. The group  $S_n$  acts in the obvious way on  $W(\kappa)$  ( $\sigma(b_1 \dots b_n) = b_{\sigma(1)} \dots b_{\sigma(n)}$ ,  $\sigma \in S_n$ ) and the vector-space  $V(\kappa)$  with the elements of  $W(\kappa)$  as basis and the action extended linearly is the representation  $\rho(\kappa)$ . We shall denote the elements of  $W(\kappa)$  and the corresponding basis elements of  $V(\kappa)$  with the same symbols.

2.2. "Reduction to the case  $m = 2$ ". It obviously suffices to prove the statement " $\kappa < \lambda \rightarrow \rho(\kappa)$  is a subrepresentation of  $\rho(\lambda)$ " in the case that  $\kappa < \lambda$  and  $\kappa < \mu < \lambda \Rightarrow \kappa = \mu$  or  $\lambda = \mu$ . In this case one easily shows that there exist  $i$  and  $j$ ,  $i > j$  such that  $\lambda_i = \kappa_i + 1$ ,  $\lambda_j = \kappa_j - 1$  and  $\lambda_r = \kappa_r$  for  $r \neq i, j$ . In this case we define a linear map

$$(2.3) \quad \beta_{\lambda, \kappa}: V(\lambda) \rightarrow V(\kappa)$$

by the formula

$$(2.4) \quad \beta_{\lambda, \kappa}(b_1 \dots b_n) = \sum b'_1 \dots b'_n$$

where the sum extends over all words  $b'_1 \dots b'_n$  such that  $b'_t = b_t$  for all but one  $t$ . And for that one  $t$  we have  $b_t = a_i$  and  $b'_t = a_j$ . I.e. the words in the sum on the right are obtained by replacing precisely one occurrence of  $a_i$  by  $a_j$ . This is obviously an  $S_n$ -equivariant map.

We shall prove that  $\beta_{\lambda, \kappa}$  is surjective if (and only if)  $\lambda > \kappa$ . This proves the theorem because the category of  $S_n$ -modules (in characteristic

zero) is semisimple. Alternatively observe that if  $\alpha_{\kappa, \lambda}: V(\kappa) \rightarrow V(\lambda)$  is defined as  $\beta_{\lambda, \kappa}$  with the letters  $a_j$  and  $a_i$  interchanged then  $\alpha_{\kappa, \lambda}$  and  $\beta_{\lambda, \kappa}$  are adjoint to each other in the sense that

$$(2.5) \quad \langle \alpha_{\kappa, \lambda} v, \omega \rangle = \langle v, \beta_{\lambda, \kappa} \omega \rangle, \quad v \in V(\kappa), \omega \in V(\lambda)$$

where the inner products on  $V(\lambda)$  and  $V(\kappa)$  are the ones for which  $W(\lambda)$  and  $W(\kappa)$  form orthonormal bases. This  $\alpha_{\kappa, \lambda}$  is an  $S_n$ -equivariant injection iff  $\beta_{\lambda, \kappa}$  is surjective and it remains to prove that  $\beta_{\lambda, \kappa}$  is surjective if  $\kappa < \lambda$ .

To do this observe that as a vectorspace  $V(\lambda)$  is the direct sum of  $\binom{n}{\lambda} \binom{\lambda_i + \lambda_j - 1}{\lambda_i}$  copies of  $V(\lambda_j, \lambda_i)$  indexed by all words in the symbols  $a_1, \dots, \hat{a}_j, \dots, \hat{a}_i, \dots, a_m, c$  ( $\hat{\phantom{a}}$  denotes deletion) such that  $a_t$  occurs  $\lambda_t$  times and  $c$  occurs  $\lambda_i + \lambda_j$  times. Similarly  $V(\kappa)$  is the direct sum of  $\binom{n}{\kappa} \binom{\kappa_i + \kappa_j - 1}{\kappa_i} = \binom{n}{\lambda} \binom{\lambda_i + \lambda_j - 1}{\lambda_i}$  copies of  $V(\kappa_j, \kappa_i)$  and the homomorphism

(2.4) maps the copies of  $V(\lambda_j, \lambda_i)$  and  $V(\kappa_j, \kappa_i)$ , labelled by the same word in  $a_1, \dots, \hat{a}_j, \dots, \hat{a}_i, \dots, a_m, c$ , into each other and is in fact the direct sum of these induced maps. Hence it is sufficient to prove the surjectivity of  $\beta_{\lambda, \kappa}$  in the case  $m = 2$ .

2.6. Proof of the surjectivity of  $\beta_{\lambda, \kappa}$  in the case  $m = 2$ . Let

$\lambda = (r-1, s+1)$ ,  $\kappa = (r, s)$ ,  $r + s = n$  and write  $x$  for  $a_j$  and  $y$  for  $a_i$ . Then  $W(r-1, s+1)$  consists of words of length  $n$  in  $(r-1)$   $x$ 's and  $(s+1)$   $y$ 's and  $\beta = \beta_{\lambda, \kappa}$  changes such a word into the sum of all words which can be obtained from this word by changing precisely one  $y$  into an  $x$ . E.g.

$$(2.7) \quad \beta(\text{xxxxyy}) = \text{xxxxyy} + \text{xxxxyx} + \text{xxxxyyx}$$

We shall now show that  $\beta$  is surjective if  $r \geq s+1$  (We only need the case  $r \geq s+2$ ). Let  $W = W(r-1, s+1) \cup W(r, s)$ . For each pair  $\omega_1 = b_1 \dots b_n$ ,

$\omega_2 = b_1' \dots b_n'$  in  $W$  we define the distance  $d(\omega_1, \omega_2)$  by

$$(2.8) \quad d(\omega_1, \omega_2) = \# \{t \mid b_t \neq b_t'\}.$$

(This distance is called Hamming distance in coding theory).

Now for  $\omega_0 = x \dots xy \dots y \in W(r, s)$  let

$$(2.9) \quad E_t = \{\omega \in W \mid d(\omega_0, \omega) = t\}$$

Then  $E_t \subset W(r, s)$  if  $t$  is even and  $E_t \subset W(r-1, s+1)$  if  $t$  is odd.

Note that  $\omega \in E_{2t}$  iff there are precisely  $t$   $y$ 's among the first  $r$  letters and  $t$   $x$ 's among the second  $s$  letters and similarly  $\omega \in E_{2t+1}$  iff there are precisely  $t+1$   $y$ 's among the first  $r$  letters of  $\omega$  and  $t$   $x$ 's among the last  $s$  letters.

Now let

$$(2.10) \quad f = r^{-1} \left( c_0 \sum_{\omega \in E_1} \omega + c_1 \sum_{\omega \in E_3} \omega + \dots + c_s \sum_{\omega \in E_{2s+1}} \omega \right)$$

where

$$(2.11) \quad c_t = (-1)^t \binom{r-1}{t}^{-1}$$

We claim that  $\beta(f) = \omega_0$ . To see this observe that since  $\omega \in W(r, s)$  and  $r \geq s+1$  the maximum distance of a  $\omega \in W$  to  $\omega_0$  is  $2s+1$ . Observe that if  $\omega' \in E_{2t+1}$  then  $\beta(\omega')$  is a sum of elements in  $E_{2t}$  and  $E_{2t+2}$  (except when  $t = s$ , then only elements of  $E_{2s}$  can occur by the maximum distance observation).

Now let  $\omega'' = b_1 \dots b_n \in E_{2t}$  ( $t \geq 1$ ) then the coefficient of  $\omega'' \in \beta(f)$  is equal to

$$(2.12) \quad r^{-1} c_t (\# \{i \in \{1, \dots, r\} \mid b_i = x\}) + r^{-1} c_{t-1} (\# \{i \in \{r+1, \dots, r+s\} \mid b_i = x\}) = \\ r^{-1} c_t (r-t) + r^{-1} c_{t-1} t.$$

(The first contribution comes from the elements in  $E_{2t+1}$  whose  $i$ -th element was  $y$  and is transformed to  $x$  to decrease the distance to  $\omega_0$ ; the second contribution comes from elements of  $E_{2t-1}$  whose  $i$ -th element was  $y$  and is transformed to  $x$  to increase the distance).

By definition of  $c_t$  the right-hand side of (2.12) is zero.

The coefficient of  $\omega \in \beta(f)$  is equal to

$$(2.13) \quad r^{-1} c_0 (\# \{i \in \{1, \dots, r\} \mid b_i = x\}) = r^{-1} \cdot 1 \cdot r = 1.$$

This proves that  $\omega_0 = \beta(f) \in \text{Im}\beta$  and hence  $\omega \in \text{Im}\beta$  for all  $\omega \in W(r,s)$  because  $\beta$  is  $S_n$ -equivariant and  $S_n$  acts transitively on  $W(r,s)$ . This concludes the proof.

#### REFERENCES

1. T. Brylawsky, The Lattice of Integer Partitions, *Discrete Math.* 6(1973), 201-219.
2. M. Hazewinkel and C. Martin, Representations of the Symmetric Groups, the Specialization Order, Systems and Grassmann Manifolds, In preparation.
3. T.Y. Lam, Young Diagrams, Schur Functions, the Gale Ryser Theorem and a Conjecture of Snapper, *J. Pure Appl. Algebra* 10(1977) 81-94.
4. R.A. Liebler and M.R. Vitale, Ordering the Partition Characters of the Symmetric Group, *J. Algebra* 25(1973) 487-489.
5. E. Snapper, Group Characters and Nonnegative Integral Matrices, *J. Algebra* 19(1971) 520-535.

Dept. of Mathematics  
 Erasmus University Rotterdam  
 P.O. Box 1738,  
 3000 DR ROTTERDAM. The Netherlands