

Corrigendum

P.W. Hemker, Remarks on sparse-grid finite-volume multigrid,
 Advances in Computational Mathematics 4 (1995) 83–110.

The author is grateful to Christoph Pflaum for pointing out a mistake in the statement and proof of theorem 2.2. The correct version is:

Theorem 2.2

If we consider an expansion of a $C^3(\bar{\Omega})$ -function, u , in piecewise constant functions on the grid Ω_n , for an arbitrary $n \in \mathbb{Z}^3$, $n > \mathbf{0}$, and if we write

$$R_n u = v_0 + \sum_{0 \leq j \leq n} u_j, \tag{1}$$

with $v_0 \in V_0$ and $u_j \in W_j$, $0 \leq j \leq n$, then

$$\|u_j\| \leq 2^{|j|} |u|, \tag{2}$$

and we get an estimate for the approximation error

$$\|u - R_n u\| \leq \frac{1}{3} \sqrt{\frac{2}{3}} (h_1 + h_2 + h_3) |u|. \tag{3}$$

Proof

We take the normalised $\{\tilde{\psi}_j^k\} = \{2^{|j-e|/2} \psi_k^j\}$ as a basis in W_j , $0 \leq j \leq n$, $j \neq \mathbf{0}$. Clearly, all these functions are orthogonal to all functions $v_0 \in V_0$ and mutually they form an orthonormal set in $W_j \subset L^2(\Omega)$. We see further $\psi_k^j \in W_j$ and $\text{support}(\psi_k^j) = \Omega_{j-e,k}$, or, in other words, $\psi_k^j \in V_j$, but ψ_k^j scales like a basis function in V_{j-e} . Hence

$$\int 2^{|j-e|/2} \psi_k^j 2^{|j-e|/2} \psi_m^j d\Omega = 0 \quad \text{for } k \neq m,$$

and

$$\int 2^{|j-e|/2} \psi_k^j 2^{|j-e|/2} \psi_k^j d\Omega = 2^{|j-e|} \int_{\Omega_{j-e,k}} d\Omega = 1.$$

Thus, we find (1) with

$$u_j = \sum_k a_{jk} \tilde{\psi}_k^j = \sum_k (u, \tilde{\psi}_k^j) \tilde{\psi}_k^j.$$

Now

$$a_{jk} = (u, \tilde{\psi}_k^j) = \int_{\Omega} u \tilde{\psi}_k^j d\Omega = \int_{\Omega_{j-e,k}} u \tilde{\psi}_k^j d\Omega.$$

By Taylor expansion around z_k^{j-e} , we have

$$\left| \int_{\Omega_{j-e,k}} u \tilde{\psi}_k^j d\Omega \right| \leq 2^{-2|j|} 2^{|j-e|/2} |u|. \quad (4)$$

For $j \geq e$ the point z_k^{j-e} lies in the interior of Ω and the estimate holds with

$$|u| = \max \left| \frac{\partial^3 u(x)}{\partial x_1 \dots \partial x_3} \right|.$$

For ψ_k^j with a j -component equal to zero, the point z_k^{j-e} lies on the boundary and the function ψ_k^j is constant in one direction over the whole domain Ω , and it is of Haar-wavelet type for the non-zero indices (or index). In this situation the same estimate (4) holds with, e.g. if $j_1 = 0$,

$$|u| = \max \left| \frac{\partial^2 u(x)}{\partial x_2 \dots \partial x_3} \right|.$$

For $j = \mathbf{0}$ the relation (4) is trivially satisfied. Hence, the estimate (4) holds for $j \geq \mathbf{0}$ if we use the seminorm (21), and we find

$$\begin{aligned} |a_{j,k}| &\leq 2^{-3/2} 2^{-3/2|j|} |u|, \\ \|u_j\|^2 &= \sum_k |a_{jk}|^2 \leq \sum_k 2^{-3|j|-3} |u|^2 = 2^{-2|j|-3} |u|^2, \end{aligned}$$

so that

$$\|u_j\| \leq 2^{|j|-3/2} |u|,$$

which leads to (2), and

$$\begin{aligned} \|u - R_n u\|^2 &= \sum_{\substack{j_1 > n_1 \\ \text{or...or} \\ j_3 > n_3}} \|u_j\|^2 \leq \sum_{\substack{j_1 > n_1 \\ j_2 \geq 0 \\ j_3 \geq 0}} + \dots + \sum_{\substack{j_1 \geq 0 \\ j_2 \geq 0 \\ j_3 > n_3}} 2^{-2|j|-3} |u|^2 \\ &\leq 3^{-3} 2(2^{-2n_1} + 2^{-2n_2} + 2^{-2n_3}) |u|^2. \end{aligned}$$

and it follows that

$$\|u - R_n u\| \leq \left(\frac{2}{3^3} \right)^{1/2} (2^{-n_1} + 2^{-n_2} + 2^{-n_3}) |u| = \frac{1}{3} \sqrt{\frac{2}{3}} (h_1 + h_2 + h_3) |u|.$$

□