

Foundations and applications of
Montague grammar
Part 2: Applications to
natural language

T.M.V. Janssen



Centrum voor Wiskunde en Informatica
Centre for Mathematics and Computer Science

1980 Mathematics Subject Classification: 03B65, 08A40, 08A55, 68S10, 03B45, 03B15.
1982 CR Categories: 1.2.7, F.4.2, J.5.

ISBN 90 6196 306 0

Copyright © 1986, Mathematisch Centrum, Amsterdam
Printed in the Netherlands

PREFACE

The present volume is one of the two tracts which are based on my dissertation 'Foundations and applications of Montague grammar'. Volume 1 consists of the chapters 1,2,3 and 10 of that dissertation, and volume 2 of the chapters 4-9. Only minor corrections are made in the text. I would like to thank here again everyone who I acknowledged in my dissertation, in particular my promotor P. van Emde Boas, co-promotor R. Bartsch, and coreferent J. van Benthem. For attending me on several (printing-)errors in my dissertation I thank Martin van de Berg, Cor Baayen, Biep Durieux, Joe Goguen, Fred Landman and Michael Moortgat, but in particular Herman Hendriks, who suggested hundreds of corrections. The illustrations are made by Tobias Baanders.

The two volumes present an interdisciplinary study between mathematics, philosophy, computer science, logic and linguistics. No knowledge of specific results in these fields is presupposed, although occasionally terminology or results from them are mentioned. Throughout the text it is assumed that the reader is acquainted with fundamental principles of logic, in particular of model theory, and that he is used to a mathematical kind of argumentation. The contents of the volumes have a linear structure: first the approach is motivated, next the theory is developed, and finally it is applied. Volume 1 contains an application to programming languages, whereas volume 2 is devoted completely to the consequences of the approach for natural languages.

The volumes deal with many facets of syntax and semantics, discussing rather different kinds of subjects from this interdisciplinary field. They range from abstract universal algebra to linguistic observations, from the history of philosophy to formal language theory, and from idealized computers to human psychology. Hence not all readers might be interested to read everything. Readers only interested in applications to computer science might restrict them selves to volume 1, but then they will miss many arguments in volume 2 which are taken from computer science. Readers only interested in applications to natural language might read chapters 1-3 of volume 1, and all of volume 2, but they will miss several remarks about the connection between the study of the semantics of programming languages and of the semantics of natural languages. Readers familiar with Montague grammar, and mainly interested in practical consequences of the approach, might read chapters 1 and 2 in volume 1 and chapters 6-10 in volume 2, but they will

miss new arguments and results concerning many aspects of Montague grammar.

Each chapter starts with an abstract. Units like theorems etc. are numbered (eg 2.3 Theorem). Such a unit ends where the next numbered unit starts, or where the end of the unit is announced (2.3 end). References to collected works are made by naming the first editor. Page numbers given in the text refer to the reprint last mentioned in the list of references, except in case of some of Frege's publications (when the reprint gives the original numbering).

CONTENTS

iii

V.	The PTQ-fragment	1
	1. Introduction	2
	2. John runs	4
	3. The woman walks	16
	4. Mary walks and she talks	20
	5. John finds a unicorn	25
	6. Every man loves a woman	32
	7. Bill walks in the garden	38
	8. John tries to find a unicorn	42
	9. John believes that Mary will run	47
VI.	Variants and deviations	53
	1. Introduction	54
	2. The use of syntactic information	56
	2.1. Introduction	56
	2.2. Easy to please	56
	2.3. The horse Cannonero	57
	3. Non-polynomially defined operators	58
	3.1. Introduction	58
	3.2. John who runs	59
	3.3. Das Mädchen gibt den Apfel dem Vater	61
	3.4. Woman such that she loves him	62
	4. Operators defined on representants	64
	5. New symbols in IL	68
	5.1. Introduction	68
	5.2. Shake John awake	69
	5.3. I and You	70
	6. Counting elements	71
	6.1. Introduction	71
	6.2. Keenan & Faltz count	71
	6.3. Partee counts	73
	7. The translation language	74
	7.1. Introduction	74
	7.2. Hausser translates	75
	7.3. Lewis translates	76
	7.4. Groenendijk & Stokhof translate	76
	7.5. Keenan & Faltz on translations	77
VII.	Partial rules	79
	1. Restrictions of the framework	80
	2. Partial algebras	83
	2.1. Partial grammars	83
	2.2. Partial models	84
	2.3. Discussion	87
	3. Incorporating transformations	90
	4. Defined for another category	93
	4.1. Introduction	93
	4.2. He_1 is loved	94

4.3. Give John a book	94
4.4. Mary shakes John awake again	95
4.5. See himself	96
4.6. Easy to see	98
5. Subcategorization and rule schemes	99
5.1. Hyperrules	99
5.2. Metarules	102
5.3. Variables	103
6. The Well-formedness constraint	108
VIII. Constituent structures	115
1. Structure-why?	116
2. Theoretical aspects	118
2.1. Trees in Montague grammar	118
2.2. Algebraic considerations	121
2.3. Practical differences	122
3. Technical aspects	125
3.1. Introduction	125
3.2. Operations on trees	126
3.3. Features and lexicon	127
3.4. Queries for information	131
4. PTQ syntax	133
IX. Relative clause formation	137
1. Introduction	138
2. The CN-S analysis	138
2.1. The discussion by Partee	138
2.2. The PTQ-rules	140
2.3. Fundamental problems	142
3. The T-S analysis	144
3.1. Cooper on Hittite	144
3.2. Bach & Cooper on English	145
3.3. Fundamental problems	147
4. The proposals of Cooper	148
4.1. Not-there	148
4.2. Left-over, proposal 1	149
4.3. Left-over, proposal 2	150
4.4. Conclusion	154
5. The variable principle	154
6. Many analyses	158
6.1. The CN-S analysis for English	158
6.2. The S-S analysis for Hittite	159
6.3. The T-S analysis for English	161
6.4. The Det-S analysis for English	163
6.5. Conclusion	164
7. Other arguments	165
7.1. Syntax: gender agreement	165
7.2. Semantics: scope	166
7.3. Conclusion	168
8. The general question	169

X.	Scope ambiguities of tense, aspect and negation	173
	1. Introduction	174
	2. The PTQ-approach	174
	2.1. Introduction	174
	2.2. Syntax of PTQ	175
	2.3. Ambiguities	175
	2.4. Model	177
	3. Basic verb modifiers	177
	4. Compound verb modifiers	184
	5. Complex constructions	187
	5.1. Introduction	187
	5.2. Conjoined verb phrases with positive verbs	187
	5.3. Conjoined verb phrases with negated verbs	189
	5.4. Terms	191
	5.5. Embeddings	192
	6. One of the rules	194
	7. The grammar	199
	7.1. Introduction	199
	7.2. Rules	200
	7.3. Morphology	205
	7.4. Fins and Verb Phrase	205
	7.5. Final remarks	206
	Appendix 1. Individual concepts in PTQ	209
	Appendix 2. Set manipulation in syntax	223
	Index of names	227
	References	229

CHAPTER V

THE PTQ-FRAGMENT

ABSTRACT

In this chapter the fragment of English described in Montague's article PTQ (MONTAGUE 1973) is presented. The method of exposition consists in starting with a very small fragment, and expanding it gradually. In each stage both the syntax and the semantics are discussed extensively. Special attention is paid to the motivation and justification of the analysis.



1. INTRODUCTION

The aim of this chapter is to present in a rigorous way the syntax and the semantics of a certain fragment of a certain dialect of English. The fragment is about the same as the one presented in MONTAGUE (1973), henceforth PTQ. On all essential points I will follow the treatment given in PTQ, in the details, however, there are some differences. The presentation, motivation and justification I will give for the treatment, differs considerably from PTQ. For the presentation I will employ a method which might be called 'concentric'. I will start with a very small fragment, and gradually expand this. For the fragments in each of the stages both the syntax and semantics are given, together with an extensive discussion. I hope that this method will make it easier to understand the sometimes difficult or subtle details of the PTQ-treatment. Certain details (concerning the problems of extension and intension) will be discussed in appendix 1 of this book. A list of the rules of the fragment (useful as a survey) can be found in chapter 8.

In the exposition I will give special attention to algebraic and algorithmic aspects of the treatment. The algebraic considerations often provide an explication why a certain detail is as it is, and not otherwise. The algorithmic aspect concerns the method to obtain simple meaning representations. I do not like some rather abstract relation between a sentence and its meaning. For instance, I am not satisfied with a two-lines-long formula, if there is a one-line-long-formula which represents the same meaning, and if a meaning is represented by a formula which has to be interpreted in models satisfying certain meaning postulates, I would like to have a formula in which these postulates are made explicit. So I prefer concise and clear meaning representations. In order to reach this aim, several rules will be given for the reduction of formulas.

The syntax of the fragment in PTQ is treated rather poorly. In this chapter only minor improvements will be given (for more fundamental changes see chapter 8). But syntax was not Montague's main interest; he was interested primarily in semantics. The fragment is rich in semantically interesting phenomena, and it deals with several famous semantic puzzles. Below I will mention some of the sentences dealt with, together with some comments.

A first kind of phenomena dealt with concerns sentences of which it is clear what their meanings are, and how these should be represented using standard predicate logic. Their challenge lies in the aim to obtain these

meanings in a systematic way. Consider (1) and (2).

(1) *John runs.*

(2) *Every man runs.*

These two sentences are closely related in form: only the subject differs. Therefore one would like to produce the sentences along the same lines. The representations of their meanings, however, are rather different. In standard logic it would be as in (3) and (4).

(3) $run(john)$

(4) $\forall x[man(x) \rightarrow run(x)]$.

This gives rise to the question how to obtain rather divergent formulas from closely related sentences. A corresponding question arises for the ambiguity of (5).

(5) *Every man loves a woman.*

This sentence may be used when one specific woman is loved by every man, (say Brigitte Bardot), or when for each man there may be another woman (say his own mother). Sentence (5) is considered as being ambiguous between these two possibilities (for arguments, see section 6). This kind of ambiguity is called 'scope ambiguity' (of quantifiers). The two readings that will be obtained for (5) are (simplified) represented in (6) and (7).

(6) $\forall x[man(x) \rightarrow \exists y[woman(y) \wedge love(x,y)]]$

(7) $\exists y[woman(y) \wedge \forall x[man(x) \rightarrow love(x,y)]]$.

A second kind of phenomena dealt with concerns sentences for which it is difficult to say how their meanings should be represented. Consider (8) and (9)

(8) *John seeks a unicorn.*

(9) *John finds a unicorn.*

These two sentences have about the same form, only the verbs they contain are different. One is tempted to expect that they have about the same meanings as well; the only difference being that they express another relation between John and a unicorn. This is not the case, however. The one sentence gives information about the existence of unicorns, which the other sentence does not. So an approach which says that the *seek*-relation is always a relation between two individuals would not be acceptable. We have

to provide a meaning for (8) from which it does not follow that unicorns exist. However, sentence (8) can also be used in a situation that unicorns exist, and it is ambiguous between these two possibilities. It has a reading from which it follows that at least one unicorn exists (the referential reading), and a reading from which this does not follow (the non-referential reading).

Some examples of the referential/non-referential ambiguity are (10), (11), and (12).

(10) *John talks about a unicorn.*

(11) *John wishes to find a unicorn and eat it.*

(12) *Mary believes that John finds a unicorn and that he eats it.*

Sentence (9) allows only for a referential reading. The same holds for sentence (13), see MONTAGUE 1973, p.269.

(13) *John tries to find a unicorn and wishes to eat it.*

The ambiguity we distinguish in sentences (8), (10), (11) and (12) is in the literature also called the '*de-dicto/de-re*' ambiguity, or the '*specific/non-specific*' ambiguity. This terminology is not felicitous, because one might associate with it a distinction that is not covered by the formal analysis that will be provided. Nevertheless, this terminology will sometimes be used in the sequel, since it is standard for some of the examples.

2. JOHN RUNS

The fragment in this section consists of very simple sentences like *John runs*. It has three categories (=sorts): the category T of terms, the category IV of intransitive verb phrases, and the category S of sentences (in PTQ a *t* is used instead of S). There are basic expressions (=generators) of the categories T and IV. The set B_T of generators of the category T contains the proper names of the PTQ-fragment, ($B_T \sim$ 'Basic expressions of category T'). Furthermore a special name is added for illustrative purposes: *Bigboss*. The sets B_T and B_{IV} are defined as follows (B_T will be extended in section 4).

2.1. $B_T = \{John, Bill, Mary, Bigboss\}$

2.2. $B_{IV} = \{run, walk, talk, rise, change\}.$

2.2. END

In the logic there is for each element of $B_{\mathbb{T}}$ a corresponding constant of type e , except for *Bigboss*. In PTQ these constants are called j, m, b respectively, but I will use full names: *john* etc.. Notice the difference in the letter type used for English (*Mary*), and the one used for logic (*mary*). One might expect that a proper name translates into the corresponding constant, but for reasons to be explained later, the translation is a complex expression containing this constant. So among the constants in IL of type e , we distinguish three special ones.

$$2.3 \quad \{john, bill, mary\} \subset \text{CON}_e.$$

2.3. END

Constants of type e get as interpretation (with respect to a point of reference) some element in the domain of individuals. This interpretation has to be restricted, for the following reason. If we will speak tomorrow about John, then we will mean the same individual as today (although he may have some other properties). For instance, if the world would have been different, say, if the Mount Everest would not be the highest mountain, then John would still be the same individual (although his opinion about the Mount Everest might be different). This conception about the individual corresponding with a proper name is expressed by the phrase 'proper names are *rigid designators*'. For an extensive discussion of this conception, see KRIPKE 1972. This idea will be incorporated in our semantics by interpreting constants like *john* 'rigidly', i.e. for each index it will denote the same individual. The name *Bigboss* is to be understood as a surname of the most powerful individual on earth. Since this will not always be the same individual, *Bigboss* is not treated as a rigid designator of type e .

The constants of intensional logic are not interpreted rigidly, on the contrary, they are interpreted index-dependent. I recall the clause for the interpretation of constants:

$$c^{A, i, g} = F(c)(i) \quad (c \in \text{CON}).$$

This means there is no guarantee that the constants corresponding with the proper names of PTQ are interpreted rigidly. Therefore not all possible models for the interpretation of IL are reasonable candidates for an interpretation of English. We will consider only those models in which the

constants *john*, *bill*, and *mary* are interpreted rigidly. This is formalized as follows. The requirement of 'rigidity' is expressed by means of an IL-formula, and we will consider only those models in which this formula holds. The formula is called a Meaning Postulate (an MP). It bears index 1 because it is the first meaning postulate in PTQ. Notice that this postulate describes in fact a collection of three formulas.

2.4. Meaning postulate 1:

$$\exists u \Box [u=\alpha] \quad \text{where } \alpha \in \{john, bill, mary\}.$$

2.4. END

This meaning postulate requires that there is one individual in the semantic domain such that the interpretation of *john* equals that individual for all indices. For the PTQ fragment this postulate may be considered sufficient. In more subtle situations this formalization of rigidity is probably too absolute. If epistemological verbs like *know* or *believe* are analysed in detail, then the notion of rigidity may have to be weakened to something like 'in all worlds compatible with the beliefs of some individual such a constant is rigid'. I will, however, follow the PTQ formalization.

An important technical consequence of MP1 is that lambda-conversion is allowed when one of the constants *john*, *bill* or *mary* occurs as argument. First I recall the notation for substitution, for a formal definition see chapter 3, definition 4.3.

2.5. DEFINITION. $[\alpha/z]\phi$ denotes the result of *substitution* of α for all free occurrences of z in ϕ .

2.6. THEOREM.

$$\models \lambda u[\phi](\alpha) = [\alpha/u]\phi$$

where

$$\alpha \in \{john, bill, mary\}.$$

PROOF. MP1 says that for all i, g : $i, g \models \exists u \Box [u=\alpha]$
 so there is a $g' \underset{u}{\sim} g$ such that : $i, g' \models \Box [u=\alpha]$
 hence for all j : $j, g' \models u = \alpha$

Let i_1 and i_2 be arbitrary. Then:

$$\begin{aligned} V_{i_1 g}(\alpha) &= V_{i_1, g'}(\alpha) = V_{i_1, g'}(u) = g'(u) = V_{i_2, g'}(u) = V_{i_2, g'}(\alpha) = \\ &= V_{i_2, g}(\alpha). \end{aligned}$$

This means that the condition of theorem 6.4 from chapter 3 is satisfied, hence the theorem allows us to apply λ -conversion.

2.6. END

In the sequel λ -conversion will be used frequently for reducing a formula to a simpler form. Besides λ -conversion several other rules will be introduced for this purpose; they are called reduction rules (RR's). Together they will constitute a procedure which simplifies the formulas obtained by translating the expressions of the fragment. For each reduction rule a correctness proof has to be given, i.e. a proof that the rule transforms a formula into a logically equivalent one. Theorem 6.1 from chapter 3 then allows us to reduce a formula as soon as it is obtained. The purpose of the reduction rules is to obtain formulas which express the intended meaning as clearly and simply as possible. The rules presented in this chapter are almost identical with the rules presented in JANSSEN 1980a. Related reduction rules are discussed in FRIEDMAN & WARREN 1980a,b and INDURKHYA 1981; these authors use the reduction rules for a somewhat different purpose (e.g. to obtain the most extensionalized form), and therefore there are some differences.

The first reduction rule concerns λ -conversion. With respect to this rule the following class of formulas is important: the formulas which contain no operators \forall , H , or W , and which contain as constants only *john*, *mary* or *bill*. Extending definition 6.2 from chapter 3, I will call these expressions modally closed, since they have the same properties with respect to λ -conversion.

2.7. DEFINITION. An IL formula is called *modally closed* if it is an element of the IL-subalgebra:

$$\langle \{john, mary, bill\}, (VAR_{\tau})_{\tau \in Ty}, R \cup \{R_{\wedge}, R_{\square}\} \rangle$$

where R consists of the operators of Ty_2 (recall that R_{\wedge} and R_{\square} indicate prefixing with \wedge and \square respectively).

2.8. Reduction rule 1

Let $z \in \text{VAR}_{\tau_1}$, $\alpha \in \text{ME}_{\tau_2}$, and $\beta \in \text{ME}_{\tau_2}$.

Then replace $\lambda z[\beta](\alpha)$ by $[\alpha/z]\beta$ if

- 1) no variable in α becomes bound by substitution of α for z in β and either
- 2) no occurrence of z in β lies within the scope of \wedge, H, W or \square
- or
- 3) α is modally closed.

CORRECTNESS PROOF

The difference between this rule and theorem 6.3 from chapter 3 is that condition 3 allows for the occurrence of the rigid designators *john*, *mary* and *bill*. Hence if conditions 1) and 2) are satisfied, the correctness of the λ -conversion follows from that theorem. Suppose now that conditions 1) and 3) are satisfied, and consider the case that α contains of the constants *john*, *bill* and *mary* only occurrences of *john*.

Let w be a variable which does not occur in α or β , and let α' and β' be obtained from α and β by substitution of w for *john*. Consider now

$$(A) \quad \lambda w[\lambda z[\beta'](\alpha')](john).$$

Since α' and β' do not contain occurrences of *john* the old conditions for λ -conversion on z are satisfied (chapter 3, theorem 6.3). So (A) is equivalent with:

$$\lambda w[[\alpha'/z]\beta'](john).$$

From theorem 2.6 above, it follows that λ -conversion on w is allowed, so this formula is equivalent with

$$[john/w][[\alpha'/z]\beta'].$$

By the definition of substitution, this is equivalent with

$$[\alpha/z]\beta.$$

So (A) is equivalent with this last formula. On the other hand, we may perform in (A) λ -conversion on w because the condition of theorem 2.6 is satisfied. So (A) is also equivalent with

$$\lambda z[\beta](\alpha).$$

The combination of these last two, with (A) equivalent, formulas proves the correctness of λ -conversion for the case that conditions 1) and 3) are satisfied, and that α contains only occurrences of *john*. For other constants and for occurrences of more than one constant, the proof proceeds analogously.

2.8. END

As said before, at different indices different persons can be Bigboss. Therefore we cannot translate *Bigboss* into a rigid constant of type e . We might translate it into a constant of type $\langle s, e \rangle$, or into a constant of type e and interpret it non-rigidly. I choose the former approach (thus being consistent with the examples involving *bigboss* given in section 7 of chapter 3). This explains the following definition

$$2.9 \quad \textit{bigboss} \in \text{CON}_{\langle s, e \rangle}.$$

2.9. END

The interpretation of the constant *bigboss* is a function from indices to individuals. Such a function is called an *individual concept*. Also $\hat{\textit{john}}$ denotes an individual concept. The individual concept denoted by $\hat{\textit{john}}$ is a constant function, whereas the one denoted by *bigboss* is not. One might expect that *Bigboss* translates into the corresponding constant. But, as for the other proper names, it will be explained later why this is not the case.

Suppose that the balance of power changes and Bresjnev becomes Bigboss instead of Reagan. Then this might be expressed by sentence (14).

(14) *Bigboss changes.*

The meaning of (14) is not correctly represented by a formula which says that the predicate *change* applies to a certain individual. Who would that be? Maybe there was a change in the absolute power of Reagan (it decreased), or in the absolute power of Bresjnev (it increased). Probably both persons changed with respect to power. Sentence (14) rather says that the concept

'Bigboss' has changed in the sense that it concerns another person. So the meaning of (14) can be represented by a formula which says that the predicate *change* holds for the individual concept related with *Bigboss*. In such an analysis *change* has to be of type $\langle\langle s,e \rangle, t \rangle$. Due to the homomorphic relation between syntax and semantics, this means that all intransitive verbs have to be of type $\langle\langle s,e \rangle, t \rangle$.

At this stage of the description of the fragment the only example of an argument of type $\langle s,e \rangle$ is the artificial example *bigboss*. In appendix 2 of this book, other examples will be given where the translation of the argument of a property has to be of this type. This discussion explains the introduction of the following constants and translations. The translation function is indicated by means of a '(prime). Note that this is a different use of ' than in PTQ (there it distinguishes English words from logical constants).

2.10 $\{run, walk, talk, rise, change\} \subset \text{CON}_{\langle\langle s,e \rangle, t \rangle}$

2.11 $run' = run, walk' = walk, talk' = talk$

$rise' = rise, change' = change.$

2.11. END

One might be tempted to take the constant *john* as translation of the proper name *John*. In the fragment consisting only of sentences like *John runs* there would be no problem in doing so. But there are more terms, and the similarity of syntax and semantics requires that all terms are translated into expressions of the same type. We already met the proper name *Bigboss*, translating into an expression of type $\langle s,e \rangle$. One might expect \hat{john} as translation for *John*. But in the sequel we will meet more terms: e.g. *every man*. If we would translate *John* into an expression denoting an individual concept (or alternatively an individual), then *every man* has to be translated into such an expression as well. Would that be possible?

The idea is discussed by LEWIS (1970). He tells us that in the dark ages of logic a story like the following was told. 'The phrase every pig names a [...] strange thing, called the universally generic pig, which has just those properties that every pig has. Since not every pig is dark, pink, grey or of another color, the universally generic pig is not of an any color (Yet neither he is colorless, since not every - indeed not any - pig is colorless)'. (LEWIS 1970, p.35). This illustrates that this approach is not sound. Therefore, we will forget the idea of universal generic

objects (for a proposal for a reconstruction, see Van BENTHEM 1981a), and we will interpret the term *every man* as the set of properties every man has. As a consequence of the similarity of syntax and semantics, all other terms will denote sets of properties as well.

On the basis of this argumentation one might expect for *John* the translation $\lambda Z[Z(\wedge^{John})]$, where Z is a variable of type $\langle\langle s, e \rangle, t \rangle$. But this is not adequate for the following reason. A variable of type $\langle\langle s, e \rangle, t \rangle$ denotes (the characteristic function of) a set of individual concepts. What we usually take to be a property cannot be adequately formalized in this way. Consider the property 'being a football player'. This would be formalized as a set of individual concepts. The same holds for the property of 'being a member of the football union': this is formalized as a set of individual concepts as well. Suppose now that (for a certain index) all football players are members of the football union. Then these two sets would be the same, so the two properties would be formalized in the same way. But we do not consider these two properties as being the same. In other circumstances (for other indices) there might be players who are not a member of the union. In order to formalize these differences, properties are taken to be of one intensional level higher hence a variable which ranges over properties has to be of type $\langle s, \langle\langle s, e \rangle, t \rangle \rangle$. This explains the following translations of proper names.

2.12. Translations

$$John' = \lambda P[[^V P](\wedge^{John})], \quad Bill' = \lambda P[[^V P](\wedge^{Bill})]$$

$$Mary' = \lambda P[[^V P](\wedge^{Mary})], \quad Bigboss' = \lambda P[[^V P](bigboss)]$$

here $P \in VAR_{\langle s, \langle\langle s, e \rangle, t \rangle \rangle}$.

2.12. END

After this discussion concerning the proper names and intransitive verbs, the rule for their combination can be given. I first quote the PTQ formulation, since this way of presentation is in the literature the standard one. The formulation of the rule contains expressions like ' $\alpha \in P_T$ ', this should be read as ' α is a phrase of the category T '. The rule is called S_4 , because it is the fourth syntactic rule of PTQ, and I wish to follow that numbering when possible.

2.13. Rule S_4

If $\alpha \in P_T$ and $\beta \in P_{IV}$ then $F_4(\alpha, \beta) \in P_S$, where $F_4(\alpha, \beta) = \alpha\tilde{\beta}$ and $\tilde{\beta}$ is the result of replacing the first verb in β by its third person singular present.

2.13. END

This formulation of the rule contains a lot of redundancy, and therefore I will use a more concise presentation. As one remembers from the previous chapters, the syntactic rules are operators in an algebraic grammar. The form of representation I will use, resembles closely the representations used in the previous chapters for algebraic operators. First it will be said what kind of function the rule is; as for S_4 it is a function from $T \times IV$ to S (written as $T \times IV \rightarrow S$). Next it will be described how the effect of the operator is obtained. I will use a notation that suggests that some basic operations on strings are available, in particular a concatenation operator which yields the concatenation of two strings as result. The semi-colon (;) is used to separate the consecutive stages of the description of the syntactic operator; it could be read as 'and next'. Furthermore the convention is used that α always denotes the expression which was the first argument of the syntactic rule. If this expression is changed in some step of the syntactic operation, it will then denote the thus changed expression. For the second argument β is used in the same way. Rule S_4 presented in this format reads as follows.

2.14. Rule S_4

$$T \times IV \rightarrow S$$

F_4 : replace the first verb in β by its third person singular present;
concatenate (α, β) .

2.14. END

The occurrence of the name F_4 is a relict of the PTQ formulation, and might be omitted here. But in a context of a long list of rules it is sometimes useful to have a name for an operation on strings, because it can

then be used in the description of other rules.

The translation rule corresponding with S_4 reads in PTQ as follows.

2.15. T_4 :

If $\alpha \in P_T$, $\beta \in P_{IV}$, and α, β translate into α', β' respectively, then $F_4(\alpha, \beta)$ translates into $\alpha'(\wedge\beta')$.

2.15. END

Also the translation rule contains a lot of redundant information.

Let us denote by α' the translation of the first, and by β' the translation of the second argument of the preceding syntactic rule. Then a translation rule can fully be described by giving just the relevant logical expression (polynomial over IL with α' and β' as parameters). What the types of α' and β' are, follows immediately from the sorts mentioned in the syntactic rule T_4 presented in this format reads:

2.16. T_4 :

$\alpha'(\wedge\beta')$

2.16. END

Now we come to the production of sentence (15), viz. *Bigboss changes*. This sentence, containing the artificial term *Bigboss*, is given as the first example because all information needed for a full treatment of this sentence is given now; sentences like *John changes* have to wait for a moment. Sentence (15) is obtained by application of S_4 to the basic term *Bigboss* and the basic verb *change*. This information is presented in the tree in figure 1. The S in {S,4} stands for the category of the obtained expression, the 4 for the number of the rule used to produce the expression (15) *Bigboss changes*.




Figure 1

The translation of *Bigboss* is $\lambda P[\vee P(\text{bigboss})]$, and the translation of *change* is *change*. If we combine *Bigboss* and *change* according to rule S_4 , thus producing (15), then the translation of the result is obtained by application of T_4 to their respective translations. Since

$T_4(\alpha', \beta') = \alpha'(\wedge \beta')$, sentence (15) translates into (16).

(16) $\lambda P[\vee P(\text{bigboss})](\wedge \text{change})$.

Now conditions 1 and 2 of reduction rule RR_1 are satisfied. So this formula can be reduced to (17).

(17) $[\vee \wedge \text{change}](\text{bigboss})$.

This formula can be simplified further using the following reduction rule.

2.17. Reduction Rule 2

Let be given a formula of the form $\vee \wedge \alpha$. Then replace this formula by α .

CORRECTNESS PROOF. $\models \vee \wedge \alpha = \alpha$ see chapter 3, theorem 7.1.

2.17. END

Using this reduction rule formula (17) reduces to (18).

(18) $\text{change}(\text{bigboss})$.

This formula expresses that the predicate *change* holds for the individual concept *bigboss*.

Instead of all this verbosity, we might present the translations immediately in the tree. Depending on the complexity of the formulas involved, these may be unreduced, partially reduced or completely reduced formulas. An example is given in figure 2.




Figure 2

Another method to present the production and translation process is to write this in an algebraic way, of which the following is an example.

$$\begin{aligned}
 [\text{Bigboss changes}]' &= [S_4(\text{Bigboss}, \text{change})]' = \\
 &= T_4(\text{Bigboss}', \text{change}') = \text{Bigboss}'(\wedge \text{change}') = \\
 &= [\lambda P[\vee P(\text{bigboss})](\wedge \text{change})] = \{RR_1\} = [\vee \wedge \text{change}](\text{bigboss}) = \{RR_2\} = \\
 &= \text{change}(\text{bigboss})
 \end{aligned}$$

The treatment of *Mary walks* proceeds, in its first stage, analogously to the treatment of *Bigboss changes*, see figure 3.




Figure 3

The formula obtained as translation for *Mary walks*, is not completely satisfactory. Intuitively one interprets this sentence as stating that a certain predicate (denoting the property of walking) holds for a certain individual (Mary). This is not reflected in the obtained translation; in $\text{walk}(\hat{\text{mary}})$ a predicate is applied to an individual concept. Since *mary* is a rigid constant, $\hat{\text{mary}}$ denotes a function which yields for all indices the same individual. Saying that this constant function has a certain property is tantamount to saying that the corresponding individual has a certain property (there is a 1-1 correspondence between individuals and functions yielding always the same individual). However, one would like to have reflected in the translation of *Mary walks* that a predicate holds for an individual. Therefore the following notation is introduced (see PTQ, p.265).

2.18. DEFINITION. Let $\delta \in \text{CON}_{\langle\langle s, e \rangle, t \rangle}$. Then δ_* is an abbreviation for $\lambda u \delta(\hat{u})$ (so $\delta_* \in \text{ME}_{\langle e, t \rangle}$).

2.18. END

Consequently we have the following rule for simplifying formulas.

2.19. Reduction rule 3

Let be given a formula of the form $\delta(\hat{\alpha})$, where $\delta \in \text{CON}_{\langle\langle s, e \rangle, t \rangle}$ and $\alpha \in \text{VAR}_e$ or $\alpha \in \{\text{john, bill, mary}\}$. Then replace $\delta(\hat{\alpha})$ by $\delta_*(\alpha)$.

CORRECTNESS PROOF. $\delta_*(\alpha) = \lambda u [\delta(\hat{u})](\alpha) = \{\text{RR}_1\} = \delta(\hat{\alpha})$. Note that λ -conversion is allowed because the mentioned constants of type *e* are rigid designators.

2.19. END

Using RR3 the translation of *Mary walks* in figure 3, reduces to (19).

(19) $walk_*(mary)$.

As last example I present the treatment of the sentence mentioned in the title of this section. For variation I use not the tree representation, but the algebraic one.

$$\begin{aligned} [John\ runs]' &= [S_4(John, run)]' = John'(\wedge run') = \lambda P[[^V P](\wedge john)](\wedge run) = \\ &= \{RR_1\} = [{}^V \wedge run](\wedge john) = \{RR_2\} = run(\wedge john) = \{RR_3\} = run_*(john). \end{aligned}$$

In PTQ more is said about the fragment presented so far. A meaning postulate (MP₃) is introduced which says that the truth of e.g. $walk(x)$ only depends on the extension of x , i.e. the subject position of $walk$ is extensional. In appendix 2 of this book the problems of extension and intension will be discussed, and this postulate will be considered. For verbs of other categories the extensionality of the subject position is guaranteed by meaning-postulates as well, (see appendix 1).

3. THE WOMAN WALKS

In this section the fragment is extended with the categories of Common Nouns (CN) and of determiners (Det). The treatment of determiners given here differs from their PTQ treatment. In PTQ determiners are introduced syncategorematically, introducing each determiner by a distinct rule. Maybe the motivation for Montague to do so, was that in logic quantifiers are usually introduced syncategorematically. From a linguistic point of view it is more attractive to have determiners in a separate category (they form a group of expressions which behave syntactically in a regular way). Since I do not know any argument against treating them categorially, the PTQ approach is not followed here. The generators of the two new categories are as follows

3.1. $B_{CN} = \{man, woman, park, fish, pen, unicorn, price, temperature\}$

3.2. $B_{Det} = \{every, a, the\}$

3.2. END

For each element in B_{CN} there is a corresponding constant, and the common nouns translate into these constants. The nouns are treated semantically in the same way as the intransitive verbs we have met in section 2. Hence the nouns translate into constants of type $\langle\langle s, e \rangle, t \rangle$. This

explains the following definitions.

- 3.3. $\{man, woman, park, fish, pen, unicorn, price, temperature\} \subset CON_{\langle\langle s, e \rangle, t \rangle}$
- 3.4. $man' = man, woman' = woman, park' = park, pen' = pen,$
 $unicorn' = unicorn, price' = price, temperature' = temperature.$
- 3.4. END

An example of a formula containing the constant *bill* is (20), in which is expressed that Bill is a man.

(20) $man(\wedge bill).$

The δ_* -notation (definition 2.18) is applicable to all constants of type $\langle\langle s, e \rangle, t \rangle$, so it can be applied to constants translating common nouns as well. So (20) may be replaced by (21).

(21) $man_*(bill).$

Out of a CN and a determiner a term can be formed, using the following rule.

3.5. Rule S_2

$Det \times CN \rightarrow T$

F_2 : concatenate(α, β)

T_2 : $\alpha'(\wedge \beta').$

Example

$F_2(a, woman) = a\ woman.$

3.5. END

We wish to use the terms produced with this rule in the same way as we used the term *John*: rule S_4 should be applicable to the result of S_2 , yielding sentences like (22), (23) and (24).

(22) *A woman runs*

(23) *Every woman runs*

(24) *The woman runs.*

The meanings associated with determiners are best understood by considering the meanings that we wish to assign to the above sentences (cf. the discussion concerning contextuality and compositionality in section 2 of chapter 1). Let us accept (for the moment without explanation) the quantification over individual concepts; then the translations of (22),

(23) and (24) are (25), (26) and (27) respectively.

(25) $\exists x[\text{woman}(x) \wedge \text{run}(x)]$

(26) $\forall x[\text{woman}(x) \rightarrow \text{run}(x)]$

(27) $\exists x \forall y[[\text{woman}(y) \leftrightarrow x=y] \wedge \text{run}(x)]$.

The last formula is somewhat complex. It says that there is an entity x which is a woman, and that for any entity y which is a woman holds that it is identical to the entity x . In other words, (27) is false when there is no woman at all, and it is false when there is more than one woman. This kind of analysis for *the* is called the Russellian analysis, because it was proposed by Russell to deal with the famous example (28).

(28) *The present King of France is bald.*

The meanings of the terms have to be such that if they take an IV-translation as argument, the resulting translations are the ones we desired for the obtained sentences. Hence their translations have to be of the same kind as the translation of the term *John*: a (characteristic function of a) set of properties of an individual concept. So we wish to translate (29) by (30).

(29) *a woman*

(30) $\lambda P \exists x[\text{woman}(x) \wedge \overset{V}{P}(x)]$.

Formula (30) is interpreted as the characteristic function of those properties P such that there is at least one woman which has this property P . Other determiners are treated analogously. As translation for the determiner a we take formula (30), but with *woman* replaced by a variable. This variable is of type $\langle s, \langle \langle s, e \rangle, t \rangle \rangle$ (the reason for this is the same as the reason given for the type of the variable P , see the translation of *John*). This explains the following translations of determiners.

3.6. Translations of determiners

every' = $\lambda Q \lambda P \forall x[\overset{V}{Q}(x) \rightarrow \overset{V}{P}(x)]$

a' = $\lambda Q \lambda P \exists x[\overset{V}{Q}(x) \wedge \overset{V}{P}(x)]$

the' = $\lambda Q \lambda P \exists x[\forall y[\overset{V}{Q}(y) \leftrightarrow x=y] \wedge \overset{V}{P}(x)]$.

3.6. END

Formulas (25), (26) and (27) are not in all respects a satisfactory representation of the meanings of sentences (22), (23) and (24) respectively. The formulas contain quantifications over individual concepts, whereas one would prefer a quantification over individuals. The conditions for application of RR_3 are not satisfied, so we have no ground for the elimination of the individual concepts by means of an application of this rule. On the contrary: as I will explain, the replacement of (31) by (32) would replace a formula by a non-equivalent one.

(31) $\exists x[woman(x) \wedge run(x)]$

(32) $\exists u[woman_*(u) \wedge run_*(u)]$.

A possible choice for the value of x in (31) would be to assign to x the same interpretation as to *bigboss*, but in (32) there is not a corresponding choice. One would prefer to have (32) as the meaning representation of the meaning of (25) because intuitively (25) gives information about individuals, and not about individual concepts. Following Montague, we obtain this effect by means of the introduction of a meaning postulate. Only those models for intensional logic are possible models for the interpretation of English in which the following meaning postulate holds.

3.7. Meaning postulate 2

$$\Box [\delta(x) \rightarrow \exists u[x = u]]$$

where $\delta \in \{man, woman, park, fish, pen, unicorn\}$.

3.7. END

This meaning postulate says that constants such as *man* can yield true only for constant individual concepts, i.e. for individual concepts which yield for every index the same individual. Note that the constants *price* and *temperature* are not mentioned in MP_2 . Arguments for this, and examples involving *price* and *temperature* will be given in appendix 1 of this volume. As a consequence of MP_2 , it can be shown that (31) and (32) are equivalent. I will not present a proof for this, because it is only one of the situations in which MP_2 will be used. In appendix 1, it will be investigated in general in which circumstances MP_2 allows us to replace a quantification over individual concepts by a quantification over individuals. For the moment it suffices to know that in all examples we will meet, such a replacement is allowed. This is expressed in the following reduction rule.

3.8. Reduction Rule 4

Let be given a formula of one of the following forms: $\exists x[\delta(x) \wedge \phi(x)]$, $\forall x[\delta(x) \rightarrow \phi(x)]$ or $\exists x[\forall y[\delta(y) \leftrightarrow x=y] \wedge \phi(x)]$.

If MP2 holds for δ , then replace these formulas by respectively $\exists u[\delta(\wedge u) \wedge \phi(\wedge u)]$, $\forall u[\delta(\wedge u) \rightarrow \phi(\wedge u)]$ or $\exists u[\forall v[\delta(\wedge v) \leftrightarrow u=v] \wedge \phi(\wedge u)]$.

CORRECTNESS PROOF. See appendix 2.

3.8. END

The production of the sentence mentioned in the title of this section is given in figure 4.




Figure 4

Note how in this simple example RR_4 and RR_3 are used in order to simplify the translation of *the woman*, and RR_1 and RR_3 to simplify the translation of *the woman walks*. In the sequel such reductions will often be performed without any further comment.

4. MARY WALKS AND SHE TALKS

In this section the fragment is extended with rules for disjunction and conjunction, and with a rule for co-referentiality. The rules for producing conjoined sentences are as follows.

4.1. Rule S_{11a} :

$S \times S \rightarrow S$

F_{11a} : concatenate (α , and, β)

T_{11a} : $\alpha' \wedge \beta'$

4.2. Rule S_{11b}:
$$S \times S \rightarrow S$$

$$F_{11b}: \text{concatenate } (\alpha, \text{or}, \beta)$$

$$T_{11b}: \alpha' \vee \beta'$$

4.2. END

Notice that the words *and* and *or* are not members of a category of connectives: they are introduced syncategorematically. It would be possible to have a three-place rule for sentence conjunction, with for the connective *and* as translation $\lambda\phi\lambda\psi[\phi \wedge \psi]$. This categorical approach is not followed here because there are rules for disjunction and conjunction for other categories as well. Furthermore, the situation is complicated by the fact that there is term disjunction in the fragment, but no term conjunction (in order to avoid plurals). In this situation it would not be a simplification to use a categorical treatment of connectives. For a categorical treatment in a somewhat different framework, see GAZDAR 1980.

The rules for forming conjoined phrases of other categories than sentences are as follows.

4.3. Rule S_{12a}:
$$IV \times IV \rightarrow IV$$

$$F_{12a}: \text{concatenate } (\alpha, \text{and } \beta)$$

$$T_{12a}: \lambda x[\alpha'(x) \wedge \beta'(x)].$$
4.4. Rule S_{12b}:
$$IV \times IV \rightarrow IV$$

$$F_{12b}: \text{concatenate } (\alpha, \text{or } \beta)$$

$$T_{12b}: \lambda x[\alpha'(x) \vee \beta'(x)].$$
4.5. Rule S₁₃:
$$T \times T \rightarrow T$$

$$F_{13}: \text{concatenate } (\alpha, \text{or } \beta)$$

$$T_{13}: \lambda P[\alpha'(P) \vee \beta'(P)].$$

4.5. END

The production of (33) is given in figure 5.

(33) *John walks and talks.*




Figure 5

Note that the produced sentence is not identical with (33). The treatment presented in figure 5 obeys the formulation of S_4 , and, therefore, only the first verb is conjugated. For an improved treatment see chapter 8, or FRIEDMAN 1979b.

An example of term disjunction is given in (34).

(34) *John or Mary talks.*

First (35) is formed according to S_{13} . Its unreduced translation is (36).

(35) *John or Mary*

(36) $\lambda P[\lambda P[\overset{V}{P}(\wedge \text{john})](P) \vee \lambda P[\overset{V}{P}(\wedge \text{mary})](P)]$.

Formula (36) contains several occurrences of the variable P , and three binders for P (viz. three occurrences of λP). However, due to the different scopes of the lambda operators, it is uniquely determined which variables occur in the scope of each of the lambda operators. The conditions for λ -conversion are satisfied, and after two applications of RR_1 formula (36) reduces to (37).

(37) $\lambda P[\overset{V}{P}(\wedge \text{john}) \vee \overset{V}{P}(\wedge \text{mary})]$.

Application of S_4 to term (35) and the verb *talk*, yields (34), which has as unreduced translation (38). This formula reduces by application of RR_1 and RR_2 to (39), and using RR_3 to (40).

(38) $\lambda P[\overset{V}{P}(\wedge \text{john}) \vee \overset{V}{P}(\wedge \text{mary})](\wedge \text{talk})$

(39) $\text{talk}(\wedge \text{john}) \vee \text{talk}(\wedge \text{mary})$

(40) $\text{talk}_*(\text{john}) \vee \text{talk}_*(\text{mary})$.

In sentences containing conjunctions or disjunctions pronouns occur often which are coreferential with some other term in that sentence. An example is the coreferentiality of *she* and *Mary* in (41).

(41) *Mary walks and she talks.*

In order to account for coreferentiality, a collection of new -artificial- terms is introduced. Since they have a relationship with logical variables, they are called syntactic variables. These variables are not words of English, and might be represented by means of some artificial symbol. Since the variables are related to pronouns, it has some advantages, to give them a representation exhibiting this relationship. The variables are written as male pronouns provided with an index (e.g. he_n). Their translations contain logical variables x_n of type $\langle s, e \rangle$. The syntactic variables he_n are generators of sort T.

4.6. $\{he_1, he_2, \dots\} \subset B_T$.

4.7. $he'_1 = \lambda P[\overset{V}{P}(x_1)]$, $he'_2 = \lambda P[\overset{V}{P}(x_2)]$,

4.7. END

One of the most important rules of PTQ is S_{14} . As for the syntax it removes the syntactic variables. As for the translation, it binds the corresponding logical variables. This rule enables us to deal with most of the ambiguities mentioned in the introduction, but in this section we will only deal with its use for coreferentiality. In fact S_{14} is not a rule, but rather a rule-scheme which for each choice of the index n constitutes a rule. This aspect will be indicated by using the parameter n in the description of the rule scheme.

4.8. Rule $S_{14,n}$:

$T \times S \rightarrow S$

$F_{14,n}$: If $\alpha = he_k$ then replace all occurrences of he_n/him_n in β by he_k/him_k respectively.

Otherwise replace the first occurrence of he_n in β by α , and replace all other occurrences of he_n in β by *he/she/it* and of him_n by *him/her/it* according to the gender of the first CN or T in α .

$$T_{14,n}: \alpha'(\wedge \lambda x_n [\beta']).$$

4.8. END

An example of the use of (an instance of) $S_{14,n}$ arises in the production of (41), as presented in figure 6.




Figure 6

The translation for (41) given in figure 6 can be reduced, using RR3, to (42).

$$(42) [\wedge \lambda x_1 [\text{walk}(x_1) \wedge \text{talk}(x_1)]](\wedge \text{mary}).$$

By application of RR2 and RR1 this reduces to (43), and by RR3, further to (44).

$$(43) \text{walk}(\wedge \text{mary}) \wedge \text{talk}(\wedge \text{mary})$$

$$(44) \text{walk}_*(\text{mary}) \wedge \text{talk}_*(\text{mary}).$$

Some syntactic details of $S_{14,n}$ give rise to problems. The rule for term disjunction allows us to produce term phrases like he_1 and *Mary*, and he_1 or he_2 . In both cases it is not clear what is to be understood by the gender of the first T or CN in such a term. And if the term *John* or *Mary* is formed, it is not correct to use the pronoun *he*, but one should use *he* or *she*, witness the following example (FRIEDMAN, 1979).

$$(45) \text{John or Mary walks and he or she talks.}$$

It would require a more sophisticated syntax than we have available here in order to account correctly for these problems (see FRIEDMAN 1979 for an improved treatment).

The detail of $S_{14,n}$ that the first occurrence of he_n/him_n has to be

replaced, is explained as follows. A pronoun may always be coreferential with a common noun or term occurring earlier in the sentence, but it may not always refer forward to terms or nouns occurring later. So it is a safe strategy to put the coreferential noun phrase always in a position which is as leftmost as possible. It is a difficult, and not completely solved task, to characterize the situations in which a pronoun may refer to a term occurring later in the sentence. Therefore S_{14} describes only reference to terms occurring earlier than the pronoun. Even this safe procedure does not avoid all problems. In some cases a personal pronoun is produced, where a reflexive pronoun is required. Sentence (46) has, according to the rules described here, a translation which expresses that John loves himself. This result is, of course, incorrect.

(46) *John loves him.*

Our aim was to deal with certain semantic problems, and therefore I will not consider here proposals for dealing with this syntactic problem (one of the proposals from the literature, viz. PARTEE 1973, will be considered in chapters 5 and 6 although not from the present point of view).

5. JOHN FINDS A UNICORN

In this section the category TV of transitive verb phrases is introduced. The generators of this category are as follows.

5.1. $B_{TV} = \{find, loose, eat, love, date, be, seek, conceive\}.$

5.1. END

Corresponding with these TV's (except for *be*), there are constants in the logic. They denote higher order functions which take as argument the intension of a term translation, and yield an element of the same type as the translations of IV-phrases. The translations of the basic verbs of the category TV are the corresponding constants; the translation of *be* is a compound expression of the same type. Let us indicate by $\tau(C)$ the type of the translation of an expression of category C. Then $\tau(TV) = \langle\langle s, \tau(T) \rangle, \tau(IV) \rangle$. This explains the following definitions

5.2. $\{find, loose, eat, love, date, seek, conceive\} \subset CON_{\langle\langle s, \tau(T) \rangle, \tau(IV) \rangle}$

- 5.3. $find' = find, loose' = loose, eat' = eat, love' = love,$
 $be' = \lambda P \lambda x [\lambda y [\lambda x [\lambda y [x = y]]]]$ where $P \in VAR_{\langle S, \tau(T) \rangle}$
 $seek' = seek, conceive' = conceive.$

5.3. END

Out of a TV and a Term and IV can be formed according to the following rule.

5.4. Rule S_5 :

$TV \times T \rightarrow IV$

F_5 : concatenate (α, β)

T_5 : $\alpha' (\wedge \beta')$.

5.4. END

An example of the use of this rule is the production of (47), partially presented in figure 7.

(47) *John seeks a unicorn.*




Figure 7

The translation obtained in figure 7 is not the traditional one: one would like to consider *seek* as a two-place relation. Therefore the following convention is introduced.

5.5. DEFINITION. $\gamma(\alpha, \beta) = \gamma(\beta)(\alpha)$, where γ is an expression translating a TV.

5.5. END

In PTQ (p.259) this convention is defined for all γ . It is however only useful for TV's (see section 11). The above definition gives rise to the following reduction rule.

5.6. Reduction rule 5

Let be given a formula of the form $\gamma(\beta)(\alpha)$, where γ is the translation of some transitive verb. Then replace this formula by $\gamma(\alpha, \beta)$.

CORRECTNESS PROOF

See definition 5.5.

5.6. END

Using RR5, the formula obtained in figure 7 reduces to (48).

$$(48) \text{ seek}(\overset{\wedge}{\text{john}}, \overset{\wedge}{\lambda P \exists u}[\text{unicorn}_*(u) \wedge \overset{\vee}{P}(\hat{u})]).$$

This translation describes the de-dicto reading of (47). The de-re reading will be considered in section 6. Below I will discuss whether the formula expresses a relation between the right kinds of semantic objects.

The first argument of *seek* is a constant individual concept. One might wish to have an individual as first argument. In analogy of the δ_* notation for intransitive verbs, we might introduce a notation for transitive verbs in which the $\overset{\wedge}$ in front of *john* disappears. PARTEE (1975, p.290) has proposed such a notation, but it is not employed in the literature, therefore I will not use it here. Notice that the interpretation of (48) is tantamount to a relation of which the first component is an individual (see section 2).

The second argument in (48) is the intension of a collection of properties. So *seek* is not treated as a relation between two individuals, and therefore (48) does not allow for the conclusion that there is a particular unicorn which John seeks. In this way the problem mentioned in section 1 is solved, so in this respect the formula is satisfactory. But one might ask whether this effect could be obtained by means of a simpler formula, viz. one without the intension sign. The need for this intension in the second argument is explained as follows (JANSSEN 1978b, p.134). Suppose that *seek* is considered as a relation between an individual and (the characteristic function of) a set of properties. Consider a world in which there exist no unicorns. Then for no property *P* it is true that $\exists u[\text{unicorn}_*(u) \wedge \overset{\vee}{P}(\hat{u})]$. Thus in these circumstances $\overset{\wedge}{\lambda P \exists u}[\text{unicorn}_*(u) \wedge \overset{\vee}{P}(\hat{u})]$ is the

characteristic function of the empty set of properties. The semantic interpretation of *John seeks a unicorn* then states that the seek-relation holds between John and this empty set. Suppose moreover that in this world also no centaurs exist. Then the semantic interpretation of

(49) *John seeks a centaur*

also expresses that the seek-relation holds between John and the empty set of properties. But this contradicts our intuition that (47) and (49) have different meanings. When we wish to describe the difference between centaurs and unicorns we cannot restrict our attention to the present state of the present world. We should also consider other worlds (or other states of the present world) for instance, those in which unicorns or centaurs do exist. In other worlds the set $\lambda P \exists u [unicorn_*(u) \wedge \forall P(\wedge u)]$ might be different from $\lambda P \exists u [centaur_*(u) \wedge \forall P(\wedge u)]$. Therefore the *seek*-relation will be considered as a relation between individuals and intensions of sets of properties. Since these intensions are different, *seek a unicorn* will get an interpretation different from the one for *seek a centaur* (even if both are extinct).

In the same way as we produced *John seeks a unicorn*, we may produce (50) with as reduced translation (51).

(50) *John seeks Mary*

(51) $seek(\wedge john, \wedge \lambda P[\forall P](\wedge mary))$.

This formula expresses that the seek relation holds between an individual concept and the collection of properties of Mary. But sentence (50) expresses that the seek-relation holds between two individuals: between John and Mary. One would like to have this aspect expressed by the obtained formula. Therefore the following definition (PTQ, p.265).

5.7. DEFINITION. $\delta_* = \lambda v \lambda u \delta(\wedge u, \wedge \lambda P[\forall P(\wedge v)])$, where $\delta \in CON_{\tau}(TV)$.

5.7. END

On the basis of this definition we have the following reduction rule.

5.8. Reduction rule 6

Let be given an expression of the form $\delta(\wedge \alpha, \wedge \lambda P[\forall P(\wedge \beta)])$, where $\alpha, \beta \in VAR_e \cup CON_e$, and $\delta \in CON_{\tau}(TV)$. Then replace this expression by $\delta_*(\alpha, \beta)$.

CORRECTNESS PROOF

$$\begin{aligned} \delta_*(\alpha, \beta) &= \delta_*(\beta)(\alpha) = \lambda v \lambda u \delta(\wedge u, \wedge \lambda P[\vee P(\wedge v)])(\beta)(\alpha) = \{\text{RR1}\} = \\ &= \delta(\wedge \alpha, \wedge \lambda P[\vee P(\wedge \beta)]). \end{aligned}$$

Note that λ -reduction is allowed because the constants of type e in the fragment are rigid.

5.8. END

Using RR6 we may reduce (51) to (52).

$$(52) \text{ seek}_*(\text{john}, \text{mary}).$$

In the same way as we produced the sentence *John seeks a unicorn*, we may produce (53), with translation (54).

$$(53) \text{ John finds a unicorn}$$

$$(54) \text{ find}(\wedge \text{john}, \wedge \lambda P[\exists u \text{ unicorn}_*(u) \wedge \vee P(\wedge u)]).$$

This result is not precisely what we would like to have. Sentence (53) gives the information that there exists at least one unicorn, and (54) does not express this information. In order to deal with this aspect we restrict our attention to those models for IL in which the following meaning postulate is satisfied.

5.8. Meaning Postulate 4

$$\exists s \forall x \forall P \square [\delta(x, P) \leftrightarrow \vee P(\wedge \lambda y \vee S(\vee x, \vee y))]$$

where $\delta \in \{\text{find}, \text{loose}, \text{eat}, \text{love}, \text{date}\}$ and $P \in \text{VAR}_{\langle s, \tau(T) \rangle}$.

5.8. END

This meaning postulate expresses that if the relation δ holds between an individual concept and a collection of properties, then there is a corresponding relation which holds between individuals. This relation is index dependent: the set of pairs which consist of a 'finder' and a 'found object', may be different for different indices. Therefore the existence of a relation between finders and found objects is formalized by means of an existential quantification over a variable which is of one intension level higher than the relation itself. An equivalent alternative would be (55), where the quantification $\exists S$ is within the scope of \square (this variant is due to P. van Emde Boas).

$$(55) \quad \Box [\exists S \forall x \forall P [\delta(x, P) \leftrightarrow P(\wedge \lambda y S(\overset{V}{x}, \overset{V}{y}))]].$$

A notation for the relation between finder and found object is already provided by the δ_* notation. This notation is introduced in the following rule.

5.9. Reduction rule 7

Let be given an expression of the form $\delta(\alpha, \beta)$ where $\delta \in \{find, loose, eat, love, date\}$ and $\alpha \in ME_{\langle s, e \rangle}$ $\beta \in ME_{\tau(T)}$. Then, replace this expression by $\overset{V}{\beta}(\wedge \lambda y [\delta_*(\overset{V}{\alpha}, \overset{V}{y})])$.

CORRECTNESS PROOF

From MP_4 follows that for all g , there is a $g' \approx_S g$ such that

$$g' \models \Box [\delta(x, P) \leftrightarrow \overset{V}{P}(\wedge \lambda y \overset{V}{S}(\overset{V}{x}, \overset{V}{y}))].$$

This means that for all expressions $\alpha \in ME_{\langle s, e \rangle}$, $\beta \in ME_{\langle s, \tau(T) \rangle}$ holds that

$$g' \models \delta(\alpha, \beta) \leftrightarrow \overset{V}{\beta}(\wedge \lambda y \overset{V}{S}(\overset{V}{\alpha}, \overset{V}{y})).$$

For this g' the following equalities hold:

$$\begin{aligned} \overset{V}{\beta}(\wedge \lambda y \delta_*(\overset{V}{\alpha}, \overset{V}{y})) &= \{\text{Def.5.5}\} = \beta(\wedge \lambda y \delta_*(\overset{V}{y})(\overset{V}{\alpha})) = \{\text{Def.5.7}\} = \\ \overset{V}{\beta}(\wedge \lambda y \lambda v \lambda u \delta(\wedge u, \wedge \lambda P [\overset{V}{P}(\overset{V}{v})])(\overset{V}{y})(\overset{V}{\alpha})) &= \{\text{choice of } g'\} = \\ \overset{V}{\beta}(\wedge \lambda y \lambda v \lambda u [\wedge \lambda P [\overset{V}{P}(\overset{V}{v})]](\wedge \lambda y \overset{V}{S}(\overset{V}{u}, \overset{V}{y}))(\overset{V}{y})(\overset{V}{\alpha})) &= \{\text{RR}_{2,1}\} = \\ \overset{V}{\beta}(\wedge \lambda y \lambda v \lambda u [\overset{V}{\lambda y \overset{V}{S}}(u, \overset{V}{y})(\overset{V}{v})](\overset{V}{y})(\overset{V}{\alpha})) &= \{\text{RR}_{2,1}\} = \\ \overset{V}{\beta}(\wedge \lambda y \lambda v \lambda u [\overset{V}{S}(u, \overset{V}{v})](\overset{V}{y})(\overset{V}{\alpha})) &= \{\text{RR}_{2,1}\} = \\ \overset{V}{\beta}(\wedge \lambda y \overset{V}{S}(\overset{V}{\alpha}, \overset{V}{y})) &= \{\text{choice of } g'\} = \delta(\alpha, \beta). \end{aligned}$$

Since S does not occur in the first and last formula, these expressions are equivalent for all g . From these equalities the reduction rule follows.

5.9. END

After the introduction of RR_7 we return to our discussion of (54). Application of RR_7 to (54) yields (56).

$$(56) \quad \overset{V}{\lambda P} \exists u [\text{unicorn}_*(u) \wedge \overset{V}{P}(\overset{V}{u})] (\wedge \lambda y [\text{find}_*(\overset{V}{john}, \overset{V}{y})])$$

This reduces further to (57), and that is the kind of formula we were looking for: it expresses that the find-relation holds between two individuals

$$(57) \exists u[\text{unicorn}_*(u) \wedge \text{find}_*(\text{john}, u)].$$

The fragment contains one single verb *be*, which is used both for the be of identity, and for the copula be. An example of the be of identity is given in (58).

$$(58) \text{John is Mary.}$$

The first step in its production is to combine *be* with *Mary* according to S_5 . This yields the IV-phrase *be Mary*. The translation of this phrase reduces by several applications of RR_1 and RR_2 to $\lambda x[\overset{V}{x}=\text{mary}]$. Combining this with *John* according to S_4 yields (58), and the corresponding translation reduces by applications of RR_1 and RR_2 to $\text{john} = \text{mary}$. One observes that the final result is an identity on the level of individuals. This shows why there is no meaning-postulate like MP_4 introduced for *be*: its translation already applies to the level of individuals rather than the level of individual concepts.

Next I give an example of the copula use of *be*.

$$(59) \text{John is a man.}$$

First the IV-phrase *be a man* is formed. Its translation reduces to the formula $\lambda x \exists u[\text{man}_*(u) \wedge \overset{V}{x}=u]$. Combining this with the translation of *John* yields as translation (60), which reduces to (61).

$$(60) \lambda P[\overset{V}{P}(\wedge \text{john})] \lambda x[\exists u \text{man}_*(u) \wedge \overset{V}{x}=u]$$

$$(61) \exists u[\text{man}_*(u) \wedge \text{john}=u].$$

In this situation one could perform one further simplification replacing (61) by (62); below I will explain why I will not do so.

$$(62) \text{man}_*(\text{john}).$$

It would of course be possible to introduce a new reduction rule performing this last reduction. But it is difficult to cover the reduction from (61) to (62) by a general rule. Suppose that a rule R would say when the occurrence of a subformula $\text{john}=u$ implies that all occurrences of u may be replaced by *john*. In order to decide whether reduction is possible, R has to take the whole formula into consideration. Reduction from (61) to (62) is allowed, but if in (61) connective \wedge would be replaced by \rightarrow the

reduction is not allowed. This supposed rule R would have a different character than all other reduction rules. The other rules are 'local': the question whether they may be applied, can be answered by inspecting a context of fixed length. But R would not be local because the whole formula has to be taken into account. I will not try to design such a rule R because I prefer to have only local reduction rules. Moreover, the set of reduction rules is incomplete, even with such a rule R, and only a partial solution of the reduction problem is possible. This one sees as follows. Suppose that we would define in each class of logically equivalent formulas one formula as being the simplest one (say some particular formula with shortest length). Then there exists no algorithm which reduces all formulas to the simplest in their class, since otherwise we could decide the equivalence of two formulas by reducing them to their simplest form and looking whether they are identical. Such a decision procedure would contradict the undecidability of IL (see also chapter 6, section 4).

6. EVERY MAN LOVES A WOMAN

The rules introduced in the previous sections allow us to produce sentence (63).

(63) *Every man loves a woman.*

In the introduction (section 1) I have described the two readings of this sentence. On the one reading, the same woman is loved by every man (say Brigitte Bardot), and on the other reading it might for every man be another woman (say his own mother). These two readings are represented by (64) and (65) respectively.

(64) $\exists v[woman_*(v) \wedge \forall u[man_*(u) \rightarrow love_*(u,v)]]$

(65) $\forall u[man_*(u) \rightarrow \exists v[woman_*(v) \wedge love_*(u,v)]]$.

Note that the difference between (64) and (65) is a difference in the scope of the quantifiers \forall and \exists . Therefore this ambiguity is called a scope ambiguity. A well known variant of this scope ambiguity is (66).

(66) *Every man in this room speaks two languages.*

A remarkable aspect of the two readings of (63) is that the one reading has the other as a special case: from (64) it follows that (65) holds. Therefore one might doubt whether the two formulas really constitute

an ambiguity we should deal with. One might say that the weaker formula (viz. (65)) describes the meaning of (63), and that, with additional information from the context, this can be narrowed down to the stronger one. This argument holds for (63), but I will illustrate, that it is not generally applicable. Consider (67), due to LANDMAN & MOERDIJK (1981,1983).

(67) *Every schoolboy believes that a mathematician wrote 'Through the looking glass'.*

This sentence is (at least) twofold ambiguous. On the one reading there is one mathematician of which every schoolboy believes that he wrote 'through the looking glass', but not every schoolboy necessarily believes that the person was a mathematician. On the other reading every schoolboy has the belief that some mathematician wrote the book, without necessarily having a special mathematician in mind. The rules needed for the production of sentences like (67) will be given in section 9. The formulas we will obtain then, are presented below in a somewhat simplified form. Formula (68) corresponds with the first reading (the believes concern the same mathematician), the second reading is represented by (69).

(68) $\exists v[\text{mathematician}_*(v) \wedge \forall u[\text{schoolboy}_*(u) \rightarrow \text{believe}_*(u, \text{wrote}_*(v, \text{'Through the looking glass'}))]]]$

(69) $\forall u[\text{schoolboy}_*(u) \rightarrow \text{believe}_*(u, \exists v[\text{mathematician}_*(v) \wedge \text{wrote}_*(v, \text{'Through the looking glass'})])]]$.

These two readings are logically independent: the one can be true while the other is false. The same situation arises for the well known example (66): if we read in that sentence *two* as *precisely two*, then the different scope readings are logically independent. These examples show that for variants of the scope ambiguity, both readings have to be produced by the grammar. Then it is not clear why (63) should get only one reading.

A part of the production of reading (65) of sentence (63) is given in figure 8. This production is called the direct production (because no quantification rule is used).

$$\begin{array}{l}
 \text{Every man loves a woman } \{S, 4\} \\
 \forall u[\text{man}_*(u) \rightarrow \text{love}(\lambda P \exists u[\text{woman}_*(u) \wedge \forall P(\wedge u)])(\wedge u)] \\
 \text{Every man } \{T, 2\} \qquad \qquad \qquad \text{Love a woman } \{IV, 5\} \\
 \lambda P \forall u[\text{man}_*(u) \rightarrow \forall P(\wedge u)] \qquad \text{love}(\lambda P \exists u[\text{woman}_*(u) \wedge \forall P(\wedge u)])
 \end{array}$$

Figure 8

The translation obtained in figure 8 can be reduced further by an application of RR_5 , yielding (70).

$$(70) \forall u[man_*(u) \rightarrow love(\wedge u, \wedge \lambda P \exists u[woman_*(u) \wedge P(\wedge u)])].$$

Application of RR_6 yields (71), and twice application of RR_2 yields (72).

$$(71) \forall u[man_*(u) \rightarrow [\wedge \lambda P \exists u[woman_*(u) \wedge \vee P(\wedge u)]](\wedge \lambda y[love_*(\vee \wedge u, \vee y)])]$$

$$(72) \forall u[man_*(u) \rightarrow [\lambda P \exists u[woman_*(u) \wedge \vee P(\wedge u)]](\wedge \lambda y[love_*(u, \vee y)])].$$

Further application of lambda conversion is not allowed because this would bring the u in $love(u, \vee y)$ under the scope of $\exists u$. In order to simplify this formula further, we first have to replace the variable u bound by $\exists u$ by another variable.

6.1. Reduction rule 8

Let be given an expression of the form $\lambda z\phi, \exists z\phi$ or $\forall z\phi$. Let w be a variable of the same type as z , but which does not occur in ϕ . Then replace $\lambda z\phi, \exists z\phi, \forall z\phi$ by respectively $\lambda w[w/z]\phi, \exists w[w/z]\phi$, and $\forall w[w/z]\phi$.

CORRECTNESS PROOF

Evident from the interpretation of these formulas.

6.1. END

Application of RR_8 to (72) yields (73). Applications of RR_1 and RR_2 yield then (74), which reduces further to (75).

$$(73) \forall u[man_*(u) \rightarrow [\lambda P \exists v[woman_*(v) \wedge \vee P(\wedge v)]](\wedge \lambda y[love_*(u, \vee y)])]$$

$$(74) \forall u[man_*(u) \rightarrow \exists v[woman_*(v) \wedge [\lambda y[love_*(u, \vee y)]](\wedge v)]]$$

$$(75) \forall u[man_*(u) \rightarrow \exists v[woman_*(v) \wedge love_*(u, v)]]].$$

A part of the production of reading (64) of sentence (63) is given in figure 9. The production uses $S_{14,n}$, and it is called (for this reason) an indirect production of (63).




Figure 9

The translation obtained in figure 9 reduces by application of RR_1 and RR_2 to (76).

$$(76) \exists u[\text{woman}_*(u) \wedge \lambda x_1[\forall u[\text{man}_*(u) \rightarrow \text{love}_*(\forall x_1, u)]](\wedge u)].$$

After change of bound variable (RR_7) we apply RR_1 , and obtain (77).

$$(77) \exists u[\text{woman}_*(u) \wedge \forall v[\text{man}_*(v) \rightarrow \text{love}_*(v, u)]]].$$

In the introduction I have already said that sentence (78) is ambiguous; its ambiguity is called the de-dicto/de-re ambiguity. From the de-re reading (79) it follows that unicorns exist, whereas this does not follow from the de-dicto reading (80).

(78) *John seeks a unicorn*

$$(79) \exists u[\text{unicorn}_*(u) \wedge \text{seek}_*(\text{john}, u)]$$

$$(80) \text{seek}(\wedge \text{john}, \wedge \lambda P\exists u[\text{unicorn}_*(u) \wedge P(\wedge u)]).$$

This ambiguity can be considered as a scope ambiguity: the difference between (79) and (80) is the difference in scope of the existential quantifier. Note that formulas (79) and (80) are logically independent, hence we have to produce both readings. These productions are analogous to the productions of the different scope readings of *Every man loves a woman*. The de-dicto reading (80) is obtained by a direct production. We have considered this production in the previous section. The de-re reading, viz. (79), is obtained by an indirect production. As a first stage of the indirect production sentence (81) is formed, which has (82) as translation.

(81) *John seeks him*₁

$$(82) \text{seek}(\wedge \text{john}, \wedge \lambda P[\forall P(x_1)]).$$

Combination according to $S_{14,1}$, of (81) with the term *a unicorn* yields (78), and combination of their translations according to $T_{14,1}$ yields (83), reducing to (84).

$$(83) \lambda P \exists u [unicorn_*(u) \wedge \forall P(\wedge u)] (\wedge \lambda x_1 [\wedge seek(\wedge john, \wedge \lambda P [\forall P(x_1)])])]$$

$$(84) \exists u [unicorn_*(u) \wedge seek(\wedge john, \wedge \lambda P [\forall P(\wedge u)])].$$

Application of RR_6 reduces this formula to (85).

$$(85) \exists u [unicorn_*(u) \wedge seek_*(john, u)].$$

Sentence (86) can be produced using the same syntactic rules as in the production of (78).

$$(86) \textit{Mary finds a unicorn.}$$

This sentence is not ambiguous; it only has a referential reading. In the previous section it was explained how the translation of the direct production reduces to such a reading. The indirect production yields, of course, a referential reading as well. An interesting aspect of the indirect production is the way in which the obtained formulas can be reduced. For this reason I will consider this production in more detail. A first stage of the indirect production of (86) is (87), which has (88) as translation.

$$(87) \textit{Mary finds him}_1.$$

$$(88) find(\wedge mary, \wedge \lambda P [\forall P(x_1)]).$$

One method to reduce (88) is to apply the same reduction rules as used in the reduction of (82). Then as last step RR_6 is applied, see the reduction of (84). But another reduction process is possible as well. We might apply RR_7 to (88) because meaning postulate 4 holds for *find*. Thus we obtain (89), reducing to (90).

$$(89) \wedge \lambda P [\forall P(x_1)] (\wedge \lambda y [find_*(mary, \forall y)])]$$

$$(90) find_*(mary, \forall x_1).$$

Combination, according to $S_{14,1}$ of (90) with the translation of *a unicorn* yields (91), which reduces to (92).

$$(91) \lambda P \exists u [unicorn_*(u) \wedge \forall P(\wedge u)] (\wedge \lambda x_1 [find_*(mary, \forall x_1)])]$$

$$(92) \exists u [unicorn_*(u) \wedge find_*(mary, u)].$$

This shows that there are two methods to reduce the formulas obtained in the indirect production of (86).

In general it makes no difference in which order we apply the reduction rules. Sooner or later we have to apply the same rule to the same (sub)expression. An exception is the introduction of δ_* for constants to which meaning postulate 4 applies. Once we have applied the meaning postulate (i.e. RR_7), we cannot apply the definition for δ_* (i.e. RR_6) any more. The reason for this is that both applications consume an occurrence of δ , and produce an occurrence of δ_* . As practice learns, these two ways of reduction always yield the same result. I have, however, not a formal proof of some formal version of this observation. The situation is difficult due to the interaction of RR_6 and RR_7 with many other reduction rules. In FRIEDMAN & WARREN (1979) related reduction rules are considered, and they provide several examples of the complex interactions of the rules (they have no normal form theorem for their system either).

Finally I consider a sentence which is not ambiguous. For sentence (93) the de-re reading is the only possible reading, and it is the only reading produced by the grammar.

(93) *John seeks a unicorn and Mary seeks it.*

The occurrence of *it* requires an application of $S_{14,n}$. A part of the production of (93) is given in figure 10.




Figure 10

The obtained translation for (93) reduces to (94).

(94) $\exists u[\text{unicorn}_*(u) \wedge \text{seek}_*(\text{john}, u) \wedge \text{seek}_*(\text{mary}, u)]$.

7. BILL WALKS IN THE GARDEN

In this section the fragment is extended with the categories Prep of prepositions, and IAV of IV-modifying adverbials. In PTQ the category 'IAV' is also called 'IV/IV'. For the basic elements of IAV there are corresponding constants of type $\langle\langle s, \tau(IV) \rangle, \tau(IV) \rangle$. The definitions concerning IAV are as follows.

- 7.1. $B_{IAV} = \{slowly, voluntarily, allegedly\}$.
 7.2. $\{slowly, voluntarily, allegedly\} \subset CON_{\tau(IAV)}$
 7.3. $slowly' = slowly, voluntarily' = voluntarily,$
 $allegedly' = allegedly.$

7.3. END

An adverb forms with an IV-phrase, according to S_{10} , a new IV-phrase.

7.4. Rule S_{10} :

$IAV \times IV \rightarrow IV$

F_{10} : concatenate (α, β)

T_{10} : $\alpha'(\wedge\beta')$.

7.4. END

An example of a sentence containing an IAV is (95).

(95) *John voluntarily walks.*

The production of (95) is presented in figure 11.




Figure 11

In PTQ the convention was introduced to write all expressions of the form $\gamma(\alpha)(\beta)$ as $\gamma(\beta, \alpha)$. This example shows that the PTQ formulation was too

liberal: it would allow to write *voluntarily* as a relation:
 $voluntarily(\hat{\text{john}}, \hat{\text{walk}})$. This result is not attractive because tradition-
 ally one does not consider *voluntarily* as a relation. Therefore in reduc-
 tion rule 5 this convention was only introduced for γ being a verb.

The translation obtained for (95) does not allow for the conclusion
 that John walks, although this would be a correct conclusion from sentence
 (95). Not all adverbs allow for such a conclusion. From (96) it does not
 follow that John walks.

(96) *John allegedly walks.*

This means that the adverb *allegedly* creates an intensional context
 for the object of a verb. Also sentence (97) does not allow to con-
 clude to the existence of a unicorn.

(97) *John allegedly loves a unicorn.*

One might expect the introduction of a meaning postulate that expresses the
 extensional character of *slowly* and *voluntarily*. Such a meaning postulate
 is not given in PTQ. I expect that it would be of a different nature than
 the postulates we have met before: it would be an implication, and I ex-
 pect that it would not give rise to simplifications of the formulas in-
 volved.

The fragment contains two prepositions, and from these new adverbial-
 phrases can be formed. Prepositions translate into constants of type
 $\langle\langle s, \tau(T) \rangle, \tau(\text{IAV}) \rangle$.

7.5. $B_{\text{Prep}} = \{in, about\}$

7.6. $\{in, about\} \subset \text{CON}_{\tau(\text{Prep})}$

7.7. $in' = in, about' = about$.

7.7. END

The rule for creating new adverbs is as follows.

7.8. Rule S_6 :

$\text{Prep} \times T \rightarrow \text{IAV}$

F_6 : concatenate (α, β)

T_6 : $\alpha' (\hat{\beta}')$.

7.8. END

An example of an application of this rule is given in figure 12, where sentence (98) is produced.

(98) *John talks about a unicorn.*




Figure 12

The translation obtained here does not imply that there is a unicorn John talks about: *about* creates an intensional context. This is the result we aimed at (see section 1).

In the same way as we produced (98), we may produce (99) with as translation (100).

(99) *Bill walks in the park*

(100) $in(\wedge \lambda P \exists u [\forall v [\text{park}_*(v) \leftrightarrow u=v] \wedge \vee P(\wedge u)]) (\wedge \text{walk}) (\wedge \text{bill})$.

This result is not completely satisfactory. If Bill walks in the park, then one may conclude that there exists a park, and if the park is the Botanical garden, then from (99) it may be concluded that Bill walks in the Botanical garden. So the locative preposition *in* does not create an intensional context. This property of *in* is formalized in the following meaning postulate.

7.9. Meaning postulate 8

$$\exists G \forall P [\forall x \square [in(P)(Q)(x) \leftrightarrow \vee P(\wedge \lambda y [[\vee G](\vee y)(Q)(x)]]]$$

7.9. END

In order to be able to give a reduction rule on the basis of this meaning postulate, a notation for the predicate denoted by $\vee G$ in MP8 is

introduced (such a notation for prepositions is not defined in PTQ). This notation is chosen in analogy of the notation δ_* for verbs.

7.10. DEFINITION.

$$\delta_* = \lambda x \lambda Q \lambda u [\delta (\wedge \lambda P [\lambda P^V (\wedge u)]) (Q) (x)] \text{ where } \delta \in \text{CON}_{\tau}(\text{Prep}) .$$

7.10. END

On the basis of this definition we have the following reduction rule.

7.11. Reduction rule 9

Let be given an expression of the form $in(\alpha)(\beta)(\gamma)$, where $\alpha \in \text{ME}_{\langle S, \tau \rangle}(\text{T})$, $\beta \in \text{ME}_{\langle S, \tau \rangle}(\text{IV})$, $\gamma \in \text{ME}_{\langle S, e \rangle}$. Then replace this expression by $\alpha (\wedge \lambda y [in_*(y)(\beta)(\gamma)])$.

CORRECTNESS PROOF. Let $v \in \text{VAR}_e$, $x \in \text{VAR}_{\langle S, e \rangle}$ and $Q \in \text{VAR}_{\langle S, \tau \rangle}(\text{IV})$. Then for all g

$$g \models in_*(v)(Q)(x) = in(\wedge \lambda P^V P(\wedge v))(Q)(x) .$$

We now apply MP8 to the right hand side of the equality: this meaning postulate says that there is a $g' \sim_G g$ such that

$$g' \models in_*(v)(Q)(x) = [\wedge \lambda P^V P(\wedge v)](\wedge \lambda y [[^V G](^V y)(Q)(x)]) .$$

The expression to the right of the equality sign reduces by means of several applications of RR₁ and RR₂. Thus we obtain

$$g' \models in_*(v)(Q)(x) = [^V G](v)(Q)(x) .$$

Consequently $g' \models in_* = ^V G$. This means that from MP₈ it follows that

$$\models \forall P \forall Q \forall x \square [in(P)(Q)(x) \leftrightarrow P(\wedge \lambda y [in_*(^V y)(Q)(x)])] .$$

7.11. END

Formula (100) can be reduced, using RR₉, to (101) and further to (102)

$$(101) [\wedge \lambda P [\exists u \forall v [park_*(u) \leftrightarrow u=v] \wedge P(\wedge u)]](\wedge \lambda y [in_*(^V y)(\wedge walk)(\wedge bill)])$$

(102) $\exists u \forall v [\text{park}_*(v) \leftrightarrow u=v] \wedge \text{in}_*(u) (\wedge \text{walk}) (\wedge \text{bill})]$.

In PTQ no examples concerning the meaning postulate for *in* are given. This example illustrates the consequence of the meaning postulate: if one stands in the relation of walking in with 'a collection of properties', then there is an 'individual' with which one has this relation.

8. JOHN TRIES TO FIND A UNICORN

In this section a new category of IV-modifiers is introduced. This new category is called IV//IV (IV modifying verbs) and contains verbs taking verbs as complements. The fragment has only two of such verbs (*try to*, *wish to*), although there are a lot more in English. The syntactic treatment of these verbs is rather primitive: *try to* is considered as a single word containing a space (so *to* is not treated as a word). But our main interest is semantics, and the verbs are interesting in this respect. They create intensional contexts even when the sentence without such a verb would only have a de-re reading. An example is (103); this sentence does not necessarily have the implication that unicorns exist.

(103) *John tries to find a unicorn.*

Corresponding with the verbs of category IV//IV there are constants in the logic of the type $\langle \langle s, \tau(\text{IV}) \rangle, \tau(\text{IV}) \rangle$. The verbs translate into these constants.

- 8.1. $B_{\text{IV//IV}} = \{ \text{try to}, \text{wish to} \}$
 - 8.2. $\{ \text{try to}, \text{wish to} \} \subset \text{CON}_{\tau(\text{IV//IV})}$
 - 8.3. $\text{try to}' = \text{try to}, \text{wish to}' = \text{wish to}.$
- 8.3. END

The members of IV//IV are used in the following rule.

8.4. Rule S_8 :

$\text{IV//IV} \times \text{IV} \rightarrow \text{IV}$

F_8 : concatenate (α, β)

T_8 : $\alpha' (\wedge \beta')$.

8.4. END

The production of (103) is partially presented in figure 13.




Figure 13

The formula obtained in this production process does not reduce further, and it does not allow to conclude for the existence of a unicorn which John tries to find. So the de-dicto aspect is dealt with adequately. But sentence (103) can also be used in a situation in which there is a unicorn which John tries to find. For reasons related to the ones given concerning *John seeks a unicorn*, the reading involving a particular unicorn has to be obtained as an alternative translation for (103). That reading can be obtained using $S_{14,n}$.

The translation obtained for (103) in figure 13 is, however, not in all respects satisfactory. We do not get information concerning the relation between John and the property expressed in the second argument of *try to*. In particular it is not expressed that what John tries to achieve is that John (he himself) finds the unicorn, and not that someone else finds the unicorn. For verbs like *promise* and *permit* the relation between the subject and the complement is much more complex. A correction of this disadvantage of the PTQ treatment can be found along the lines of DOWTY (1978) and BARTSCH (1978b), see also section 4.1 in chapter VII.

In section 4 we introduced the rules for IV conjunction and disjunction. The verb phrases involved may concern two coreferential terms as in (104).

(104) *John finds a unicorn and eats it.*

The coreferentiality can be dealt with by means of quantifying in the term *a unicorn*. This yields the reading (105).

(105) $\exists u[\text{unicorn}_*(u) \wedge \text{find}_*(\text{john},u) \wedge \text{eat}_*(\text{john},u)]$.

This formula expresses that there is a particular unicorn which John finds and eats.

The conjoined verb phrase underlying (106) can be embedded in a *try to* construction.

(106) *John tries to find a unicorn and eat it.*

This sentence does not allow for the conclusion that there is a unicorn. The occurrence of a pronoun, however, invites us to produce this sentence with quantification rule S_{14} , and that would result in a referential reading, viz. (107)

(107) $\exists u[\text{unicorn}_*(u) \wedge \text{try to}(\wedge \text{john}, \wedge [\text{find}(\lambda P[\vee P(\wedge u)]) \wedge \text{eat}(\lambda P[\vee P(\wedge u)])])]$.

A new quantification rule makes it possible to produce (106) in a reading which does not imply the existence of a unicorn. The following rule scheme describes the quantification of a Term into an IV-phrase.

8.5. Rule $S_{16,n}$:

$T \times IV \rightarrow IV$

$F_{16,n}$: If α does not have the form he_k
 then replace in β the first occurrence of he_n or him_n
 by α , and all other occurrences of he_n by *he/she/it* and of
 him_n by *him/her/it* according to the gender of the first T
 or CN in α
 else replace all occurrences of he_n by he_k and of him_n by him_k .

$T_{16,n}$: $\lambda y[\alpha'(\wedge \lambda x_n[\beta'(y)])]$.

8.5. END

In order to produce (106) we first produce the verbphrase (108).

(108) *find a unicorn and eat it.*

The production of (108) is partially given in figure 14.




Figure 14

Now we return to the production of sentence (106). Its production from (108) is presented in figure 15.




Figure 15

A sentence related with (106) is (109).

(109) *John tries to find a unicorn and wishes to eat it.*

Montague argues that only a referential reading of this sentence is possible (except for the case that the pronoun *it* is considered as a pronoun of laziness). A production of sentence (109) might be given in which S_{14} is used. Then it is not surprising that a referential reading is obtained. But this is also the case for a production using S_{16} , as will be shown below. The first step is to form verb phrase (110), with translation (111).

(110) *try to find him₁ and wish to eat him₁*

(111) $\lambda x [\text{try to}(x, \overset{\wedge}{\text{find}}(\overset{\wedge}{\lambda P^V P}(x_1))) \wedge \text{wish to}(x, \overset{\wedge}{\text{eat}}(\overset{\wedge}{\lambda P^V P}(x_1)))].$

Combination of (110) with *a unicorn* according to $S_{16,1}$ yields (112). The translation is (113), which reduces to (114).

(112) *try to find a unicorn and wish to eat it*

$$(113) \lambda y[\lambda P\exists u[\text{unicorn}_*(u) \wedge \forall P(\wedge u)](\wedge \lambda x_1[\text{try-to}(y, \wedge \text{find}(\wedge \lambda P^V P(x_1))) \wedge \text{wish to}(y, \wedge \text{eat}(\wedge \lambda P^V P(x_1)))]]]$$

$$(114) \lambda y\exists u[\text{unicorn}_*(u) \wedge \text{try to}(y, \wedge \text{find}(\wedge \lambda P^V P(\wedge u))) \wedge \text{wish to}(y, \wedge \text{eat}(\wedge \lambda P^V P(\wedge u)))].$$

Combination of (112) with *John* according to S_4 yields sentence (109). The translation is (115).

$$(115) \exists u[\text{unicorn}_*(u) \wedge \text{try-to}(\wedge \text{john}, \wedge \text{find}(\wedge \lambda P^V P(\wedge u))) \wedge \text{wish-to}(\wedge \text{john}, \wedge \text{eat}(\wedge \lambda P^V P(\wedge u)))].$$

The formula obtained here can be simplified by replacing $\delta(\wedge \lambda P^V P(\wedge u))$ by $\lambda y \delta(\wedge \lambda P^V P(\wedge u))(y)$, where δ is the translation of a transitive verb. The advantage of this replacement is that now RR_5 and RR_6 can be used. In this way (115) reduces to (116).

$$(116) \exists u[\text{unicorn}_*(u) \wedge \text{try-to}(\wedge \text{john}, \wedge \lambda y \text{find}_*(\wedge y, u)) \wedge \text{wish to}(\wedge \text{john}, \wedge \lambda y \text{eat}_*(\wedge y, u))]$$

This method is formulated in a reduction rule as follows.

8.6. Reduction rule 10

Let be given an expression of the form $\delta(\wedge \lambda P^V P(\wedge u))$, where δ is the translation of a TV for which MP_4 holds. Then replace this expression by $\lambda y \delta_*(\wedge y, u)$.

CORRECTNESS PROOF. By definition of interpretation the two expressions are equivalent.

8.6. END

The possibilities for application of RR_{10} are limited by mentioning explicitly the argument of δ . One might omit this argument; then the rule would be applicable in many more circumstances, for instance to the formula obtained in figure 13. I have not used this more general version because it would not give rise to simpler formulas (in the sense of more concise formulas), but one might judge that the general rule would give rise to more understandable formulas.

9. JOHN BELIEVES THAT MARY WILL RUN

A new construction considered in this section arises from verbs of the category IV/S; I.e. verbs taking a sentence as complement. There are several such verbs, but only two of them are incorporated in the fragment.

- 9.1. $B_{IV/S} = \{believe\ that, assert\ that\}$
 9.2. $\{believe\ that, assert\ that\} \subset CON_{\langle\langle S, \tau(S) \rangle, \tau(IV) \rangle}$
 9.3. $believe\ that' = believe\ that, assert\ that' = assert\ that.$
 9.3. END

The rule producing IV phrases from these verbs reads as follows.

9.4. Rule S₇:

$IV/S \times S \rightarrow IV$

F_7 : concatenate (α, β)

T_7 : $\alpha' (^{\wedge}\beta')$.

9.4. END

An example of a sentence with a verb of category IV/S is (117).

(117) *John believes that Mary runs.*

Part of the production of (117) is given in figure 16.




Figure 16

Believe is considered as a relation between an individual concept and a proposition (i.e. a function from indices to truth values). It is not said what kind of relation this is. There are several proposals in the

literature analyzing the believe relation in more detail (e.g. LEWIS 1970), but Montague did not analyze it any further.

The formula obtained in figure 16 expresses that *believe* is a relation with as first argument $\hat{\ }^{\text{john}}$. To this, the same comment applies as to the first argument of the *seek*-relation: there is no generally accepted notation which expresses that for this argument *believe* can be considered as a relation with as first argument an individual. The second argument is an expression of type $\langle s, t \rangle$. Would it have been an expression of type t , then we could replace it by any other expression which denotes (for the current index) the same truth value. So if someone would believe a truth, he would believe all truths (for the current index). Now that the second argument of the believe-relation is a proposition, this is not the case. If John and Mary walk, then one may believe that John walks, without having the formal implication that one believes that Mary walks. Nevertheless, the use of a proposition is not completely satisfactory. It implies that in case John believes a tautology, he believes all tautologies. This is a fundamental shortcoming of this kind of approach; there is, however, not a generally accepted alternative.

The aspect that makes the introduction of *believe* and *assert* interesting in the present fragment, even with the present semantics, is that these verbs introduce intensional contexts in which a de-re reading is impossible. Sentence (118) does not allow for the conclusion that there exists a unicorn.

(118) *Mary believes that John finds a unicorn and he eats it.*

The relevant part of the production of sentence (120) is given in figure 17.

$$\begin{array}{l} \text{Mary believes that John finds a unicorn and he eats it} \\ \text{believe that}(\hat{\ }^{\text{mary}}, \hat{\ }^{\exists u}[\text{unicorn}_*(u) \wedge \text{find}_*(\text{john}, u) \wedge \text{eat}_*(\text{john}, u)]) \\ \text{Mary} \qquad \qquad \qquad \text{believe that John finds a unicorn and he eats it} \\ \lambda P[\overset{\vee}{P}(\hat{\ }^{\text{mary}})] \quad \text{believe that}(\hat{\ }^{\exists u}[\text{unicorn}_*(u) \wedge \text{find}_*(\text{john}, u) \wedge \text{eat}_*(\text{john}, u)]) \end{array}$$

Figure 17

A further extension of the fragment are the restrictive relative clauses: terms will be produced like *Every man such that he runs*. This *such that* form is not the standard form of relative clauses, but it avoids the syntactic complications arising from the use of relative pronouns. The

following rule scheme describes how relative clause constructions are formed out of a CN and a sentence.

9.5. Rule $S_{3,n}$:

$CN \times S \rightarrow CN$

$F_{3,n}$: replace in β all occurrences of he_n by *he/she/it* and him_n by *him/her/it* according to the gender of the first CN in α .

$T_{3,n}$: $\lambda x_n [\alpha'(x_n) \wedge \beta']$.

9.5. END

An example is the production of term (119), which is given in figure 18.

(119) *a man such that he runs.*




Figure 18

The obtained translation can be reduced, using RR_4 , to (120).

(120) $\lambda P \exists u [\text{man}_*(u) \wedge \text{run}_*(u) \wedge \overset{V}{P}(\hat{u})]$.

Rule $S_{3,n}$ takes a CN as one of its arguments, and yields a CN as result. This means that the rule can be applied more than one time in succession. Then terms are obtained like the one in (121)

(121) *Every man such that he walks such that he talks.*

In (121) both the relative clauses are attached to the head *man*; this phenomenon is called 'stacking'. A situation that may arise in connection with stacking is as follows. The second relative clause contains a pronoun which is coreferential with a term in the first relative clause, whereas the pronoun is (semantically) within the scope of the determiner of the whole term. An example, due to Bresnan (PARTEE 1975, p.263) is (124).

(122) *Every girl who attended a womans college who gave a donation to it, was put on the list.*

Sentence (124) exhibits co-reference within the compound CN phrase: *it* in the second relative clause refers to the college in the first relative clause. The whole term has a reading in which the college needs not to be the same for all girls. Suppose that we obtained coreferentiality by means of quantifying in the term *a womans college* for him_1 in sentence (123).

(123) *Every girl who attended him_1 who gave a donation to him_1 was put on the list.*

In that production process a reading would be obtained with for the existential quantifier wider scope than for the universal quantifier. That is not the intended reading. In order to obtain the intended reading, a new quantification rule is introduced: quantification into a CN phrase.

9.6. Rule $S_{15,n}$:

$T \times CN \rightarrow CN$

$F_{15,n}$: Replace the first occurrence of he_n/him_n in β by α .

Replace all other occurrences of he_n by *he/she/it*, and of him_n by *him/her/it*, according to the gender of the first CN or T in α .

$T_{15,n}$: $\lambda y \alpha' (\wedge \lambda x_n [\beta' (y)])$.

9.6. END

An extensive discussion of relative clause formation will be given in chapter 8; examples in which rule S_{15} is used, will be considered in appendix 1. There also will be solved a problem that I neglected above: reduction rule RR_4 applies to the translation of terms like (120), but not to such terms with the determiners *every* or *the*.

In the remainder of this section I mention some rules which are introduced only to incorporate the complete PTQ fragment. The first rule concerns the sentence modifier *necessarily*.

9.7. $B_{S/S} = necessarily$

9.8. $necessarily' = \lambda p \square [^V p]$.

9.9. Rule S₉:
$$S/S \times S \rightarrow S$$

$$F_9: \text{concatenate } (\alpha, \beta)$$

$$T_9: \alpha' (\wedge \beta')$$

9.9. END

An example is the production of (124) which gets as its translation (127).

(124) *Necessarily John runs.*(125) $\square \text{run}_*(\text{john})$.

This example illustrates how sentence modifiers can be incorporated in the fragment. The translation of (126) is not correct since that sentence cannot mean that John always runs. For an alternative of the semantics of *necessarily* see e.g. VELTMAN 1980.

Up till now we have met sentences in the positive present tense. PTQ has rules for some other tenses as well. These rules have several shortcomings, and I will mention them without further discussion.

9.10 Rule S_{17a}:
$$T \times IV \rightarrow S$$

$$F_{17a}: \text{replace the first verb in } \beta \text{ by its negative third person singular present; concatenate}(\alpha, \beta)$$

$$T_{17a}: \neg \alpha' (\wedge \beta')$$
9.11 Rule S_{17b}:
$$T \times IV \rightarrow S$$

$$F_{17b}: \text{replace the first verb in } \beta \text{ by its third person singular future; concatenate}(\alpha, \beta)$$

$$T_{17b}: W\alpha' (\wedge \beta')$$
9.12 Rule S_{17c}:
$$T \times IV \rightarrow S$$

$$F_{17c}: \text{replace the first verb in } \beta \text{ by its negative third person singular future; concatenate}(\alpha, \beta)$$

$$T_{17c}: \neg W[\alpha' (\wedge \beta')]$$

9.13 Rule S_{17d} :
$$T \times IV \rightarrow S$$

F_{17d} : replace the first verb in β by its third person singular perfect.
concatenate(α, β)

$$T_{17d}: H[\alpha'(\wedge\beta')]$$
9.14 Rule S_{17e} :
$$T \times IV \rightarrow S$$

F_{17e} : replace the first verb in β by its negative person singular present perfect; concatenate(α, β)

$$T_{17e}: \neg H[\alpha'(\wedge\beta')].$$

9.14. END

This completes the exposition of the PTQ fragment. One should realize that the sentences we have discussed, constitute a special selection of the sentences of the fragment. Besides those rather natural examples, there are a lot of remarkable sentences in the fragment. An example is (128).

(126) *The park walks in John.*


Whether this is a shortcoming or not, depends on the opinion one has about the acceptability of (126). And how this should be dealt with, depends on the opinion one has about the question which component of the grammar should deal with such phenomena. Since these questions are completely independent of the problems we were interested in, I have not discussed this aspect. Several more fundamental aspects of the system which were not completely satisfactory, have been mentioned in the discussions. Other such aspects will arise in the discussion in later chapters, for instance in appendix . As for the main aim of the enterprise, I conclude that Montague has for the problematic sentences mentioned in section 1 indeed provided an analysis which has the desired semantic properties, and which is in accordance with the compositional framework.

CHAPTER VI

VARIANTS AND DEVIATIONS

ABSTRACT

In this chapter the impact of the algebraic framework on the design of grammars, is illustrated by considering several proposals from the literature. Most of these proposals contain details which are not in accordance with the framework. It will be shown that these proposals can be improved by adopting an approach which is in accordance with the framework, without losing the semantic effect the proposal was designed for. Other proposals present acceptable variants for certain details of the framework.



1. INTRODUCTION

On the basis of several proposals from the literature, I will illustrate in this chapter what the practical consequences are of the framework we have developed in chapters 1 and 2. Some of the proposals were already discussed in JANSSEN 1978a. The rules from the proposals will not be adapted to the way of presentation used up till now, but they are quoted in the way they were formulated in the original papers. I expect that this will cause no problems. Only the formulas of IL are sometimes adapted to our notations (e.g. $\hat{\lambda}$ instead of $\tilde{\lambda}$). Some of the proposals concern variants which are in accordance with the framework, but most are not. The objections against these proposals, however, concern in most cases only a minor detail of the paper, and my criticism should not be taken as a criticism on the paper as a whole. On the contrary, most of the papers I like very much, and that was a reason for studying them in detail. I will not consider proposals which are presented in such a way that it is evident that they are intended as a non-compositional component of the system (e.g. the indexing component for variables of COOPER & PARSONS 1976, and the interpretation strategy for pronouns of BARTSCH 1979). Rather I will discuss aspects of proposals which seem at first glance to be in accordance with the framework, but which at closer investigation appear not to be. Such examples exhibit that the practical consequences of the framework are sometimes not well understood. These examples are collected here to provide as illustrations of the framework: non-examples too can be very instructive. I hope that the examples give the reader an improved understanding of what it means to design a Montague grammar. As a matter of fact, my personal experience with the examples discussed here, was a great stimulant for the research presented in this book: discovering the foundations of Montague grammar, and investigating the practical consequences of these fundamental properties.

The structure of our framework, as developed in chapters 1 and 2, is presented in figure 1. The arrows 2,5, and 7, are homomorphisms, and the arrows 3 and 6 are derivers. The examples we will consider are grouped according to the arrow representing the component where the deviation from the framework can be located. The number of the arrow indicates the section where that group of examples will be considered.




Figure 1. The framework

The framework of Montague grammar constitutes a framework which guarantees that one is working in accordance with the principle of compositionality. Deviations from this framework are not just deviations from some arbitrary mathematical system, but from a framework that is designed with the purpose of both obeying the principle, and being at the same time as general as possible. If one violates this framework, then there is a great risk that one does not only disturb the framework, but also the underlying principle of compositionality. The ultimate consequence may be that one does not describe a semantics at all. In the discussion it will turn out that the practical examples of violations of the framework in most cases indeed yield an incorrect (i.e. unintended) semantics, or no semantics at all. In such cases the framework guides us toward a correct solution. In other cases, where the proposal did not give rise to an incorrect semantics, the principle suggests another kind of solution that is simpler than the original proposal. These aspects exhibit the value of (the formalization of) the principle of compositionality as a heuristic tool.

In the light of the above remarks, it is useful to give a characterization of what are harmful deviations of Montague's framework, and what are harmless variants. This characterization can be given at the hand of figure 1. It is harmless to change the language of which the semantics is given; to change the kind of logic used as auxiliary language, or to change the kind of meanings obtained. All algebras in the figure may be replaced by other algebras. But the algebraic relations between them should not be changed; the algebraic properties of the arrows should not be disturbed. Homomorphisms should remain homomorphisms, and derivers should remain derivers. These are the properties which guarantee that the principle of compositionality is obeyed.

2. THE USE OF SYNTACTIC INFORMATION

2.1. Introduction

Some proposals from the literature contain a translation rule which depends on the actual expression on which the syntactic rule operates. This means that there are different semantic operations for the various syntactic possibilities. Hence there is a one-many correspondence between the syntactic operations and the semantic operations. Then the mapping from the syntactic algebra to the semantic algebra cannot be a homomorphism. Consequently the framework is not obeyed: the relation indicated in figure 1 by arrow 2 has to be a homomorphism. But also the principle of compositionality itself is violated. In this situation the meaning of the compound expression is not determined by the information which syntactic rule is used and what the meanings of the parts of the expression are, but also information about the actual expressions operated upon is needed. This situation is not a source of great practical problems, since, at least in the examples considered below, the rule can easily be reformulated in such a way that the framework is obeyed.

2.2. Easy to please

This example concerns a variant of Montague grammar proposed in PARTEE 1973. The expressions generated by the grammar contain labelled brackets which indicate the syntactic structure of the expressions. Partee wants to account for the occurrence of verb phrases in conjunctions and infinitives. Examples are given in (1) and (2)

(1) *Few rules are both explicit and easy to read.*

(2) *John wishes to see himself.*

For the production of these sentences a rule called 'derived verb phrase rule' is used. The rule is so close to a correct formulation that I would not like to call it a violation of the framework. It is rather an illustrative slip of the pen.

Derived verb phrase rule (PARTEE 1973)

If $\phi \in P_t$ and ϕ has the form ${}_t[{}_T[he_i]_{IV}[\alpha]]$, then $F_{104}(\phi) \in P_{IV}$, where $F_{104}(\phi) = \alpha'$, and α' comes from α by replacing each occurrence of he_i , him_i , $him_i self$ by he^* , him^* , $him^* self$ respectively.

Examples:

$F_{104}(he_1 \text{ sees } him_1 self) = \text{see } him^* self$

$F_{104}(he_7 \text{ is easy to please}) = \text{be easy to please.}$

Translation rule

If $\phi \in P_t$ and ϕ translates into ϕ' , then $F_{104}(\phi)$ translates into $\lambda x_i \phi'$.

From the formulation of the translation rule it might not be evident that the translation rule uses syntactic information. But this becomes clear if one realizes that in order to decide what the actual translation is ($\lambda x_1 \phi$ or $\lambda x_2 \phi$ or ...), one needs to know the index of the first word of ϕ . So syntactic information is used. The correction of this rule is rather simple, in analogy of term-substitution in PTQ we give the syntactic operation an index as parameter: so F_{104} is replaced by $F_{104,i}$. In a later paper (PARTEE 1977a) the rule is corrected in this way.

2.3. The horse Cannonero

DELACRUZ (1976) considers expressions like *the horse Cannonero*. Such expressions belong to a category \bar{T} and they are generated by the following rule:

S3.1 If $\alpha \in B_T$ and $\zeta \in B_{CN}$ then $F_{21}(\zeta, \alpha) \in P_{\bar{T}}$, provided that whenever α is of the form he_n , $F_{21}(\zeta, \alpha) = \alpha$; otherwise $F_{21}(\zeta, \alpha) = \text{the } \zeta \alpha$.

Examples:

$F_{21}(\text{horse}, \text{Cannonero}) = \text{the horse Cannonero}$

$F_{21}(\text{horse}, he_1) = he_1$.

Translation rule:

T3.1 If $\alpha \in B_T$, $\zeta \in B_{CN}$ and α, ζ translate into α', ζ' respectively, then

$F_{21}(\zeta, \alpha)$ translates into α' if α is of the form he_n ; otherwise

$F_{21}(\zeta, \alpha)$ translates into

$$(3) \quad \lambda P \exists y [\forall x [[\zeta'(x) \wedge \lambda P \lambda z [\forall P](\wedge \lambda x [\forall z = x])(\wedge \alpha')(x)] \leftrightarrow x=y] \wedge [\forall P](y)].$$

Translation rule T3.1 depends on the form of the input expressions of the syntactic rule, so it violates the framework. An attempt to formulate the translation rule as a single polynomial in which no syntactic information is used, would require an extension of IL with an if-then-else construction, and with a predicate which discriminates on semantic grounds between variables and constants. I doubt whether the latter is possible. But a simple solution can be found in the syntax. The construction

described by Delacruz provides evidence that we should distinguish among the terms the (sub)categories Proper Names and Indexed Pronouns. Rule S3.1 applies only to the category of Proper Names, or alternatively, rule S3.1 is a partial rule which only applies to the subcategory of Proper Names. This approach describes more clearly what the situation is, than the original rule does, or a semantic reformulation would do. A final remark about the formula (3) given by Delacruz. It is not the simplest polynomial expressing the intended semantic operation. I would use instead:

$$(4) \quad \lambda P \exists y \forall x [\zeta'(x) \wedge \alpha'(\wedge \lambda z [\overset{V}{x} = \overset{V}{z}]) \leftrightarrow x=y] \wedge [\overset{V}{P}](y).$$

3. NON-POLYNOMIALLY DEFINED OPERATORS

3.1. Introduction

The algebra of formulas into which we translate, is obtained from the algebra of IL-expressions by means of restructuring this algebra. This means that new operations may be added, another type structure may be put on the elements, and old operations may be omitted. Following MONTAGUE 1970b, we require that in this process of restructuring, all operations are polynomial operations over IL. This restriction ensures that the interpretation homomorphism for IL determines a unique interpretation for the derived algebra. If one uses operations on IL expressions which are not defined as a polynomial, then there is a great risk of disturbing the homomorphic interpretation. This would mean that we have no interpretation for the derived algebra, thus we are not doing semantics at all! Therefore it is advisable to use only polynomially defined operators.

When we consider examples of operators which are not polynomially defined, it will turn out that in all cases the operator can be replaced by a polynomially defined operator which has the desired properties. Replacement of a non-polynomially defined operator by a polynomially defined one, is (in all these cases at least) a simplification. Thus the requirement of working in accordance with the framework guides us toward a simpler treatment than originally was proposed. This consequence illustrates the heuristic value of the principle of compositionality. So there is, from a practical point of view, no reason to use nonpolynomially defined operators. Theoretical aspects of non-polynomially defined operators will be discussed in section 4.

3.2. John who runs

The approach to natural language followed in BARTSCH 1979 is closely related to the approach followed in the field of Montague grammar. The differences which appear in this and the next example are that the treatment of intensions is different, and that the generated language is somewhat more abstract since it contains brackets and other auxiliary symbols. These differences do not influence the aspect I wish to discuss. Bartsch presents a rule for the formation of term phrases containing non-restrictive relative clauses. Such expressions are formed from a term and a relative sentence by the following rule (BARTSCH 1979, p.45).

S4. If α is a term and β a relative sentence, then $\beta(\alpha)$ is a term. [...].

The corresponding translation rule reads

T4. If α' is the translation of the term α and $RELT(\lambda x \beta'(x))$ is the translation of the relative clause β from S4, then $(RELT(\lambda x \beta'(x)))(\alpha')$ is the translation of $\beta(\alpha)$, and for all terms α with $\alpha' = \lambda P(\dots P(v)\dots)$ we have: $(RELT(\lambda x \beta'(x)))(\lambda P(\dots P(v)\dots)) = \lambda P(\dots \beta'(v) \& P(v)\dots)$.

The translation rule evidently is no polynomial over IL. The rule works well for the translation one usually obtains for term phrases. For *every man* the standard translation is (5), and for this case the rule is perfect.

$$(5) \quad \lambda P \forall v[\text{man}'(v) \rightarrow P(v)].$$

In case an alphabetical variant of formula (5) is used, the situation changes. Consider (6).

$$(6) \quad \lambda Q \forall \mu[\text{man}'(\mu) \rightarrow Q(\mu)].$$

Translation rule T4 as formulated above does not apply: it is not defined for this representation. Probably we have to be more liberal and consider T4 to be defined for all expression of the indicated form. But there are also formulas which are equivalent to (5) and which are certainly not of the same form. Let R be a variable of the same type as the translation of terms, and consider (7)

$$(7) \quad \lambda Q \forall v[\lambda R[R(\text{man}') \rightarrow R(Q)](\lambda P[P(v)])].$$

Rule T4 is not defined for this representation. Moreover, application of

the rule to the subexpression $\lambda P[P(v)]$ would yield a semantically incorrect result.

This discussion shows the consequence of T4 that it is no longer allowed to exchange logically equivalent formulas. The rule defines a partial function between IL formulas; it is an instruction for formula manipulation, not for compositional semantics. A reaction to these objections against a rule like T4 might be that one adds to the rule a clause stating that in case a formula is not of the mentioned form, it must be reduced to that format. This instruction obscures a lot of problems since it does not say how such a reduction is to be performed. A discussion of the problems arising with this attempt to correct in this way a non-polynomial rule, will be given in section 4.

Can the basic idea of the operation be described in a polynomial way? The desired effect is the replacement of $P(v)$ by $\beta'(v) \wedge P(v)$. This can be obtained giving $\lambda z[\beta'(z) \wedge P(z)]$ as argument of $\lambda P[\dots P(v)\dots]$. We must take care furthermore of the binding of the variable P . Thus we come to a version of T4 which is in accordance with our formalisation of the semantic compositionality principle:

T4'. Let α' be the translation of the term α and γ' the translation of the relative clause γ . Then the translation of the compound expression $\gamma(\alpha)$ is:

$$(8) \quad \lambda Q(\alpha'(\lambda z[\gamma'(z) \wedge Q(z)])).$$

One observes that it is not needed to follow the method hinted at above: to define the intended semantic operator by defining an operator on specially selected representations. The formulation of T4' uses the polynomial expression (8). It is more exact and simpler than the original formulation, and it works well for all formulas equivalent with α' or γ' .

RODMAN(1976) also considers the formation of terms containing a non-restrictive relative clause. He presents a rule which produces such terms out of a term and a sentence, where the sentence contains a suitable variable. His translation rule reads:

If $\alpha \in P_T$, $\phi \in P_t$ and α, ϕ translate into α', ϕ' respectively, then $F_{3,n}(\alpha, \phi)$ translates into $\lambda P \lambda Q [\overset{V}{P}(\wedge \lambda x_n [\phi' \wedge \overset{V}{Q}(x_n)])](\wedge \alpha')$.

This is not the simplest formulation of the polynomial. By one time λ -reduction one obtains

$$(9) \quad \lambda_Q[\alpha'(\wedge \lambda_{x_n}[\phi' \wedge \vee_Q(x_n)])].$$

One observes that this rule is almost identical with the version given above of Bartsch's rule. The differences are due to a different treatment of intensions, and the fact that Bartsch uses the intermediate stage of a relative sentence. Concerning the meaning of the relative clause construction the two solutions are essentially the same. This shows another advantage of the restriction to use only polynomials. It gives us a uniform representation of meanings, and different polynomials can be compared with each other by using known methods.

3.3. Das Mädchen gibt den Apfel dem Vater

BARTSCH (1979) represents a rule which combines an n -place verb-phrase with a term to produce an $(n-1)$ -place verb-phrase. The rule does not in advance fix the order in which the term positions should be filled in: a rule has as parameter a number indicating which position is to be filled. By varying the sequence of 'filling in' one can generate the German versions of *The girl gives the father the apple*, *The girl the apple the father gives*, etc. (the German versions all seem to be parts of correct sentences). The result of substituting the term α on the i -th place of a verb β is indicated by $(\alpha, i)\beta$. The syntactic rule reads (BARTSCH 1979, p.27)

(S1) If β' is a V^n (n -place verb) with the set of n term-places, $K, i \in K$, and if α' is a T (term), then $(\alpha', i)(\beta')$ is a V^{n-1} with the set of term-places $K - \{i\}$.

For this we write $(\alpha'_{T(i)}(\beta' V^n) V^{n-1})$.

(T1) If α'' is the translation of α' as a T, and $\lambda_{x_j, \dots, x_m} \beta''(x_j, \dots, x_m)$ with n places, the translation of β' as a V^n , then the translation of $(\alpha', i)(\beta')$ is

$$\lambda_{x_j, \dots, x_i', x_i, \dots, x_m} (\alpha''(\lambda_{x_i} (\beta''(x_j, \dots, x_m))))),$$

with x_i' as the variable that precedes x_i and x_i as the variable that follows x_i .

This rule is defined only for special representations of the meaning of the term, and, for reasons related to the ones mentioned in the previous example, it is not acceptable as a rule for compositional semantics. Again the idea behind the formulation of the rule can be formulated by means of a polynomial, thus becoming an acceptable rule and obtaining a shorter formulation:

If α'' is the translation of α' as a T and γ'' is the translation of γ' as a V^n , then the translation of $(\alpha', i)\beta'$ is

$$\lambda y_1, \dots, y_i' y_i, \dots, y_m (\alpha'' (\lambda y_i \gamma'' (y_1, \dots, y_m)))$$

with y_i' as the variable that precedes y_i and y_i as the variable that follows y_i .

3.4. Woman such that she loves him

Below we have the rule for the formation of restrictive relative clauses from PTQ (MONTAGUE (1973)). This rule reads as follows (notation adapted)

S3,n: $CN \times S \rightarrow CN$

F3,n: Replace he_n in β by *he/she/it* and him_n by *him/her/it*, according to the gender of the first CN in α ;

Concatenate $(\alpha, \textit{such that}, \beta)$

T3,n: $\lambda x_n [\alpha'(x_n) \wedge \beta']$.

This rule gives rise to an incorrect translation in case β' contains an occurrence of x_n which should not be bound by the λ -operator which is introduced by the translation rule. An example is the production presented in figure 2.




Figure 2. Incorrect binding of x_2

The translation obtained reduces to (10).

$$(10) \quad \lambda x_2 [\textit{woman}(x_2) \wedge \textit{love}_*(x_2, x_2) \wedge \textit{run}(x_2)].$$

The produced CN-phrase may be used in the production of some sentence, and

in this process *John* might be substituted for him_2 . Then the sentence contains a CN-phrase *woman who loves John*. But the translation contains (10), expressing that the woman loves herself.

In order to avoid this collision of variables, Thomason has presented the following translation rule (footnote to PTQ, THOMASON 1974, p.261, presentation adapted)

$T3',n: \lambda x_m [\alpha'(x_m) \wedge \psi]$

where ψ is the result of replacing all occurrences of x_n in β' by occurrences of x_m , where m is the least even number such that x_m has no occurrence in either α' or β' .

One observes that $T3'$ uses an operation on expressions which is not an operation of IL: the replacement of certain variables by a variable with a special index. We might try to describe the required change by means of IL operators. It is easy to obtain the replacement: λ -conversion does the job:

$T3'' \quad \lambda x_m [\alpha'(x_m) \wedge \lambda x_n [\beta'](x_m)]$.

Where m is as in $T3'$

It is, however, impossible to extend IL with a operator Gr which yields the greatest non-used even index. This can be shown by providing two equivalent formulas for which this operator would yield non-equivalent results. Let ϕ be an arbitrary formula. $Gr(\phi \wedge x_2 = x_2)$ would be x_4 , whereas $Gr(\phi \wedge x_4 = x_4)$ would be x_6 , what contradicts the law concerning substitution of equivalents.

We just observed that Thomason's rule contains instructions which essentially use the particular IL representation of the meaning of the relative clause. Nevertheless the rule as a whole is correct in the sense that it corresponds with an operation on the set of meanings. If the translation of the common noun or of the sentence is replaced by an equivalent formula, the application of $T3'$ (or $T3''$) gives for all cases an equivalent result. This is due to the fact that the syntactic operation we called Gr is used only in the context of renaming bound variables. So $T3'$, although violating the framework, does not violate the compositionality principle.

But there is a more direct solution to the problem raised by Montague's rule. A translation rule for relative clause formation which does obey the restriction of using polynomially defined operators is

$T3''' \quad \lambda P \lambda x_n [P(x_n) \wedge \beta'](\alpha')$.

This translation rule yields a correct result for the problematic case presented in figure 2, due to the conditions for λ -conversion. In case α' does not contain free occurrences of x_n , then $T3'''$ reduces to $T3$, otherwise

λ -conversion evokes change of bound variables. One observes that the formulation of $T3'''$ is simpler and much more elegant than the formulation of $T3'$ (or $T3''$). Moreover $T3'''$ is in accordance with the *variable principle*, whereas $T3''$ and $T3'$ are not (see chapter 8). The simple formulation of $T3'$ is possible because the syntactic problem of collision of variables is not dealt with in the translation rule, but on a more appropriate level: in the rules for λ -conversion which, by their nature, are syntactic operations on IL expressions.

4. OPERATORS DEFINED ON REPRESENTANTS

In all examples from section 3, a rule was defined which works well in the situations one usually meets in practice. In two of the examples the rule does not work well in unusual situations. Often one is tempted to design rules in such a way that as a consequence they have this character. One defines a rule for the formulas one is familiar with, using well known properties of these formulas. Then one hopes that an operation defined on these special formulas in fact defines an operation on the associated meanings. In the present section it will be investigated under which circumstances this hope will become reality. It will be shown that it is not easy to create such circumstances.

Besides the practical motivation given above for considering non-polynomially defined operators, there is a theoretical argument. In the introduction of section 3 I mentioned that an operator which is not defined by means of a polynomial over IL violates the framework, and bears the risk of violating the compositionality principle itself as well. But not all non-polynomial operators do so. In 3.4 we have met a non-polynomially defined operator which could be replaced by a polynomially defined one. From chapter 2, section 7, we know that an operator which is defined on the language IL, and which respects all homomorphic interpretations, can be described by means of a polynomial. But this does not imply that all non-polynomially defined operators which respect the interpretation of IL, indeed can be described by means of a polynomial. This is due to the rather strong condition of the theorem that all homomorphic interpretations are respected. We are not interested in all meaning algebras we might associate with IL, but only in some of them. We want to interpret $P(x)$ as the application of a function to an argument, we want the interpretations of $\phi \wedge \psi$ and of $\psi \wedge \phi$ to be the same, and we want several meaning postulates to be

satisfied. Therefore the theorem does not give us the guarantee that every operation on IL which respects the usual interpretations of IL indeed can be defined by means of a polynomial. These considerations constitute a theoretical argument for considering non-polynomially defined operators. But the practical argument given above is, in my opinion a more interesting reason for doing so.

The definition of an operation on IL formulas is acceptable if (and only if) it does not disturb the compositionality principle, i.e. if with the operation on formulas we can associate an operation on meanings. This can only be done if for logically equivalent formulas the operation yields equivalent results. So when defining an operation by defining it for special formulas, every special formula ϕ has to be considered as a representant of the class of formulas equivalent to ϕ .

A mathematical description of the situation is as follows. The set of formulas (of a certain type) is divided into equivalence classes. A class consists of formulas which have the same meaning in all models. Remember that we defined the meaning of an IL formula of type τ as a function which assigns to an index and a variable assignment some element in D_τ (so expressions in the same class represent the same function). In each class representants are defined, and we wish to define an operation on the whole class by defining an operation for the representants. If in each class there is only one representant, we are in the situation presented in figure 2b, if there are more, then we are in the situation of figure 2a. We want to know when a definition on a representant defines an operation on the whole class.




Figure 2a Several representants




Figure 2b One representant

When defining an operation on an equivalence class by a definition on its representant, two aspects can be distinguished.

A) the definition of an operation on the representants

B) A proof that this defines an operation on the whole class.

As for A) we have to fulfill the following two requirements.

- A1) One has to describe exactly the set of formulas for which the operation is defined, i.e. one has to define a *recursive* set of representants.
- A2) One has to define for all representants what the effect of the operation is, i.e. we have to define an *effective* operation on the set of representants.

This kind of requirements we have met before: define a set and operations on this set (e.g. define a language and a translation of this language). Therefore it seems attractive, in the present context, to define the set of representants by means of a grammar generating the expressions in the subset. In order to be sure that the operation is defined for all formulas in the subset, one might formulate the clauses of the operation parallel to the grammatical rules generating the subset. This will probably be more complicated than a polynomial definition. But other techniques for defining the set of representants and the operation are possible as well.

As for B) I will consider first the situation described in figure 2a: one representant in each class. Here the definition of an operation on the representant automatically determines a semantic operation on the whole class: the interpretation of the operation on the representant is the semantic operation on the interpretations of all expressions in the class. But how can we be sure that we are in the situation of figure 2a? Proving that we are in such a situation means that we have to prove the existence and unicity of a representant for each class. I do not know whether there exists for each type a *recursive* set of unique representants. Assuming the possibility of such a set, it remains the question how to prove the existence and unicity of such representants. It seems attractive to do this by providing an algorithm which transforms a given formula into the representant of its equivalence class. This expresses the idea we met in section 3.2: if a formula is not in the required form, it should be transformed into the required form. This approach is, however, in the relevant cases impossible, as follows from the following theorem.

4.1. THEOREM. Let $\sigma, \tau \in \text{Ty}$. Define the equivalence relation \sim on $\text{ME}_{\langle \sigma, \langle \tau, t \rangle \rangle}$ by $\phi \sim \psi$ iff $\models \phi = \psi$. Let $R \subset \text{ME}_{\langle \sigma, \langle \tau, t \rangle \rangle}$ be a (recursive) set of representants such that for each equivalence class there is one element in R . Let $f: \text{ME}_{\langle \sigma, \langle \tau, t \rangle \rangle} \rightarrow \text{ME}_{\langle \sigma, \langle \tau, t \rangle \rangle}$ be a function which assigns to each formula the representant of its equivalence class. Then f is not recursive. The

same result holds if the expressions are from $ME_{\langle \tau, t \rangle}$.

PROOF. Let $\phi \in ME_{\tau}$, $P \in VAR_{\sigma}$, $Q \in VAR_{\tau}$. Then the following statements are equivalent

$$(11) \quad \models \phi \quad (\phi \text{ is logically true})$$

$$(12) \quad \models \lambda P \lambda Q [\phi] = \lambda P \lambda Q [\forall x [x=x]]$$

$$(13) \quad \models f(\lambda P \lambda Q [\phi]) = f(\lambda P \lambda Q [\forall x [x=x]]).$$

Note that in (13) semantic equality of the formulas is expressed. Since for each class there is one representant, (13) is equivalent with (14)

$$(14) \quad f(\lambda P \lambda Q [\phi]) \equiv f(\lambda P \lambda Q [\forall x [x=x]]).$$

Note that in (14) syntactic identity of the by f obtained formula is expressed. So, if f is recursive, the question whether ϕ is logically true is decidable: calculate $f(\lambda P \lambda Q [\phi])$ and $f(\lambda P \lambda Q [\forall x [x=x]])$, and see whether they are identical. Since IL is undecidable, f cannot be recursive. For $ME_{\langle \tau, t \rangle}$ analogously. Note that we did not use the recursiveness of the set of representants.

4.1. END

The translations of expressions of the PTQ fragment are all of the form $\langle \sigma, \langle \tau, t \rangle \rangle$ or $\langle \tau, t \rangle$. The same holds for the expressions of the fragments considered in all examples. The theorem says that there is no algorithm which transforms a formula into its representant. If one tries to define an operation on a class by an operation defined on its representants, then one has to find some other way of proving the existence and uniqueness of a representant.

Next we will consider the situation described in figure 2b: a multiple set of representants is allowed for. There is no doubt that such a set exists: ME_{τ} itself is a recursive set of multiple representants of ME_{τ} . But also here a complication arises.

4.2. THEOREM. Define \sim as in the previous theorem. Let R be a (recursive) set such that for every equivalence class there is at least one equivalent element in R . Let $f: ME_{\langle \sigma, \langle \tau, t \rangle \rangle} \rightarrow ME_{\langle \sigma, \langle \tau, t \rangle \rangle}$ be a recursive function that

assigns to every formula an equivalent formula in R. Then for $r_1, r_2 \in R$ it is undecidable whether $\models r_1 = r_2$.

PROOF. As observed in the proof of the previous theorem $\models \phi$ is equivalent with $\models f(\lambda P \lambda Q[\phi]) = f(\lambda P \lambda Q[\forall x[x=x]])$. If equality is decidable for elements of R, then the equality is decidable for these two formulas, so it is decidable whether ϕ holds. This gives rise to a contradiction since IL is undecidable. Note that we did not use the recursiveness of the set of representants.

4.2. END

This result means that we have to prove that an operation defined for representants yields for equivalent formulas an equivalent result, without knowing what the equivalent formulas look like.

The above discussion shows that it is, generally spoken, a complicated and extensive task to define a function by defining a function on specially selected representations. Probably this can only be done in practice, if one considers a situation with a special structure which has the effect that all proofs become drastically simplified. But if the situation is such a special one, it may be expected that the same effect can be obtained in a more direct way: by using polynomials. This is illustrated in the examples considered in section 3. So far there is no evidence that there is any advantage in defining operations in a non-polynomial way.

5. NEW SYMBOLS IN IL

5.1. Introduction

Some authors extend the language of IL with new symbols. These symbols should obtain an interpretation by means of an extension of the interpretation homomorphism for IL. For each point of reference some semantic object of the right type has to be associated with the new symbol. If the new symbol is an operator, its interpretation has to be a function operating on the interpretation of its argument. If these requirements are not met, then the interpretation homomorphism of IL cannot be extended to an interpretation homomorphism for the extension of IL. Consequently arrow 5 in figure 1 is no longer a homomorphism. Hence arrow 7 is not a homomorphism either. Then the translation homomorphism 2 cannot be continued to an interpretation homomorphism, and this means that the principle of compositionality

is violated. Below we will consider two examples of new symbols.

5.2. Shake John awake

DOWTY (1976) treats, among others, the semantics of factive constructions such as *shake John awake*. In order to do so, he extends the language of intensional logic with two operators: CAUSE and BECOME. Interesting for our discussion is his treatment of CAUSE. In order to define its interpretation Dowty adds "to the semantic apparatus of PTQ a selection function f that assigns to each wff ϕ and each $i \in I$ a member $f(\phi, i)$ of I . [Intuitively $f(\phi, i)$ is to be that i' most like i with the (possible) exception that ϕ is the case [...]]". Then the interpretation of CAUSE reads:

"If $\phi, \psi \in ME$, then $(\phi \text{ CAUSE } \psi)^{A, i, j, g}$ is 1 if and only if $[\phi \wedge \psi]^{A, i, j, g}$ is 1 and $[\neg \psi]^{A, f(\neg \phi, i), j, g}$ is 1."

The function f is defined on IL-expressions and not on the interpretations of these expressions. As a consequence CAUSE is an operator on IL-expressions and not on the meanings they represent. This is illustrated as follows. The definition of f allows that for some ϕ, η, i holds that $f(\neg[\phi \wedge \eta], i) \neq f(\neg[\eta \wedge \phi], i)$. This may have as a consequence that $[(\phi \wedge \eta) \text{ CAUSE } \psi]^{A, i, j, g} = 1$ whereas $[(\eta \wedge \phi) \text{ CAUSE } \psi]^{A, i, j, g} = 0$. The main features of an example of such a situation are as follows. Let $[(\phi \wedge \eta) \wedge \psi]^{A, i, j, g} = 1$, so $[(\eta \wedge \phi) \wedge \psi]^{A, i, j, g} = 1$. Suppose that $f(\neg[\phi \wedge \eta], i) = i'$ and $[\neg \psi]^{A, i', j, g} = 1$. Then $[(\phi \wedge \eta) \text{ CAUSE } \psi]^{A, i, j, g} = 1$. Suppose moreover that $f(\neg[\eta \wedge \phi], i) = i''$ and $[\neg \psi]^{A, i'', j, g} = 0$. Then $[(\eta \wedge \phi) \text{ CAUSE } \psi]^{A, i, j, g} = 0$.

In the above example the principle of compositionality is not obeyed: two equivalent formulas cannot be interchanged 'salva veritate'. Moreover the meaning of CAUSE described above is incorrect since, intuitively, $[(\phi \wedge \eta) \text{ CAUSE } \psi]^{A, i, j, g} = [(\eta \wedge \phi) \text{ CAUSE } \psi]^{A, i, j, g}$. A correction is possible by taking as domain for f the intensions of formulas: f assigns to each $d \in D_{\langle s, t \rangle}$ and $i \in I$ a member $f(d, i) \in I$. Then a situation as described above is automatically excluded. The interpretation of CAUSE now becomes as follows.

$$\begin{aligned} [\phi \text{ CAUSE } \psi]^{A, i, j, g} &= 1 \quad \text{if and only if} \\ [\phi \wedge \psi]^{A, i, j, g} &= 1 \quad \text{and} \quad [\neg \psi]^{A, \underline{i}, j, g} = 1, \quad \text{where} \\ \underline{i} &= f([\wedge \neg \phi]^{A, i, j, g}, i). \end{aligned}$$

This definition has the property that if ϕ CAUSE ψ , then for all tautologies η holds that $(\phi \wedge \eta)$ CAUSE ψ , a problem of the same nature as the problem we met in chapter 4 concerning the complements of belief-sentences.

5.3. I and You

GROENENDIJK & STOKHOF (1976) give a treatment of the pronouns *I* and *You*. For this purpose, they extend the model for IL. Usually the denotation of an IL expression is defined with respect to a world i and a time j ; these i and j are called 'indices'. Groenendijk & Stokhof extend the set of indices with three components: j_0, s and h . Here $j_0 \in J$ is the moment 'now', i.e. the moment of utterance, $s \in A^{I \times J}$ is a function which for a point of reference (i, j) yields the speaker at that moment, and $h \in A^{I \times J}$ is a function yielding the hearer. The interpretation of α may depend on i, j, j_0, s and h , and we write $\alpha^{A, i, j, g, s, h, j_0}$ for the interpretation of α . The language of IL is extended with the constants i and y of type $\langle s, e \rangle$; these constants occur in the translations of the pronouns *I* and *you* respectively. The goal they wish to reach is described as follows. (op.cit.p.308). 'What we want our interpretations to express is that the extension of the constants i, y are the possible individuals which are speaking now, spoken to now respectively. This would explain the tautological character of a sentence like (15) and the contingent character of sentences like (16)'.:

(15) *I am the speaker*

(16) *I will not be the speaker*

Groenendijk & Stokhof define $F(i) = s$ and $F(y) = h$. Furthermore they define

$$i^{A, i, j, g, s, h, j_0} = F(i)(i, j_0) \quad (=s(i, j_0))$$

and

$$y^{A, i, j, g, s, h, j_0} = F(y)(i, j_0) \quad (=h(i, j_0)).$$

So for any point of reference the interpretation of i is the speaker now, and the interpretation of y is the hearer now.

The corresponding intensions, however, are separately defined: as $(\hat{i})^{A, i, j, g} = F(i)$ and $(\hat{y})^{A, i, j, g} = F(y)$ respectively. One observes that no longer holds that for all α : $(\hat{\alpha})^{A, i, j, g} = \lambda(i, j) \alpha^{A, i, j, g}$. This combination of the definition of interpretation of \hat{i} and \hat{y} with the interpretation of i and y violates the recursive interpretation of the IL, thus

disturbing its homomorphic interpretation. This has drastic consequences: several tautologies become invalid. It is no longer true that for constants of type $\langle s, e \rangle$ holds that $\bigwedge^V c = c$, nor that $\alpha = \beta \rightarrow \bigvee^V \alpha = \bigvee^V \beta$ is valid (for $\alpha, \beta \in ME_{\langle s, e \rangle}$). The interpretation of the logic is not a homomorphism (since $h(\bigwedge^V \alpha) \neq \bigwedge^V h(\alpha)$); therefore the interpretation of the natural language is not a homomorphism either. This means that the principle of compositionality is violated.

Let us consider the first goal of the approach of Groenendijk & Stokhof. Sentence (15) is true when evaluated with respect to the moment 'now', but not with respect to a point of reference where the speaker is someone else. The sentence expresses not a tautology (as a matter of fact, this is not claimed by Groenendijk & Stokhof). What they probably wish to express by the phrase 'tautological character' is that for every choice of the moment 'now', the sentence is true when evaluated with respect to this moment, whereas not the same can be said about the second sentence. This effect can be obtained in a compositional way by stipulating that $F(i) = \lambda i, j [s(i, j_0)]$ and $F(y) = \lambda i, j [h(i, j_0)]$. Then the translation of (16), being something like $\bigvee^V i = \bigvee^V s$, becomes true for the point of reference (i, j_0) , no matter what j_0 is, whereas this is not the case for the translation of (16), being something like $\neg \forall^V [i = s]$.

6. COUNTING ELEMENTS

6.1. Introduction

In the present section I will consider two examples of counting the number of elements in the model. In the first example this is done in a way which suggests a misunderstanding of the framework. As a contrast I present the second example in which the counting proceeds correctly. These examples illustrate the role of the derived algebra M' which is obtained from the algebra M in which we interpret intensional logic.

6.2. Keenan & Faltz count

KEENAN & FALTZ (1978) present a system for the description of syntax and semantics that is related to Montague's system. An important difference is that they obtain their semantic domain by application of algebraic operations (join, meet, homomorphism) on certain basic elements. One way in which they compare their system with Montague's is by ways of

counting the number of elements in the semantic domain of their system and Montague's sets D_τ . They base an argument in favour of their system on the fact that a certain domain D_τ contains many more elements than the corresponding set in their own system. There are several objections against this comparison. The stage at which Keenan & Faltz carry out their comparison (viz.p.130) does not do justice to Montague's enterprise. They compare their model for an extensional fragment of English with Montague's domains developed for an intensional fragment. Furthermore they do not take Montague's meaning postulates into account. So the numbers they obtain are not the relevant numbers. I will, however, not correct their calculations, since I am primarily interested in the method of comparison. This method will be discussed below.

Keenan & Faltz have a theory which says e.g. how many verb phrase meanings are possible (for a given domain of individuals): it is the number of homomorphisms between certain sets (which are built from the set of individuals). Keenan and Faltz count in their framework the number of elements in some of such sets, i.e. they count the number of possible denotations of certain types. In Montague's framework they count the number of elements in D_τ for the corresponding types. I have fundamental objections against this comparison since in this way sets are compared that are incomparable. The sets D_τ in Montague's system are sets in the algebra M (see figure 1). Out of algebra M a derived algebra is defined. This derived algebra M' consists precisely of the elements which are used for the interpretations of expressions produced by the grammar for the fragment. In the process of forming the derived algebra M' elements of D_τ may be thrown out; e.g. a set D_τ may consist of all functions of a certain type, whereas in M' only the homomorphisms may be left over. If one wants to count the number of possible denotations for the expressions of a certain category, then one has to count the number of elements of the corresponding type in the union of all derived algebras. One should not count the auxiliary set D_τ instead. The method of counting of Keenan & Faltz neglects the role of arrow 6 in figure 1.

The number of elements in a derived algebra can easily be counted. The derived algebra M' is the image of the syntactic algebra for the fragment. Therefore the number of elements in M' cannot be larger than the number of expressions in the syntax. Since the latter is denumerable, the former is. And for a given category, say the verb phrases, the number of expressions

of this category gives an upperbound for the number of elements of the corresponding type in the semantic algebra.

6.3. Partee counts

As a contrast to the previous example, I would like to consider PARTEE (1977b), where the difference between M and M' is taken into account. Partee discusses the psychological plausibility of possible world semantics. She argues that the finiteness of our brains requires that the theory of linguistic information we have should be finitely representable. The possible world semantics, however, gives rise to sets of rather large cardinalities (For instance if $|A| = \aleph_0$ and $|I| = \aleph_0$, then $|D_{\langle s, e \rangle}| = 2^{\aleph_0}$ and $|D_{\langle \langle s, e \rangle, \langle s, e \rangle \rangle}| = 2^{(2^{\aleph_0})}$). These cardinalities make it impossible to assume that we have finite representations of all sets D_τ in our brains. Partee gives a way out of this dilemma: assume that we have finite representations of the form of the sets D_τ , but not of all their elements. PARTEE (1977b, p.317-318) says: 'In the acquisition of any particular language, not all of the in-principle possible denotations need to be considered as a hypothesis about the actual denotation of an actual expression of the language. The intensional logic into which the natural language is translated, will contain at most denumerable many expressions of any given type, and the finite perceptual and cognitive apparatus will make at most denumerable many members of $D_{a,A,I,J}$ finitely representable, and it will only be correspondences between these two at most denumerable sets that will have to be empirically determined by the language learner'.

It is striking to compare these psychologically motivated opinions with the mathematical properties of the framework. These predict that in an interpretation of a particular language there are only denumerable many meanings because there are denumerable many expressions in the language of which we give the meaning. So for any particular language the number of meanings in the model, and the number found on psychological considerations agree.

In relation with the previous discussion Partee considers the following question (PARTEE 1977b p.318). 'One might ask at this point, if the 'available' members of $D_{a,A,I,J}$ are always going to form a denumerable set, why shouldn't all the sets $D_{a,A,I,J}$ be constrained to be denumerable by the semantic theory?' The answer Partee would argue for is 'that there is no telling in advance which possible world the native speaker will find her-

self in; [...] her semantic component must equip her for a language in any of them.

An alternative is to consider the psychological arguments as an invitation to change one of the algebras of the framework in figure 1. It seems to be an argument against taking the standard model for intensional logic because of the large cardinality of its sets, in favor of taking as semantic algebra some generalized model with denumerable many elements (generalized model in the sense of HENKIN 1950, see sections 1 and 3 of ch.3). This would have the interesting consequence that the axiomatization of intensional logic (given in chapter 3) would be a complete axiomatization for this class of models. This is a direct consequence of lemma 3.3.1 in GALLIN 1975. But no matter which conclusion is drawn from the psychological arguments, this whole discussion remains an interesting excursion because Montague's system was not designed, as I explained in chapter 1 and 3, to formalize any psychological theory.

7. THE TRANSLATION LANGUAGE

7.1. Introduction

An intensional language is a language of which the denotation of an expression depends on an index, and an extensional language is a language where this is not the case. An example of an extensional language is predicate logic, an example of an intensional language is IL. Our approach makes English an intensional language: a sentence denotes a truth value; which one this is depends on the current index (point of reference). In an extensional approach we would say that a sentence denotes an function from indices to truth values (an intension). Is it possible to give an extensional treatment of English, and what are the consequences? In other words, is it possible to change the relation indicated in figure 1 by arrow 7?

It will turn out that the answer to the above question is positive. This gives us the choice between (at least) two different approaches. When making a choice, we have to realize what the role is of the translation level in the whole framework. We aim at defining a systematic relation between English expressions and their meanings. In order to be able to express this relation conveniently, we use the translation into intensional logic. In chapter 2 is explained how this logic is interpreted

homomorphically: e.g. an expression of type t has as its homomorphic image (has as its meaning) some function in $\{0,1\}^{I \times J \times G}$. Fundamental to this whole approach is the relation between expressions and meanings. If a translation into an extensional language gives rise to the same relation, it is acceptable as well. Another translation is just another tool, and a choice has to be made on the basis of technical arguments.

7.2. Hausser translates

As an introductory step of treating English as an extensional language, I consider an approach which is very close to the PTQ translation: translate into an IL expression denoting the meaning of that expression. Let the intensional translations of two sentences be A and B respectively. Then the translation of their conjunction has to be $\hat{A} \wedge \hat{B}$. Most of the translation rules have the format $\alpha(\hat{\beta})$. With the new translations this becomes $\hat{A}(\hat{B})$. These examples illustrate that this approach does give rise to somewhat more complex formulas. A next step is to use a logical language in which the operators on elements of type $\langle s, \tau \rangle$ are defined. For instance \wedge , where $A \wedge B$ is interpreted as the complex conjunction formula given above, so as indexwise evaluation of the parts of the conjunction. For function application is used $A(B)$; to be interpreted as denoting the same as $\hat{A}(\hat{B})$. Such operators are used in JANSSEN & VAN EMDE BOAS 1977 for dealing with semantics of programming languages.

Following TICHY 1971, HAUSSER (1980) argues for an extensional approach to English. In HAUSSER (1979a, 1984) this idea is worked out. He does not use the standard logical operators (e.g. conjunction on truth values) any more, and this gives him the opportunity to use the standard symbols with a new meaning. Now $\phi \wedge \psi$ means indexwise valuation of the parts of the conjunction, and $\alpha(\beta)$ is the variant of function application which is described above as $\alpha(\hat{\beta})$. In this way one obtains a simplification of the formulation of the translation rules since no intension symbols have to be used. The price one has to pay for this, is that the logical symbols obtain a somewhat deviant interpretation, what is the normal price, and what is quite acceptable. But the presentation of the translation rules is not the only aspect of a new translation. What happens if one wishes to take the meaning postulates into account, or if one wishes to simplify the formulas one obtains? To understand the dangers, one should realize that the new translation causes the intension operators to be invisible whereas semantically

they still are there. Therefore new reduction rules have to be found. In HAUSSER (1979b,1984) indeed a simple translation is obtained but not all meaning postulates are expressed in the translation. Therefore his results are not convincing, and further investigations are required before it is clear whether this extensional approach gives a simplification.

7.3. Lewis translates

A simplification I expect from the approach in LEWIS 1974. He discusses the consequences of another kind of extensionalized translation. He considers using the extra possibilities given by an extensionalized translation: namely the possibility to relate to an expression a meaning that is not an intension (i.e. not a function in $D^{I \times J \times G}$). The verb *run* gets in the PTQ approach a translation of type $\langle\langle s, e \rangle, t \rangle$. In the Hausser approach it is translated into an expression of type $\langle s, \langle\langle s, e \rangle t \rangle \rangle$. But in the Lewis approach its translation would be of type $\langle\langle s, e \rangle, \langle s, t \rangle \rangle$. So the translation of *run* would be a function which assigns to an individual concept a proposition. In this way one gets rid of the remarkable non-constant interpretation of constants of IL, where $run^{A, i, j, g} = F(run)(i, j)$. In Lewis approach it would be just $run^{A, i, j, g} = F(run)$. The translation of *He₁ runs* would be $run(x_1)$ being an expression of type $\langle s, t \rangle$. Note that here the function application has its standard meaning. This illustrates the advantage of Lewis approach, but further investigations are required in order to decide whether it is a real simplification. A remarkable aspect of Lewis approach is that it gives a completely different relation between expressions and meanings than we considered up till now, so investigating these matters here, might bring us far from the current work (cf. HAUSSER 1984, p.82,83).

7.4. Groenendijk & Stokhof translate

A last version of what might be called an extensionalized translation is used in GROENENDIJK & STOKHOF 1981. They do not translate into IL, but into Ty2 (see GALLIN 1975). Such a translation can be called extensional since the interpretation of a Ty2-expression does not depend on an index, but only on the variable assignment (including the assignment to 'index' variables). Also in this translation we get rid of the non-constant interpretation of constants since the index dependency of predicates as *run* is made explicit by translating *run* into an expression containing an index-variable. The phenomena described by Groenendijk & Stokhof seem to require

the expressive power of Ty2, and it is to be expected that this power will be required for the treatment of other phenomena as well (e.g. VAN BENTHEM (1977) argues that explicit reference to moments of time is needed for tense phenomena). Since we will not consider these phenomena, we will not investigate the details of such a translation. From the way in which we introduced IL in chapter 3 (using a translation into Ty2), it is evident that this would cause no problems at all (on the contrary, several aspects would become simplified).

7.5. Keenan & Faltz on translations

In KEENAN & FALTZ 1978, several requirements are given concerning the logical form of a natural language, e.g. criteria concerning the correspondence between a natural language expression and its logical form. They criticize the logical form which is obtained in a Montague Grammar. An example is their comment on the translation of *John* which is in an extensional fragment $\lambda P[P(j)]$. They say (p.18) '... this assignment of logical structure fails the Constituent Correspondence Criterion, since it contains three logical elements, namely j , P and λP , none of which corresponds to a constituent of John.' Such criticism plays an important role in the argumentation in favour of their framework.

The argumentation of Keenan & Faltz is, however, based upon a misconception of the framework. I assume that they understand by 'logical form', that level of description at which the meaning of an expression is completely determined. In fact, there is no unique level of description in Montague Grammar for which this holds. The analysis tree of an expression, its immediate, unreduced translation, its reduced translation (and all the stages in between), all determine the meaning of that expression completely. That, in particular, the translation of an expression into IL cannot be claimed to have a special status as *the* logical structure of that expression, becomes clear if one realizes that this level of representation, in principle, can be dispensed with altogether. Grammar provides a correlation between syntactic structures and meanings. In Montague Grammar this is done by providing a homomorphism from the set of syntactic structures into the set of abstract settheoretical entities, modelling the meanings. In the PTQ-system this homomorphism is defined in two steps. First a homomorphic translation from syntactic structures into logical expressions is provided, second the logical expressions are interpreted, i.e. related in the usual homomorphic


way to the abstract entities defined in the model. These two homomorphisms together determine one homomorphism from the syntactic structures into the meanings, viz. the composition of the two. This two-step approach is chosen for reasons of convenience only, it is not necessary. As a matter of fact, the EFL-system (MONTAGUE 1970a) is an example of a system in which the homomorphism from syntactic structures into abstract meanings is defined in one fell swoop, without an intermediate stage of translation into a logical language. All this means that within the PTQ-framework it is not possible to talk of *the* logical structure, or *the* logical form, of an expression. So Keenan & Faltz criticize a non-existing aspect of Montague grammar.

CHAPTER VII

PARTIAL RULES

ABSTRACT

In the framework the syntactic and semantic rules are considered as algebraic operators. As a consequence of the definitions given in the first chapters, the syntactic rules have to be total. This is investigated and compared with linguistic requirements. Partial syntactic rules from the literature are considered and alternatives for them are presented. One of the methods to avoid partial rules is the use of rule schemes. It turns out that the requirement of using total rules is a valuable heuristic tool. Consequences of this requirement are compared to consequences of Partee's well-formedness constraint.



1. RESTRICTIONS OF THE FRAMEWORK

Based upon the principle of compositionality, we have developed an algebraic framework for the description of syntax and semantics. The algebras of the framework have operators: i.e. functions from carriers to carriers. This implicates that an operator can be applied without any further restriction to any element of the sorts required by the operator. In this chapter I will consider consequences of this aspect of the framework, and especially its consequences for the syntactic algebra. Some of these consequences are closely related with those of the 'well-formedness constraint', (PARTEE 1979b), which will be considered in section 6.

In linguistics one often conceives of a grammar as a generating device for producing all and only the expressions of a language. With this conception it is rather natural to think of restrictions on this production process. One might think of restrictions on the order of application of the rules. Two examples are the following. One might have rules of which the applicability depends on the way in which an expression is produced (such conditions are called 'global constraints'). One might have a filter which throws away some of the produced elements (e.g. one which filters out all expressions which contain a certain symbol). The description of the possible sequences of application of the rules constitutes an important component of a transformational grammar (for instance certain rules might be obligatory, others ordered cyclically), and filters are also often used in that field. If one wishes to use the syntactic knowledge from the field of transformational grammar in the field of Montague grammar, then one is tempted to incorporate these restrictions on the generation process in Montague grammars. Would that be possible, and at what price?

In our framework the syntax has to be a many-sorted algebra, i.e. a set of carriers with operations defined on these carriers. An algebra is not a generating device, it rather is the description of a situation. By describing what the syntactic algebra is, it is said what the relevant expressions are, and what the functions are which describe the relations between these expressions. The expressions can be determined in any way one likes, and nothing has to be said about their production. One might for instance define an algebra by mentioning all the elements and describing the effects of all operators (we did so in the beginning of chapter 2). A simpler method is to give a collection of generators, and tell what the operators are. Several choices of generators may be possible, one more

clever than the other. But no matter how the algebra is defined, the elements remain the same elements, the operators remain the same operators, and the algebra remains the same algebra.

The operators of the algebra are mappings from carriers to carriers. The range of an operator (the expressions obtained as results of an application of the rule) consist by definition of elements of a certain carrier. Therefore it is in our framework not possible to have a filter which says that certain outcomes are not acceptable. The domain of an operator (the expressions it operates upon) is some n -tuple of carriers. How we obtained the information that an expression belongs to a carrier is of no influence. The applicability of an operator cannot depend on the information which rules were applied previously, because there are no 'previously applied rules' for an element of an algebra. For this reason, there cannot be a prescribed ordering on the rules, there cannot be rules that are explicitly required to be used in all derivations, and the derivational history cannot influence the applicability of the rules.

Of course, the generation of expressions is an important aspect of syntax, and therefore we paid special attention to it. The notion of a generated algebra was defined, and theorems were proved about such algebras. In a generated algebra it might be meaningful to speak about filtering, ordering of the application of rules, the influence of derivational history, and obligatory rules. But if we would allow this, we would describe a generation mechanism, and not operators of an algebra: in an algebra there is no place for such aspects. So this discussion brings us to reject certain methods which are customary in the tradition of transformational grammars. But the rejection only concerns the method, not the ideas. It is possible to organize an algebra in such a way that the same effects are obtained in another way. Below I will give some examples.

An explicit ordering of rules is not possible in an algebra. But in a generated algebra there is a certain natural ordering among the operators. If an operator R takes as its argument an expression of category C , then the operators which yield expressions of the category C are used before R . In this way the categorial system of the algebra has as effect a certain implicit ordering of the operators.

If one wants a certain ordering on the rules, this effect can be obtained by a suitable refinement of the categorial system. Let R_a and R_b be two rules, both operating on sentences and yielding sentences. Suppose that

we want R_a to be applied precisely one time before R_b . This effect can be obtained by distinguishing among the sentences two categories: S_1 and S_2 . Here S_1 is the category of sentences to which R_a has not yet been applied, and S_2 of sentences to which R_a has applied. Then R_a can be defined as a rule which operates on expressions of category S_1 and yields expressions of category S_2 , whereas R_b is defined to operate on expressions of category S_2 , yielding expressions of this category again. The definitions of the other rules have to be adapted for these categories as well. I expect that by means of a refined categorization system the effect of any ordering can be obtained. Since in the field of Montague grammars explicit rule ordering hardly is employed, I will not consider this topic any further.

As explained above, the applicability of a syntactic rule to an expression cannot depend on the derivational history of that expression. Notice that, on another level, we already met a situation where it was important to have derivational histories available. The meaning of an expression may depend on the derivational history of that expression. We did not define the translation homomorphism on the algebraic grammar for a language because in that grammar such histories are not available. The translation homomorphism is defined on the associated term-algebra, i.e. the algebra of derivational histories. This suggests us what to do when derivational histories would be important in syntax: use an algebra in which the elements represent derivational histories. But in the field of Montague grammars I know of only one rule which uses information about the derivational history (rule 3 of THOMASON 1976), so the issue does not seem to be important. Moreover, this aspect of Thomason's rule can probably be avoided by following the proposal of PARTEE (1973) to let a grammar produce not unstructured strings, but labelled bracketings. For these reasons the role of derivational histories in the syntax will not be considered here any further.

Above we have considered some restrictions on the circumstances in which a rule may be used. The conclusion was that such rules violate basic aspects of our framework. Another request from linguistics is to allow restrictions on the expressions to which a rule is applied. In the field of transformational grammars it is standard to put conditions on the possible inputs of a transformation. In the field of Montague grammar many rules are proposed as well which do not apply to all expressions of the category required by the rule, but only to some of them. In the field of semantics one has proposed to use functions which are not defined for all arguments of the required type (see section 2.2). In contrast to the constraints on

applicability discussed above, one might argue that our framework should allow for operators which are not defined for all arguments of the required sort. Such partial operators are known in the theory of universal algebras; the algebras in which they occur are called partial algebras. In the next sections it will be investigated whether we could be more liberal than we have been, and whether we should allow for partial algebras within our framework.

2. PARTIAL ALGEBRAS

2.1. Partial grammars

Contrary to what one might expect, it is not just a minor variation of the system to allow for partial algebras (i.e. algebras with partial operations). Such a step would disturb important parts of theory we have developed so far. I will illustrate this by means of two examples which show that certain theorems we proved concerning properties of the syntax are not valid when partial rules are allowed. In 2.2 it will be shown that certain theorems of intensional logic lose their validity when partial operators are allowed in the logic.

2.1. EXAMPLE.

$$G = \langle \langle [\{a_A\}, \{b_B\}, \{c_C\}], \{F_a, F_b, F_c, F\} \rangle, D \rangle.$$

Here $F_a : A \rightarrow A$ is defined by $F_a(\alpha) = \alpha a$

$F_b : B \rightarrow B$ is defined by $F_b(\beta) = \beta b$

$F_c : C \rightarrow C$ is defined by $F_c(\gamma) = \gamma c$.

So by repeated application of F_a strings of arbitrary length consisting of a 's are produced. Analogously for F_b and F_c . Furthermore the partial rule F is defined as follows:

$$F: A \times B \times C \rightarrow D$$

where

$$F(\alpha, \beta, \gamma) = \begin{cases} \alpha\beta\gamma & \text{if the lengths of } \alpha, \beta \text{ and } \gamma \text{ are equal.} \\ \text{undefined} & \text{otherwise} \end{cases}$$

The language $L(G)$ generated by G is $\{a^n b^n c^n \mid n \in \mathbb{N}\}$. This is a non-context-free language (see HOPCROFT & ULLMANN 1979 example 6.1). So when partial operations are allowed in the syntax, theorem 5.6 from chapter 2 does not hold.

2.2. EXAMPLE. Let L be some recursively enumerable language over alphabet A . According to theorem 3.7 from chapter 2, there is an algebraic grammar

G such that $L(G) = L$. Suppose that $G = \langle\langle [B_s]_{s \in S}, (F_\gamma)_{\gamma \in \Gamma}, s_0 \rangle\rangle$.

Let $\sigma \in A^*$ be arbitrary, and define the algebraic grammar H_σ by

$H_\sigma = \langle\langle [B_s]_{s \in S}, (F_\gamma)_{\gamma \in \Gamma} \cup \{f\}, s_1 \rangle\rangle$

where s_1 is a new sort ($s_1 \notin S \cup \{s_0\}$), and where f is a partial operation defined by

$$f: s_0 \rightarrow s_1 \text{ where } f(\alpha) = \begin{cases} \sigma & \text{if } \alpha \equiv \sigma \\ \text{undefined} & \text{otherwise.} \end{cases}$$

Note that H_σ produces a language which is either empty (if $\sigma \notin L(G)$) or consists of σ (if $\sigma \in L(G)$). So $L(H_\sigma) \neq \emptyset$ iff $\sigma \in L(G)$.

Suppose now that it was decidable whether $L(H_\sigma) = \emptyset$; then it was decidable as well whether $\sigma \in L(G)$. Since $L(G)$ is an arbitrary recursively enumerable language, the latter is not decidable, and consequently it is not decidable whether $L(H_\sigma) = \emptyset$. This means that theorem 5.5 from chapter 2 (which states the decidability of the emptiness of $L(G)$) is not valid if we allow for partial operations.

2.2. Partial models

The following example concerns the use of partially defined operations in the semantics. They arise, for instance, if one wishes to deal with sortal incorrectness: certain combinations of a verb phrase with a subject do not fit well together, although most expressions of their categories give no rise to problems. An example (THOMASON 1972) is (1).

(1) *The velocity of light is shiny*

It is not attractive to say of such a sentence that it is 'false', since then its negation would be 'true'. Either, one should consider (1) as being syntactically incorrect, or the strangeness should be dealt with in the semantics. THOMASON (1972) followed the latter approach and has proposed

to assign to such sentences no truth values. This idea is worked out in the framework of Montague grammar by WALDO (1979). In his proposal several semantic domains contain partial functions, and the function corresponding with *shiny* is not defined for arguments such as 'the velocity of light'. So (1) is not associated with a truth value.

Waldo's approach gives rise to strange consequences. Formulas which one might expect to be equivalent, are not. I will discuss two examples, and indicate how the problems could be solved by using total functions in the model.

The first example concerns formula (2), where $\phi \in \text{ME}_t$.

$$(2) \quad \phi = \phi.$$

Suppose that the interpretation of ϕ is undefined (e.g. because it is the translation of (1)). Then, due to the interpretation of $=$, also (2) is undefined. However, due to the interpretation of connectives (which uses 'extended interpretations'), formula (3) gets the interpretation *true*: This difference in interpretation is, in my opinion, a strange result.

$$(3) \quad \phi = \phi \wedge \phi = \phi.$$

The second example is based upon a suggestion of R. Scha (pers.comm.). It concerns formula (4), where $z \in \text{VAR}_t$, and where $\phi \in \text{ME}_t$ is as in (2).

$$(4) \quad \lambda z[z=z](\phi).$$

The possible assignments to z are, in Waldo's partial model, the truth values *true* and *false*. Therefore the expression $z = z$ is equivalent with some tautology not containing z , for instance $\forall w[w=w]$. Hence (4) is equivalent to (5)

$$(5) \quad \lambda z[\forall w[w=w]](\phi).$$

According to the standard conditions for λ -conversion, formula (5) can be reduced to (6), which clearly gets the interpretation *true*.

$$(6) \quad \forall w[w=w].$$

Also in (4) λ -conversion is, according to the standard conditions, an allowed operation. Then (2) is obtained, but the interpretation of that formula is undefined. So the formulas (6) and (2), obtained by reduction of (4) are not equivalent, an unacceptable result. We have to conclude that one of the reductions steps is not allowed. This problem is, in my opinion due to the fact that for the variable z in (4), there are two possibilities (true and false), whereas for the arguments ϕ there are three possibilities (true, false, and undefined). Note that the variable z cannot be undefined, because its range consists of all elements in the model of the correct type, and *undefined* is no value in the model.

The above examples show that the laws of logic we have met before, cannot be used in this system without further investigations. In any case the conditions for λ -conversion have to be changed. Unfortunately, Waldo does not provide laws for his system. This causes a difficulty in the study of his proposal. He presents several examples, and each consists of a sentence accompanied by its reduced translation. Since I do not know which reduction rules hold in an approach with partial functions in the semantics, I cannot check the correctness of the examples. Also other authors who describe a fragment with the use of partial functions in the semantic domains, do not present reduction rules (HAUSSER (1976), COOPER (1975)). The last author mentions at least that not all standard reductions remain valid. I expect that it will be very difficult to reformulate the reduction rules. An obvious attempt to improve the conditions for λ -conversion would be to require that the reduction of $\lambda z[\alpha](\beta)$ is allowed only if β is defined. This is, however, not a syntactic condition on β , and I doubt whether it is possible to give a syntactic characterization of 'undefined'.

I already explained that the problem is due to the fact that a variable cannot take the value *undefined*, whereas an argument ϕ (which might be substituted for that variable) can be undefined. Therefore I expect that the problems will be solved when a third truth value is introduced, say a value *error*. In any case, the two problems mentioned above disappear. If the value *error* is assigned to z , then the interpretation of $z = z$ is always the same as the interpretation of $\phi = \phi$, even in case ϕ is undefined. Now λ -conversion is allowed both in (4) and in (5), and furthermore, all formulas (i.e. (2)-(6)) get the same interpretation for all values of ϕ . Note that this plea for using a third value is not an argument for using some of the existing tables for three valued logic. Waldo uses supervaluations (Van FRAASSEN 1969), and one might try to reformulate super-

valuations for an approach with a third truth value.

The idea of using a third truth value is not new; it goes back to ŁUKASIEWICZ (1920), who gives tables for the connectives in a three-valued system. In the theory of topoi one introduces a value representing 'undefined' (GOLDBLATT 1979, p.268). In the theory of semantics of programming languages the problems of working with 'undefined' are well-known. Undefinedness arises, for instance, when a process is defined for calculating the value of a function, whereas the process does not terminate normally because not enough memory capacity is available. The standard approach in this field is not to use partial functions, but to make the functions total by means of the introduction of an extra element in the semantic domain, called 'errorvalue' or 'bottom' (SCOTT & STRACHEY 1971, GOGUEN 1978). In the field of Montague grammars the situation is as follows. A model for intensional logic with 'undefined' as possible value, is presented in Von KUTSCHERA (1975). It is however common practice to consider undefinedness not as a value (see KAMP 1975, COOPER 1975, HAUSSER 1976, WALDO 1979, KLEIN 1981). I know of only one author who presents a treatment of a certain fragment and uses a model with 'undefined' as value: Ter MEULEN (1980).

2.3. Discussion

The examples given in sections 2.1 and 2.2 show that it will be a considerable change of the framework to allow for algebras with partial operations. Moreover, it is not obvious in which way we have to proceed. GRAETZER (1968,p.80) says the following. 'For algebras there is only one reasonable way to define the concepts of subalgebra, homomorphism, and congruence relation. For partial algebras we will define three different types of subalgebra, three types of homomorphism, and two types of congruence relation [...] all these concepts have their merits and drawbacks'. This situation constitutes an argument for my expectation that it will be a considerable task to develop a framework based upon the use of partial algebras. What I have seen of the literature concerning partial algebras did not give me the confidence that an elegant framework can be built using this notion (e.g. ANDREKA & NEMETI (1982), MIKENBERG (1977), ANDREKA, BURMEISTER & NEMETI (1980)). The example concerning partial functions in the semantics gives me the conviction that it is not a good idea to base a semantic theory on partial functions. For these three reasons I do not sympathize with the idea of basing the framework on partial algebras.

As for the introduction of partial rules in the syntax only, the situation seems to be different. It is just a minor change of the framework because the homomorphic relations between the algebras of the framework are hardly disturbed. An argument in favor of the introduction of partial rules is that such rules are used frequently in practice. But there also are arguments against the introduction of partial rules in the syntax. Below I will mention some of them, thereafter they will be discussed.

1. *Consistency of argumentation*

The first argument concerns the consistency of our argumentation. In a Montague grammar we distinguish categories, and the rules give the information in which way the expressions of certain categories combine to form expressions of other categories. An argument for distinguishing such categories (given e.g. in chapter 1) was that certain groups of expressions behave differently from other groups in syntactic or semantic respects. Designing partial rules would mean that among a single category we distinguish two subgroups (these expressions of a category to which the rule can be applied, and those to which the rule cannot be applied). A consistent reaction in such a situation would be to conclude that the system of categories was not refined enough, and that the system has to be refined in such a way that the partial rules are no longer partial.

2. *Filtering power*

A partial rule introduces a kind of filter in the grammar, and filters form a powerful tool which can easily be abused. In a Montague grammar the syntactic rules provide the information which kinds of expressions may be combined to form new expressions. But partial rules would make it possible that the syntactic rules combine rubbish to rubbish, whereas a final partial rule would filter out the undesired expressions. In this way, the other rules would not give information about the combinations which make sense and which not. The filtering power of partial rules in syntax is employed in the first two examples given above.

3. *Generation of expressions*

Often one wishes to conceive a grammar as a generating device. The rules of the fragment presented in chapter 4 can easily be conceived in this way. A rule like S_4 is considered as an instruction stating that if one wants to generate a sentence, one has to generate a term and an

IV-phrase, and next combine them. The rules for term-formation and IV-formation are, in the same way, considered as instructions for a generating process. The processes of generating a term and of generating an IV-phrase may be carried out independently, and every outcome of the processes is acceptable. Details of a computer program based upon these ideas can be found in JANSSEN (1980a). Suppose now that the grammar contains a partial variant of S_4 , say a rule which imposes restrictions on the possible combinations of a term with an IV-phrase. Then the simple algorithm just sketched cannot be used. One has to design an algorithm that gives the guarantee that after a finite amount of time an acceptable combination is found (provided that there is one). This requirement would make the algorithm rather inefficient: the only possibility I see for such an algorithm is one which tries out all possible combinations of a term with an IV-phrase. So in the perspective of a generation process partial rules are unattractive.

4. *Consequences*

An important argument in favor of total rules is that this requirement has attractive consequences. On a more theoretical level it gives rise to an interesting restriction on the possibility to incorporate transformations in a Montague grammar (see section 3). On a more practical level the requirement of using total rules turns out to be a valuable heuristic tool. Several partial rules from the literature can be reformulated or eliminated, and the requirement suggests how this can be done. Thus several proposals from the literature can be replaced by a simpler treatment (see section 4).

5. *No theory*

The introduction of partial rules, even if only in the syntax, constitutes a considerable change of the framework. As the given examples have shown, the theory which we have developed, cannot be used without corrections. Since a theory about partial syntactic algebras is not available, there is no guarantee that all consequences are acceptable.

None of these five arguments is decisive. As for 'consistency', it is indeed more elegant to use the argument for the introduction of categories in all situations with the same conclusion. But with respect to other considerations there might be arguments of elegance in favor of partial rules (e.g. that in that way linguistic generalizations can be captured). That partial rules introduce a powerful filter, is not an impressive theoretical argument since the algebraic grammars have a universal generative capacity

anyhow. As for the argument of 'generation', it is not a primary aim of our grammar to develop an efficient generating device. From a practical point of view, a parser might even be of more importance than a generator. The fact that the practical consequences of using total rules turns out to be attractive in the situations considered, is not a guarantee that in other cases this will be the case as well, and that there is no theory about partial algebraic grammars might be a challenge to develop such a theory.

The arguments against the introduction of partial rules and the arguments in favor of doing so, have to be weighed against each other. The arguments given above show that there are several unattractive aspects related with the introduction of partial rules. I do not know of convincing arguments for the need or attractiveness of partial rules. In the remainder of this chapter I will show that there are several alternatives for the introduction of partial rules. These alternatives are: reformulating as a total rule (section 3), reformulating as a rule operating on another category (section 4) and a refined system of subcategories (section 5). It will turn out that the use of these alternatives gives, in most cases, rise to a simpler treatment than originally proposed: the requirement of using total rules turns out to be a valuable heuristic tool. So the situation can be summarized as follows: there are arguments against the introduction of partial rules, and attractive alternatives are available. Therefore I do not feel enthusiastic about the introduction of partial rules in the syntax. I do not state that I will never use partial rules myself, but I would first try to use total rules.

3. INCORPORATING TRANSFORMATIONS

In the field of transformational grammars, the use of partial rules is standard. As part of their specification the transformations always contain a restriction on the inputs to which they may be applied (a SC: i.e. structural condition). One might wish to incorporate transformations in Montague grammar in order to benefit from the syntactic insights obtained in that field. In this section I will present a general method for the incorporation of a class of transformations in a Montague grammar in which all rules have to be total.

Some characteristics of transformations are as follows

1. Transformations define mappings from trees to trees; these trees represent constituent analyses of sentences.
2. If several transformations can be applied, then their order of application may be prescribed.
3. A transformation is applied to one input tree at a time.
4. A transformation imposes structural conditions determining the possible input trees.

In order to take care of the first point, it is required that a Montague grammar does not produce plain strings, but trees, (or, equivalently, labeled bracketings). Let us assume that Montague's framework is adapted in the way proposed by PARTEE (1973). So the grammar produces trees. This change of the system turns all rules into rules which operate on trees, so in a certain sense all the rules in the grammar become transformations. In order to avoid confusion of terminology, I will use the name C-transformation ('Chomskyan') for transformations used in transformational grammars. Once that they are incorporated in a Montague grammar, they are called M-transformations.

The second characteristic point is not acceptable in our framework. As explained in section 1, explicit rule ordering does not fit into the algebraic approach. But an implicit rule ordering which has the same effects might be possible. The third point does not give rise to problems. Although the rules in a Montague grammar mostly take two arguments, there is no objection against rules taking one argument. The fourth point is problematic since it implies that C-transformations are partial rules. This is an important characteristic of C-transformations which makes them very attractive for practical use. It makes it possible to indicate in a simple way what the relevant input trees are, without the need to bother about irrelevant inputs.

I will incorporate a class of C-transformations in a Montague grammar which requires total rules, by reformulating them in a way which makes them total. The reader might be surprised by this reformulation and at first glance consider it as a sneaky trick employed in order to obey the letter of the principle. This is not completely true. The reformulation expresses a different view on transformations than the standard one, and it has interesting consequences.

The reformulation proceeds as follows. Suppose that a C-transformation

is given in the following form.

If the input sentence satisfies structural condition SC, then we may apply transformation T in order to obtain a new sentence, otherwise T cannot be applied.

Its reformulation as a total rule has the following form.

To the input sentence we apply operation T'. Operation T' is defined as follows. If the input sentence satisfies the structural condition SC, then transformation T is applied, and otherwise the 'do nothing' transformation is applied.

By the 'do-nothing' transformation is understood an operation which does not produce another expression, but which gives its input unchanged as output. The reformulation expresses the view that an M-transformation applies to all expressions of the required category, and that its application yields always a result.

Corresponding with a M-transformation T' there has to be a translation rule τ . For the cases that we did 'nothing' in the syntax, we do 'nothing' in the semantics: the input formula is given unchanged as output. This means that for these cases the translation rule τ can be represented as the polynomial $x_{t,1}$. Since in our framework τ has to be represented by means of a single polynomial expression, τ yields for each input formula, that formula as output. So the M-transformations (obtained with the method described here) do not change meanings. Consequently, if one wants to incorporate C-transformations in this way in a Montague grammar, then these transformations have to be meaning preserving! This requirement is a well-known hypothesis in the so called standard theory of transformational grammars (see PARTEE 1971 for a discussion); it is, however, nowadays not generally accepted.

The conclusion that transformations have to be meaning preserving, holds only for the method described above. But I do not know of any other uniform method for incorporating transformations in a Montague grammar with total rules. To illustrate this, I consider one attempt. Instead of requiring that the translation rule corresponding with a do-nothing transformation is the identity operation on formulas, we might require that it is the identity operation on meanings (but not necessarily the identity on formulas). This would make it possible that the polynomial is not the identity when interpreted for the real transformation. Such a rule τ has the following effect:

$$\tau(\phi) = \begin{cases} \rho(\phi) & \text{if } \phi \text{ is the translation of an expression which satisfies the conditions for application of the transformation } (\rho \text{ formalizes the semantic effect of the transformation}) \\ \phi & \text{otherwise.} \end{cases}$$

The first objection is that this effect cannot be obtained by means of polynomial over IL. In order to obtain the effect of such a choice, IL has to be extended with something like the *if-then-else* construction. There would be, however, no problem in doing so. A more essential objection is that in the description of the translation rule τ information about the (syntactic) expressions is used. This has to be replaced by information concerning their meanings. For most transformations there is probably no semantic property corresponding to the condition on the transformation. In any case, we have no uniform method for obtaining a logical condition which is equivalent with the structural condition of the transformation. So a uniform method for finding the polynomial cannot be given.

I described a uniform method for the incorporation of a class of transformations in Montague grammar by means of a do-nothing transformation. This method might be generalized to a method to eliminate certain partial rules from a Montague grammar. For rules with one argument the method can be used if the rule is meaning preserving. For rules with more than one argument the use of a kind of do-nothing transformations implies that (at least) one of the inputs should have the same category as the output. The do-nothing transformation has to correspond to a translation rule which is the identity on formulas. Therefore the translation rule which corresponds with the original partial rule has to be the identity translation for one of its arguments. So this method can be used only for very limited class of the partial rules with more than one argument.

4. DEFINED FOR ANOTHER CATEGORY

4.1. Introduction

In this section I will consider several rules from the literature which are partial, and for which the corresponding translation rule is not meaning preserving. This implicates that the method developed in the previous section cannot be used for them. The method employed in this section is to

reformulate the rule for another category than where it was originally formulated for. It turns out that in all cases the new version of the rule is simpler than the original formulation of the rule, and sometimes the original rule was incorrect whereas the new rule is correct. This shows the heuristic value of the framework, and of the requirement of using total rules in particular. The examples are presented in the notation of the original proposal; most examples were already mentioned in JANSSEN (1978a).

4.2. He₁ is loved

PARTEE (1973) considers the M-transformation 'Passive Agent Deletion'. An example is

$$F_{102}(he_1 \text{ is loved by him}_3) = he_1 \text{ is loved.}$$

Translation:

If $\phi \in P_t$ and ϕ translates into ϕ' , then $F_{102}(\phi)$ translates into $\exists x_j \phi'$.

On the one hand this transformation applies only to input trees of a special form, on the other hand the translation rule is not the identity mapping. This means that we cannot reformulate this transformation as a total rule, and that Partee's way of dealing with agentless passive is disallowed by the requirement of using total rules. For the example under discussion, the literature provides an alternative. THOMASON (1976) presents rules for generating passive directly, i.e. without a passive transformation and without a passive agent deletion.

4.3. Give John a book

The C-transformation of dative shift changes sentence (7) into (8).

(7) *Mary serves the cake to John*

(8) *Mary serves John the cake.*

A refined category system for sentences in which dative-shift would be a total rule is very difficult to design (since each new subcategory would require rules producing expressions of that subcategory). Also here the

literature contains an alternative. DOWTY (1979a) shows that the partial rule of dative shift on the level of sentences, can be replaced by a rule on the level of lexical elements. That rule changes the category of the verb *serve* from DTV (verbs which take a Dative and a Term) to TTV (verbs which take two Terms). By having a sufficiently refined category system, these lexical rules become total rules. Many examples of transformations which are reformulated on the lexical level can be found in DOWTY 1978, 1979a, and in BARTSCH 1978b, thus they can easily be reformulated as total rules.

4.4. Mary shakes John awake again

In chapter 5, section 5.2, we considered some semantic aspects of the proposals of DOWTY (1976) concerning the treatment of factives. Now I will consider some syntactic aspects (of course, in doing this, the semantic aspects cannot be neglected). Dowty produces the factive sentence *Mary shakes John awake* from the term *Mary* and the IV-phrase *shake John awake*. This IV-phrase in its turn is obtained from the TV-phrase *shake awake*. The first rule Dowty presents for generating this TV-phrase is as follows.

S_{30} : If $\alpha \in P_{IV}$ and $\phi \in P_t$ and ϕ has the form he_n is γ
then $F_{30,n}(\alpha, \phi) \in P_{TV}$ where $F_{30,n}(\alpha, \phi) = \alpha\gamma$.

An example is:

$F_{30,1}(\text{shake}, he_1 \text{ is awake}) = \text{shake awake}$.

The corresponding translation rule reads:

T_{30} : If α translates into α' and ϕ
translates into ϕ' then $F_{30,n}(\alpha, \phi')$ translates into:
 $\lambda P \lambda x [^V P(\lambda x_n [\alpha' (x, \lambda P [^V P(x_n)]) \text{ CAUSE}[\text{BECOME}[\phi']]])]$.

This rule is a partial rule which is not meaning preserving, so we have to find another approach. Can the above result be obtained by means of a total rule? For generating expressions like *shake awake* one only needs an adjective and a TV-phrase. So it lies at hand to try the following rule

S_{601} : If $\alpha \in P_{TV}$ and $\beta \in P_{adj}$ then $F_{601}(\alpha, \beta) \in P_{TV}$ where $F_{601}(\alpha, \beta) = \alpha\beta$.

The corresponding translation rule would be

T_{601} = If α translates into α' and ϕ translates into ϕ' then $F_{601}(\alpha, \beta)$
translates into
 $\lambda P \lambda x [^V P(\lambda y [\alpha' (x, \lambda P [^V P(y)]) \text{ CAUSE}[\text{BECOME}(\beta' (y))]])]$.

Why did Dowty propose a production of *shake awake*, with as intermediate stage the sentence *He₁ is awake*? This has probably to do with the ambiguity of *Mary shakes John awake again*. On the one reading Mary has done it before, on the other John has been awake before. Dowty treats *again* as a sentence modifier and he needs two different sentences in the derivation in order to deal with the ambiguity. He starts his investigations along this line probably for historical reasons: it is the way in which such constructions are treated in generative semantics. But, as in the previous examples, we need not to follow the old pattern. By rule R_{601} we are guided to another approach to this ambiguity. The one reading can be obtained by combining *again* with *Mary shakes John awake*, the other by combining it with *shake awake*. I do not go into details of this approach for the following reason. After considering several phenomena concerning factives, Dowty observes that his first approach is not completely adequate. He discusses extensively several alternatives and escapes. Finally he concludes 'there would be no reason why we should not then take the step of simplifying rules S30-S32 drastically by omitting the intermediate stage in which a sentence is produced'. Next he presents as the rule which he considers as the best one, a rule which is identical with S_{601} . So the framework has led us immediately to the solution which is the simplest and best one. This example suggests that we might derive from the framework the advice 'when designing a syntactic rule, ask for what you need as input and not for more'.

4.5. See himself

In chapter 5, section 2.1, we considered the derived verb phrase rule of PARTEE (1973). This rule makes verb phrases out of sentences. An example is

$$F_{104}(he_1 \text{ sees } him_1 \text{ self}) = \text{see } him^* \text{ self}.$$

The syntactic part of this rule reads as follows:

If $\phi \in P_t$ and ϕ has the form ${}_t[{}_T[he_i]_{IV}[\alpha]]$, then $F_{104}(\phi) \in P_{IV}$, where $F_{104}(\phi) = \alpha'$, and α' comes from α by replacing each occurrence of he_i , him_i , $him_i \text{ self}$ by he^* , him^* , $him^* \text{ self}$ respectively.

At the one hand the derived verb phrase rule is a partial rule, at the other hand its output belongs to a different category than its input. Therefore we cannot reformulate this rule as a total one using a do-nothing transformation. The derived verb phrase rule is disallowed by the requirement

of using total rules, and we have to find another treatment for the cases where Partee uses this rule. Let us, in accordance with the advice given in section 4.4, just ask for what we actually need and not for more. In the above example we only need a TV-phrase. So we might try the following rule.

S_{602} If $\alpha \in P_{TV}$ then $F_{602}(\alpha) \in P_{IV}$ where $F_{602}(\alpha) = \alpha \textit{ him}^* \textit{ self}$.

The corresponding translation rule reads:

T_{602} If α translates into α' , then $F_{602}(\alpha)$ translates into $\lambda x[\alpha'(x, \wedge \lambda P[\vee P(x)])]$

Would this rule be an acceptable alternative?

Let us consider why one would like to generate *see himself* from the source sentence *he sees himself*. There are semantic arguments for doing so. The sentence *John sees himself* is obviously semantically related to the sentences *John sees John* and *He₁ sees him₁*. In transformational grammar this might be an argument for producing these sentences from the same source: no other formal tool is available. The effect of Partee's rules is that such a transformation is split up into several stages; it amounts to the same relations. Montague grammar has a semantic component in which semantic relations can be laid formally. So if we do not have to ask for a sentence as source for syntactic reasons, we are not forced to do so on semantical grounds. So this cannot be an argument against S_{602} .

PARTEE (1975) provides as an explicit argument for the derived verb phrase rule the treatment of sentence (9)

(9) *John tries to see himself.*

This sentence is generated, using the derived verb phrase rule, from sentence (10)

(10) *he₃ tries to see him₃self.*

The translation of (9) becomes in this case (11)

(11) *try to*($\wedge \textit{john}, \wedge \lambda x_3[\textit{see}(x_3, \wedge \lambda P[\vee P(x_3)])]$)].

Sentence (9) can also be generated according to the rules of PTQ. If we do not change the syntactic details of the rule the following sentence is produced:

(12) *John tries to see him.*

In (12) *him* is coreferential with *John*. The translation is

(13) *try to*($\wedge John, \wedge see(\lambda P[\vee P(\wedge John)])$).

Partee provides arguments for her opinion that interpretation (11) might be preferable to (13). Let us assume that her arguments hold and consider whether S_{602} is compatible with that. The combination of *try to* with the translation of *see himself* (obtained by T_{602}) yields

(14) *try to*($\wedge \lambda x[\wedge see(x, \wedge \lambda P[\vee P(x)])$).

So the translation of *John tries to see himself* is, as desired, equivalent to (11). As Partee notices, the derived verb phrase rule does not prohibit the unwanted reading (13). Rule S_{602} is an improvement since it only allows for reading (11). Of course, S_{602} does not give a complete treatment of reflexives, and I am not sure whether I would like to treat them in this way. For the purpose of the discussion this aspect is irrelevant: I just would like to demonstrate that the requirement of using total rules, and in particular the advice 'ask for what you need', guides us to a better rule than originally proposed.

4.6. Easy to see

PARTEE (1975) presents another example for the derived verb phrase rule:

$F_{104}(he_7 \text{ is easy to please}) = be \text{ easy to please}.$

This example may seem somewhat strange since it produces the IV-phrase *be easy to please* from a sentence containing this IV-phrase. The reason is that the sentence is obtained by some transformation from the source

(15) *To please him₇ is easy.*

This transformation is not sufficient for producing all sentences containing the phrase *be easy to please*. Phrases resulting from F_{104} have to be produced as such, in order to generate (16) and (17).

(16) *few rules are both explicit and easy to read*

(17) *try to be easy to please.*

In PARTEE (1977a) other constructions are considered which contain expressions of this kind, such as

(18) *John is being hard to please.*

In order to deal with such expressions Partee needs another rule, called

the derived adjective rule, which has the following effect

$${}_{IV}[be\ easy\ to\ please] \rightarrow {}_{ADJ}[easy\ to\ please].$$

This is again a partial rule which cannot be brought in accordance with the restriction of total rules. So for (15)-(18) an alternative has to be given.

The advice given in section 3.4 stimulates us to ask just for what we need for generating *easy to please*. We need an expression like *easy* and some TV-phrase. Let us, following PARTEE 1977a, assume that we have a special category \overline{ADJ} which contains *easy*, *tough* etc. The resulting expression *easy to please* will be of the category ADJ' . Then we are guided to the following rule:

$$S_{603}: \text{ If } \alpha \in P_{\overline{ADJ}} \text{ and } \beta \in P_{TV} \text{ then } F_{603}(\alpha, \beta) \in P_{ADJ'}: \text{ where}$$

$$F_{603}(\alpha, \beta) = \alpha \text{ to } \beta.$$

The translation of (this) *easy* must be such that it may be combined with an TV-translation in order to obtain an expression of the type of translations of adjectives. Then the translation rule reads

$$T_{603}: \text{ If } \alpha \text{ translates as } \alpha' \text{ and } \beta \text{ as } \beta' \text{ then } F_{603}(\alpha, \beta) \text{ translates into}$$

$$\lambda x \alpha' (\lambda y \beta' (y, \lambda P [{}^V P(x)])).$$

Rule S_{603} makes it possible to generate the expressions containing *easy to please* we mentioned above. Unfortunately, Partee does not provide an explicit semantics for the source of all her constructions (sentence (15)) so we cannot compare it with the semantic consequences of S_{603} ; but I expect that she will finally end up with something close to the result of T_{603} . Concerning the syntax, it is demonstrated that our requirement guides us to a much simpler treatment.

In section 3.5 and 3.6 we have considered two examples concerning the derived verb phrase rule. These examples do not cover all possible applications of the rule. But the treatment given here shows that in any case the two kinds of examples for which Partee has used the derived verb phrases rule can be dealt with in a better way by means of total rules.

5. SUBCATEGORIZATION AND RULE SCHEMES

5.1. Hyperrules

An argument for distinguishing categories (given for instance in

chapter 1, section 1.3) is that certain groups of expressions behave (syntactically or semantically) differently than other groups of expressions. If for some rule it turns out that the rule can only be applied to a subset of the expressions of its input category, then this can be considered as an indication that the system of categories is too coarse. A method to avoid partial rules consists of refining the system of categories. In this section we will consider examples of this method, and present tools which are useful when it is employed.

There are several arguments for distinguishing among the category of nouns the groups of mass nouns and of count nouns. One of the differences between the expressions of these two groups is their behaviour with respect to determiners. Let us compare, as an example, the count noun *ring* with the mass noun *gold*. Well-formed are *a ring* and *every ring*, whereas ill-formed are *a gold*, and *every gold*. In larger expressions the same differences arise: well-formed are *a beautiful ring* and *every ring from China*, whereas ill-formed are *a beautiful gold* and *every gold from China*. These differences constitute an argument for introducing in the grammar the separate categories 'Mass Noun' and 'Count Noun'.

In many respects, however, mass nouns and count nouns behave analogously. Expressions of both categories can be combined with relative clauses and with adjectives. If we treat mass nouns and count nouns as being two independent categories, then the consequence is that the rules for relative clause formation and for adjective addition are duplicated. Thus the grammar will contain a lot of closely related rules. This effect will be multiplied if more categories are distinguished among the nouns. Therefore it is useful to have a tool for controlling this proliferation. Such a tool are rule schemes.

Rule schemes are not new in Montague grammars; recall the rule for relative clause formation given in chapter 4.

$S_{3,n}$: $CN \times S \rightarrow CN$

$F_{3,n}$: replace in α all occurrences of he_n by *him/she/it*, and of him_n by *him/her/it*, according to the gender of the first noun or term in β ; concatenate (α , *such that*, β).

This cannot be considered as a rule because $F_{3,n}$ deals with occurrences of he_n , whereas this expression does not occur in the lexicon of the fragment: examples of relevant expressions of the fragment for this rule are he_1 , he_2 . So we have to consider $S_{3,n}$ as a rule scheme out of which rules can be obtained.

This can be done by replacing all occurrences of n in the scheme by some number. Thus $S_{3,n}$ stands for an infinite collection of actual rules.

In $S_{3,n}$ three characteristic features are illustrated of the kind of rule schemes that I will use. The first one is that a rule scheme differs from a real rule by the occurrence of a parameter. $S_{3,n}$ contains the parameter n , which stands for a number. Schemes may contain several occurrences of one or more parameters, and I will put no restrictions on where a parameter stands for. The second characteristic feature is that out of a scheme an actual rule can be formed by means of substituting the same expression for all occurrences of a parameter. If it is not required that all occurrences are replaced by the same expression then the occurrences of the parameter will be indexed (e.g. with n_1, n_2, \dots), and then occurrences with different indices may be replaced by different expressions. The third feature is that a parameter may stand for a part of a (formally simple) symbol. The expression he_n is, formally spoken, a single generator in the syntactic algebra, but in the scheme given above it is treated as a compound symbol with he and n as parts. This does not change the role of he_1 in the algebra; it remains a simple generator. One should distinguish the formal position in the algebra, and the presentation of an infinite collection operators (or generators) by means of schemes.

A rule scheme involving nouns is the following.

$S_{604,n}$: Adj \times *cm* Noun \rightarrow *cm* Noun

$F_{604,n}$: Concatenate (α, β) .

From this scheme two actual rules can be obtained. If *cm* is replaced by 'Count', then we obtain a rule which says that an adjective in combination with a Count Noun forms a new Count Noun. If *cm* is replaced by 'Mass', then we obtain a rule which says that an adjective in combination with a Mass Noun forms a new Mass Noun. This scheme exhibits again the feature that a compound symbol in the sense of the scheme, can be a single symbol in the algebraic sense. In the algebra 'Count Noun' is a category symbol, whereas in $S_{604,n}$ it is a compound with 'Count' and 'Noun' as individual parts. Notice that the above scheme contains two parameters: n and *cm*.

The new formal aspects introduced in this section are the use of compound category symbols and the possibility to use parameters for parts of these symbols. The practical impact of this is that partial rules can be avoided by increasing the number of categories, and that rule schemes can be used for handling these categories.

Now I will introduce some terminology. The parameters in the rule schemes are called *metavariables*. To distinguish the rule schemes of the kind just described, from others, the former are called *hyperrules* (i.e. they are rules containing metavariables). Hyperrules without the occurrence of a variable are considered as a special case; by means of an 'empty' substitution they become actual rules. I will give the hyperrules a 'name' which starts with an H, and its parameters will not be included in the name. So rule $S_{604,n}$ mentioned above will be called H_{604} . The distinction between Count Nouns and Mass Nouns is in linguistics called subcategorization. I will use this term with the following formal interpretation. A category C_1 is called a subcategory of a category C_2 if the carrier of sort C_1 is a subset of the carrier of sort C_2 .

5.2. Metarules

Suppose that we have a hyperrule which contains some metavariable. In the example from section 4.1 concerning nouns, I explicitly listed the two possible substitutions. But often the situation will be more complex. There are arguments for distinguishing among the nouns many more subcategories, and we will meet examples where infinitely many substitutions are possible. Therefore it is useful to have a handsome tool for telling what the possible substitutions for a metavariable are. In the sequel we will use rewriting rules for this purpose. Besides the grammar consisting of hyperrules I will give a second grammar, called *metagrammar*. This grammar consists of a collection context-sensitive rewriting rules, and in these rules the metavariables of the grammar occur as auxiliary symbols. If we take some metavariable as start symbol, then the metagrammar determines a language: the set consisting of all strings which can be produced from the metavariable which was taken as start symbol. The possible substitutions for a metavariable in some hyperrule are all strings from the language generated by the metagrammar using that metavariable as starting symbol.

The benefit of using a metagrammar becomes especially clear in cases where there are several levels of subcategorization and crosslinks in the category system. As example I present the metagrammar for the subcategorization system given in CHOMSKY (1965, p.85); it is striking to observe that Chomsky used rewriting rules as well for the presentation of the subcategorization.

$$\begin{array}{l}
 \textit{common} \rightarrow \textit{sgn count} \\
 \textit{sgn} \rightarrow \left\{ \begin{array}{l} + \\ - \end{array} \right. \\
 - \textit{count} \rightarrow \textit{sgn Abstract CN} \\
 \textit{count} \rightarrow \left\{ \begin{array}{l} - \textit{Animate CN} \\ \textit{anim} \end{array} \right. \\
 \textit{anim} \rightarrow \textit{sgn Human CN}
 \end{array}$$

According to the convention for substitution, this metagrammar implicates that a hyperrule containing *anim* as metavariable represents two actual rules (for the subcategories + Human CN and -Human CN), and that a hyper-rule containing *common* represents 5 actual rules.

A grammar designed in the way sketched above is a system with two levels in the grammar: the level of metarules and the level of the (hyper)rules. The conception of a grammar with two levels is due to Van Wijngaarden, and was developed for the formal description of the syntax of the programming language ALGOL 68 (see VAN WIJNGAARDEN 1975). He used these notions hyper-rule and metarule with about the same meaning (for a linguistically oriented example see VAN WIJNGAARDEN 1970). The same terminology, although with a somewhat different meaning, is used in GAZDAR & SAG 1981 and GAZDAR 1982. The concept of a two-levelled grammar gives rise to an elegant method handling a lot of rules, even an infinite number. The method could easily be generalized to multi-level grammars. In Van Wijngaarden's original system the metarules have to be context-free, whereas I allowed for context sensitive rules. This liberty has no consequences since the generative power of system lies in the rules, and not in the metarules. In the example given above (Chomsky's subcategorization) the context sensitive rules turned out to be useful. If we would be more liberal, and allow to use a type-0 grammar as metagrammar instead of a context sensitive grammar, then this would have the consequence that the by the metagrammar produced language would be undecidable. Then it would not be decidable whether a substitution for a metavariable is allowed, and consequently the set of actual rules would not be recursive. Therefore type-0 grammars are in our framework not acceptable in the metagrammar.

5.3. Variables

The use of variables in a Montague grammar gives rise to certain problems. I will consider here two of them. A more extensive discussion will be

given in chapter 8.

1. 'Left over'

According to the PTQ rules we may generate the sentence He_3 runs. This is not a correct English sentence because it contains he_3 , which is not a correct English word.

2. 'Not there'

One might apply a rule which involves variables in a situation in which such variables are not there. In this way one obtains relative clauses, which do not contain a reflexive pronoun. An example is *the man such that Mary seeks a unicorn*.

In order to eliminate these two problems, in chapter 8 a restriction will be proposed that contains the following two conditions:

- (I) The production of a sentence is only considered as completed if each syntactic variable has been removed by some syntactic rule.
- (II) If a syntactic rule is used which contains instructions which have the effect of removing all occurrences of a certain variable from one of its arguments, then there indeed have to be such occurrences.

It is evident that requirement (II) can be guaranteed by means of a partial rule. To this aspect I will return later. Requirement (I) says that all stages of the derivation process have to meet a certain condition. So it appears to be a global filter. Since one can tell from the final result whether the condition is met, it reduces however to a final filter. As I explained in chapter V, filters are not acceptable in our framework. But the effect of (I) can be obtained by means of a partial rule as follows. Replace everywhere in the grammar the category of Sentences by the category of Protosentence (so the grammar produces Protosentences). Then we add an extra rule which produces a sentence out of a protosentence in case requirement (II) is fulfilled, and which is not applicable when this requirement is not fulfilled. Thus only sentences obeying (I) are produced. Since I aim at avoiding partial rules, I have to provide an alternative method for the incorporation of the above two restrictions. This will be given below.

Categories are defined to be complex symbols consisting of two parts: a category name as we used before (e.g. S), and a representation of a set of integers. The set indicates which indices occur in the expressions of

that (complex) category. So he_2 or he_3 is an expressions of the category $(T, \{2, 3\})$. Other examples are He_4 runs of the category $(S, \{4\})$, and $John$ of the category (T, \emptyset) . The language generated by the grammar is defined as the set of expressions of the category (S, \emptyset) .

The hyperrules of the grammar contain variables for integers (n) and variables for sets (set_1, set_2, \dots). The following notations are used.

$set_1 \cup set_2$ denotes the set obtained as union of the sets set_1 and set_2
 set with n is a compound expression indicating that set contains element n
 $set - n$ is a compound expression denoting the set obtained by removing the element n from set .

The hyperrule corresponding with S_4 reads

$$H_4: (T, set_1) \times (IV, set_2) \rightarrow (S, set_1 \cup set_2)$$

F_4 : replace the first verb in β by its third person present singular, concatenate (α, β) .

This hyperrule states that set of the syntactic variables in the sentence is the union of the syntactic variables in the T-phrase and the IV-phrase. An example of an actual rule obtained from H_4 is

$$H_4: (T, \{1, 2\}) \times (IV, \emptyset) \rightarrow (S, \{1, 2\})$$

F_4 : see above.

This rule may be used in the production of He_1 or he_2 runs. Corresponding with S_2, S_5, \dots, S_{13} and S_{17} we have analogous hyperrules. The hyperrules corresponding with the rules S_{14} and S_3 are:

$$H_{14}: (T, set_1) \times (S, set_2 \text{ with } n) \rightarrow (S, set_1 \cup [set_2 - n])$$

F_{14} : substitute $(\alpha, \text{first occurrence of } he_n \text{ in } \beta)$;
 replace all occurrences of he_n in β by $he/she/it$ and of him_n by $him/her/it$ according to the gender of the first noun or term in α .

$$H_3: (CN, set_1) \times (S, set_2 \text{ with } n) \rightarrow (CN, set_1 \cup [set_2 - n]).$$

F_3 : Replace he_n in β by $he/she/it$ and him_n by $him/her/it$, according to the gender of the first CN in α ;
 concatenate $(\alpha, \text{such that, } \beta)$.

An actual rule obtained from H_{14} is

$$H_3: (T, \emptyset) \times (S, \{2, 3\}) \rightarrow (S, \{3\}).$$

An application of this rule is the production of $John$ loves him_3 from $John$

and he_2 loves him_3 .

A formalist might object to the hyperrules given above since they implicitly assume that the reader knows what sets are, and what is meant by the symbols \cup , *with* and $-$. This is, however, not knowledge about operations of the grammatical system, but set theoretical knowledge, and the rules should not be dependent on this knowledge. In appendix 3 of this book it will be shown how these notions can be described by means of purely grammatical tools (*viz.* by rewriting rules).

Clause (I) required that the expressions of the generated language do not contain any occurrences of syntactic variables. In my approach this requirement is not formalized as a filter or as a condition in a partial rule, but within the system of categories. This is theoretically more attractive, and practically somewhat simpler. Clause (II) requires that in case a rule is applied which removes variables, then there are such occurrences. This clause is also dealt with in the categorial system, as one can see from the following. Let us suppose that the categorial information given in the rules corresponds with the syntactic operations performed by these rules (*i.e.* if the rule removes all occurrences of a variable, its index is removed from the set mentioned in the category of the produced expression). This assumption can easily be checked from the rules. Assuming this correspondence, the condition *set₂ with n* in H_{14} and H_3 guarantee that these rules are applied only to expressions containing the required occurrences of variables. So instead of formalizing (I) as a condition in a partial rule, it is formalized within the categorial system. This is theoretically more attractive, but practically somewhat more complex.

One observes that the requirements concerning variables can be dealt with in accordance with the aim of using total rules. This is made manageable by using a two-level grammar. Within this system the requirements can be handled about as easy as in a system with partial rules. But the introduction of two levels did not make the system simpler. Therefore I would not say that the requirement of using total rules has led us here to a simpler treatment. In order to see practical advantages of using a two-level grammar, one has to consider a much more complicated situation. Such a situation will be described in chapter 9: the interaction of tense scope, and quantifier scope. But in the present situation the advantage is only of theoretical importancy. Therefore one might take in practice the following position. It has been shown that the requirements concerning variables can be incorporated within a system with only total rules. This implicates

that in practical cases there is no need to treat the requirements explicitly in this way. One might use requirements (I) and (II) as they are formulated, assuming the present formalization.

5.4. A theoretical result

The method introduced in this section for eliminating partial rules consists in refining the system of categories. For nouns I gave an example with five subcategories, and for the treatment of variables even an infinite number. One might consider the possibility of applying this method up to the very limit (every expression constituting a single category on its own). By proceeding that far, all partial rules are eliminated from the grammar. This simple idea is followed in the proof of the following theorem (LANDSBERGEN 1981).

5.1. THEOREM. *For every enumerable algebraic grammar G with partial rules, there is a general algebraic grammar G' with total rules, such that $L(G) = L(G')$.*

PROOF. We obtain G' as follows. For each category C of G and each expression w of this category, we define a new category in G' , denoted by the compound symbol (C,w) . The only expression of this category is w . Since for each sort of G , the expressions are recursively enumerable, the sorts of G' are recursively enumerable as well (but in general not recursive). For each rule R in G there is a collection of rules in G' . If according to a rule of G the expression w_0 (of category C_0) is formed out of the expressions w_1, w_2, \dots, w_n of the categories C_1, \dots, C_n , then there is in G' a rule producing expressions of the category (C_0, w_0) out of expressions of the categories $(C_1, w_1), \dots, (C_n, w_n)$. Of course, this rule can be used in only one production, but it is a total rule. Since the rules of G and the expressions of $L(G)$ are recursively enumerable, the rules of G' are recursively enumerable as well. Suppose that the distinguished category of G is S (so $L(G) = G_S$). Then we add for each category (S,w) , where w is arbitrary, a new rule which takes as input an expression of category (S,w) and yields as output the expression w of category S . From this construction it is evident that $L(G) = L(G')$.

5.1. END

The theorem states that every language generated by a grammar with

partial rules can be generated by a grammar with total rules. As such the theorem is not surprising: even finite grammars have a universal generating capacity. The merit of the theorem lays in the method used in its proof. The grammars G and G' do not only generate the same language, but they do so in the same way. The derivational history of a given expression has in G and in G' the same structure. Several properties of G are carried over to G' ; for instance, if G consists of concatenation rules only (i.e. if the rules correspond with a context free rules), then the same holds for G' . This correspondence between G and G' means that the proof can be used for restricted classes of grammars as well.

One might be tempted to conclude from the theorem that grammars with partial rules are just notational variants of grammars with total rules, and that it constitutes a justification for writing partial rules in a framework that requires total rules. This is however not the case, since an important property of G can be lost by transforming it to G' . If G is a recursive grammar, where its generated language $L(G)$ is not recursive, then G' is not a recursive grammar. In chapter 2 we have restricted our attention to the class of recursive grammars. Hence the method used in the theorem may bring us outside the class of grammars we are working with. For this class the grammars with partial rules cannot be considered as a notational variant of the grammars with total rules. So the requirement to use total rules is a substantial one. It has a consequence that not every condition on applicability is acceptable: only those are acceptable which can be reformulated as total rules in a recursive algebraic grammar. In previous sections it has been demonstrated that such a reformulation gives rise to a simpler, a better grammar.

6. THE WELL-FORMEDNESS CONSTRAINT

In this section I will discuss some aspects of a principle for syntax due to Partee. It is called 'the well-formedness constraint', and it reads as follows (PARTEE 1979b, p.276):

Each syntactic rule operates on well-formed expressions of specified categories to produce a well-formed expression of a specified category.

The motivation for this principle is related with the aim 'to pursue the linguists goal of defining as narrowly as possible the class of possible grammars of natural languages' (op. cit. p.276). Although this is a completely different aim than the theme of the present chapter, it turns out

that the well-formedness constraint has practical consequences which can be compared with consequences of our algebraic approach, in particular with the requirement of using total rules. I will restrict the discussion of the well-formedness constraint to these aspects.

Our investigations started from algebraic considerations, and the requirement of using total rules constitutes a formal restriction. The value of the requirement was its impact on heuristics. What is the position of the well-formedness constraint in this respect? Is it a formal restriction, heuristic guideline, or something in between these two? I will first try to answer this question by considering the constraint as it is formulated; Partee's interpretation will be considered thereafter. In order to answer the question concerning the formal position of the well-formedness constraint, it is important to have a formal interpretation for the phrase 'well-formed expression'. I will consider two options.

One might decide to associate the phrase 'well-formed expression' with the meaning that this phrase has in formal language theory. The rules of a grammar produce strings over some alphabet, and these strings are called the well-formed expressions over this alphabet. The epithet 'well-formed' is used to distinguish these strings from the other strings over this alphabet. Un-well-formed generated expressions do not exist by definition. It is possible to tell what the well-formed formulas of predicate logic are, but it is not possible to give examples of un-well-formed formulas of predicate logic: if a string is not well-formed, it is no formula at all. If we apply this interpretation to the PTQ grammar, then we have to conclude that *love him₁* is a well-formed expression (of the category IV) because it is a string produced by the grammar, whereas *love her* is not well-formed (because it is not produced as IV-Phrase). With this interpretation the phrase in the constraint stating that the rules produce well-formed expressions is a pleonasm. The same holds for the input: the only possible expressions of specified categories are the expressions generated by the grammar. With this interpretation the well-formedness constraint just describes how the framework operates, and it is no constraint at all.

One might relate the notion well-formedness with the language generated by the grammar. Then the generated language consists of well-formed expressions, and also all substrings of well-formed expressions are considered as well-formed. Following this interpretation, the constraint says that all intermediate stages in the production process have to be substrings of the produced language. So an acceptable grammar for English has not only to be

adequate (i.e. produce the correct language), but also all the intermediate stages arising from the grammar have to be adequate in a certain sense. This mixture of the notions 'possible grammar' and 'adequate grammar' makes the constraint an unusable one. Suppose that a list of rules of a grammar for English is presented, and one is asked whether they conform the constraint. In order to answer this question one may start to produce some strings by means of the rules, and ask for each application of a rule, whether it is applied to well-formed expressions of English. Suppose that this is the case, then one cannot thereby conclude that all rules from the list obey the constraint, since not all possible derivations have been considered. One has to try and try again, but the definite answer 'yes' cannot be given. It may be undecidable whether an arbitrary grammar satisfies the constraint or not. Of course, this is not a mathematical proof. Such a proof cannot be provided, since the set of English sentences is not a mathematically defined set, but related questions in formal language theory are known to be recursively undecidable. Since the constraint is an undecidable constraint, it cannot be accepted as a restriction on the class of possible grammars (otherwise a more attractive, undecidable, constraint would be 'is an adequate grammar for English').

Partee gives no formal definition of the notion 'well-formed expression'. Conclusions about her interpretation have to be based upon the examples she gives concerning the constraint. As an illustration of the constraint she presents a rule which forms adnominal adjectives from relative clauses. (PARTEE 1979b,p.277). Its syntactic function F_1 has the effect that:

$$F_1(\textit{immigrant who is recent}) = \textit{recent immigrant}.$$

The input for this operation is an ill-formed expression (*immigrant who is recent*), and therefore she judges that this rule is prohibited by the well-formedness constraint. From this example is clear that she does not follow the first interpretation given above, the second one is closer to her intentions. But she would not consider all substrings of well-formed expressions as being well-formed as well (Partee, personal communication). I expect that *John and Peter* is well-formed, whereas *John and* is not. Probably the judgement what well-formed expressions are, is to be based upon linguistic intuitions. In any case, Partee does not give a formal interpretation for the notion 'well-formed expression'. If this notion is not formally interpreted, then the constraint itself cannot be a formal

restriction either. Furthermore, both our attempts to give a formal interpretation were not successful.

I conclude that the constraint has to be considered as a guideline for designing rules. As such it might be useful for its heuristic value, but it has not the position of a formal constraint on the possible kinds for grammars. As a guideline it is a very appealing one, since it aims at a natural way of production; in which no artificial expressions occur as intermediate forms. However, following this interpretation is not without problems. As HAUSSER (1978) remarks, the intuitions concerning the well-formedness of incomplete expressions are rather weak. Hausser gives as example *and about the seven dwarfs quickly*; well-formed or not? Furthermore, the well-formedness constraint does, even in clear cases, not guarantee that only natural production processes are obtained. Hausser gives as example an operation with the following effect.

$$F_n(\text{John kissed Mary}) = \text{Bill walks.}$$

This operator F_n is according to the well-formedness constraint an acceptable operator: an intuitively well-formed expression is transformed into well-formed expression. In order to give real content to the principle, restrictions have to be put on the possible effects of a rule. PARTEE (1979a) gives some proposals for such constraints, and her ideas will be followed in chapter 8.

A consequence of Partee's interpretation of the well-formedness constraint brings us back to the discussion of this chapter. Her interpretation says that in all stages of the production process only well-formed expressions are formed. So there is no need to filter out some of them. Neither there is a need to have obligatory rules which have in transformational grammars the task to transform ill-formed expressions into well-formed ones. So in a grammar satisfying the constraint obligatory rules and filters are not needed. Partee even goes further and interpretes the constraint in such a way that they are disallowed. As we found in section 1, such requirements are a direct consequence of the algebraic framework.

Partee's proposal deviates in an important aspect from our framework. Following linguistic practice, she allows for partial rules. As explained in the previous sections, I would not like to follow this idea and I would prefer to use total rules. Some effects of the well-formedness constraint can be dealt with by means of the requirement of using total rules, as will be shown below.

Suppose that in a grammar with total rules there is a rule S_i of which the syntactic operation F_i has the following effect.

$$F_i(\textit{immigrant who is recent}) = \textit{recent immigrant}.$$

So the rule operates on a common noun phrase which, according to rule $S_{3,n}$ must be constructed from the common noun *immigrant* and a sentence of the form he_n *is recent*. This sentence has to come from the IV-phrase *be recent*. Since we require that the rules are total, we may also combine this IV-phrase with other term-phrases. So the sentence *John is recent* also is generated by the grammar, and this is not a correct sentence of English. This example suggests that an adequate grammar for English with total rules cannot contain a rule which generates *recent immigrant* in the way S_i does, because one cannot get rid of the phrase *be recent*. But an easy solution for the production of *recent immigrant* is available. Follow the advice given in section 4.3, and ask for what we need to produce this phrase. This advice suggests us to ask for an adjective (*recent*) and a noun phrase (*immigrant*). So the requirement of using total rules has the same practical impact here as the well-formedness constraint: it is a guideline for obtaining a non-artificial production process. (Note that I did not prove that it is impossible to have a rule like F_i in a system with total rules; I expect that a refined subcategorization might make this possible).

PARTEE (1979b) discusses certain aspects of the formation of (13).

(13) *Fewer of the women came to the party than of the men.*

Following BRESNAN (1973), this sentence is derived from the (ill-formed) sentence (14) by means of an operation called Comparative Ellipsis.

(14) *Fewer of the women came to the party than of the men came to the party.*

This is in its turn derived from (15) by Comparative Deletion.

(15) *Fewer of the women came to the party than x many of the men came to the party.*

As Partee says, the production of (13) is a difficult case for the well-formedness constraint since it uses the ill-formed source (14). Partee says: 'Unless further analysis of these constructions leads to a different kind of solution, they would seem to require the admissibility of ungrammatical intermediate stages. (Note that the derivations in question give semantically reasonable sources, so any reanalysis has a strong semantic as well as syntactic challenge to meet).' (PARTEE 1979b, p.303,304).

For our requirement of using total rules this production process is problematic as well. It is no problem that the rules of comparative deletion and comparative elipsis are partial rules, since they are meaning preserving. But the production of the ill-formed sentence (14) is problematic since we cannot get rid of this sentence: we cannot filter this sentence out, we may not have it as an expression of the generated language, and we may not use its embeddings (cf. the discussion concerning *recent immigrant*). But why follow this approach? Maybe one judges that a source like (14) or (15) expresses the semantic content of the comparatives more completely than comparatives. Or one wishes to explain the semantic relations between all variants of comparatives by generating them from the same kind of source. In transformational grammar this might be valid arguments, no other formal tools than transformations are available. In a Montague grammar there is a semantic component in which such semantic relations can be formally expressed. So if we do not need such a source for syntactic reasons we may try another approach. The requirement of using total rules guides us toward asking what we need. In order to make a sentence of which the kernel consists of two terms and a verb phrase, we need two terms and a verb phrase. Therefore we should introduce a three place rule

F₆₀₅ (*John, Bill, see women*) = *John sees more women than Bill.*

The semantic component has to express what is compared; the syntax needs no to do so.

Another rule might compare of two nouns in which degree they are involved in a certain property.

F₆₀₆ (*man, boy, come to the party*) = *fewer of the men come to the party than of the boys.*

One may also compare two terms for two verb phrases

F₆₀₇ (*John, Bill, see men, meet women*) = *John sees more men than Bill meets women.*


These examples do not provide for a treatment of the comparative. They just illustrate the kind of solution one might search for in a framework with total rules. Variants are possible: for instance, one might introduce compound quantifier phrases like *fewer of the man than of the boys*, and use instead of F₆₀₆ a rule with two arguments. Note that all these attempts to find total rules, are in accordance with the well-formedness constraint.

CHAPTER VIII

CONSTITUENT STRUCTURES

ABSTRACT

Some proposals from the literature for assigning constituent structures to the expressions produced by a Montague grammar are shown to violate the framework. A treatment of the syntax of the PTQ fragment is presented which assigns constituent structures to the produced expressions and which meets the requirements of the framework. Furthermore a restricted set of syntactic operations is introduced for the description of the syntactic rules.



1. STRUCTURE - WHY?

The syntactic rules of PTQ make a primitive impression in comparison to the kind of rules used in transformational grammars. A first point of difference is that the syntactic operations make no reference to the constituent structure of the involved expressions. A second one is that the syntactic operations are described without any formalism: the desired effects are described by English sentences. On the one hand English is a rather poor tool since in this way the description of the syntactic operation can hardly use any abstract syntactic information. At the other hand it is a very unrestricted tool, since it allows any operation that can be described in the English language. Since the earliest times of Montague grammar, it has been tried to bring the syntax of Montague grammar closer to that of transformational grammar. This would open the possibility to incorporate syntactic knowledge from transformational grammars in Montague grammar, and to discuss the differences. In this chapter I will present the first steps of an approach which makes the syntax of Montague grammar less primitive: by developing a formalism for the formulation of the syntactic rules, and by introducing constituent structures in the syntax.

An example of the kind of structure used in transformational grammars is given in figure 1. The tree is not taken from any proposal in that field (then several details would be different), but it can be used to illustrate what kind of information is provided by such trees. The words attached to the end nodes of the tree yield, when read in the given order, the sentence of which the tree represents the constituent analysis. Constituents are groups of words which have a certain coherence. This appears for instance from the fact that it is rather easy to replace a constituent of a sentence by another group of words, whereas this is not the case for arbitrary groups of words from the sentence. The tree in figure 1 indicates what the constituents of the sentence are: all words of a certain constituent are connected to the same node in the tree. This node is labelled by a symbol: the name of the category of the constituent. Thus the tree gives the information that each word is a constituent, and that e.g. *a unicorn* is a constituent, whereas *seeks a* is not.




Figure 1 Tree like those in transformational grammar

A first argument for the introduction of constituent structures in the syntax of Montague grammars is that it would make it possible to incorporate ideas, or even particular rules, from transformational grammars into Montague grammar. I will not try to sketch the role such structures play in the syntax of transformational grammars; the reader should accept that constituent structures have proven their usefulness. A second argument is that, even without the aim of incorporating ideas from transformational grammar, it is useful to have structural information available about the expressions dealt with. An example, based upon a phenomenon from the PTQ grammar, is the following (PARTEE 1973).

Rule S_{11a} from PTQ, the rule for verb-phrase conjunction, produces the IV-phrase

- (1) *walk and talk*.

Rule S_8 produces from (1) and the verb *try to* the IV-phrase

- (2) *try to walk and talk*.

From the term *John* and the IV-phrase (2) we can produce, according to rule S_4 , the sentence

- (3) *John tries to walk and talk*.

Another derivation is to produce first (using S_8)

- (4) *try to walk*.

Next we produce (5), using S_{11} .

- (5) *try to walk and talk*.

Application of S_4 to (5) yields (3), but the correct form of a sentence with the intended conjunction would be

(6) *John tries to walk and talks.*

In order to correct rule S_4 for this, it is useful to distinguish the conjunction of *try to walk* and *talk* from the IV phrase *try to walk and talk*. So it is useful to assign structure to the strings (5) and (6).

This second argument shows that it is useful to have some kind of structure available, not that it has to be the kind of structures used in transformational grammars. As has been shown by FRIEDMAN (1979), the kind of problems mentioned above can be dealt with by un-labelled trees. A completely different kind of syntactic structures is used in JANSSEN 1981b, where the present framework is combined with the structures used in Dik's 'functional grammar' (DIK 1978,1980). However, the kind of structures I will consider in this chapter are constituent structures of the kind described above.

2. THEORETICAL ASPECTS

2.1. Trees in Montague grammar

The rules of a Montague grammar determine how basic syntactic units are combined to larger ones. Such production processes can be represented by a tree. The tree for the de-dicto reading of *John seeks a unicorn* is given in figure 2.




Figure 2 tree from Montague grammar

Such trees are representations of derivational histories. For this reason PARTEE(1975) compares them with the T-markers from transformational grammar, and not with the produced trees themselves. In transformational grammars trees are produced, and if one wishes to compare the approach of Montague grammar

to the approach of transformational grammar, then one has to compare trees. Trees like the one in figure 2 are the only trees one finds in publications of Montague. Therefore one is tempted to compare such trees with the trees obtained in transformational grammars.

The tree in figure 2 is not of the form of the trees of transformational grammars. The main difference is that in transformational grammars the nodes are not labelled with expressions, but with category symbols (except for the end-nodes). Therefore one considers the tree from figure 2 as an unusual representation of the tree given in figure 1. Then the tree from figure 2 is taken as the syntactic structure assigned to the sentence by the PTQ grammar. Proceeding in this way, using the only trees available in Montague grammars, it becomes possible to compare the structures in Montague grammar with the structures in transformational grammars. This view on syntactic structure in Montague grammar can be found in work of several authors. In the next chapter we will see that PARTEE (1973) has compared the relative clause formation in Montague grammar and in transformational grammar by comparing trees like the one in figure 2, with those of transformational grammars. This way of discussion was followed up in by BACH & COOPER (1978). The same approach can be found in COOPER & PARSONS (1976). They describe a transformational grammar that is claimed to be equivalent with the PTQ system. The base rules of their transformational grammar produce (roughly) the same trees as the derivational histories of PTQ.

If one just compares the trees in the two approaches one soon will find great differences, and problems arise if one wishes to take the trees from Montague grammar as serious proposals for the syntactic structure assigned to a sentence. Consider the tree for the de-re reading of *John seeks a unicorn*, given in figure 3, or alternatively the one in figure 4.




Figure 3: de-re reading




Figure 4: variant of figure 3

This tree cannot be taken as a serious analysis of the constituent structure of the sentence since it does not even fulfill the weakest requirement: that the lexical material is presented in the correct sequence.

Cooper has developed a variant of Montague grammar in which no quantification rules are used, and which seems to eliminate the problem just mentioned. I give an example from COOPER 1978 (the idea originates from COOPER 1975). Consider the tree in figure 2. The interpretation of this tree follows its structure. The lexical items are interpreted first, and next the interpretations of larger constituents are formed. The usual interpretation yields the de-dicto reading, roughly presented as $John'(seek'(a unicorn'))$. The de-re interpretation is obtained by means of a mechanism which gives the translation of *the unicorn* wide scope. This mechanism simply is a storage mechanism which allows the translation of the noun phrase *a unicorn* to be stored, putting a variable placeholder in the regular translation. The stored translation of a unicorn is carried up the tree, until it can be retrieved at a suitable point where quantifying in is allowed. The store is a set of pairs consisting of an interpretation of a term and a variable. The way of processing is as follows.

$$\begin{array}{lcl}
 a \text{ unicorn} & \rightsquigarrow & \langle \lambda P^V P(x_0), \langle a \text{ unicorn}', x_0 \rangle \rangle \\
 seek \ a \ \text{unicorn} & \rightsquigarrow & \langle seek'(\lambda P^V P(x_0)), \langle a \text{ unicorn}', x_0 \rangle \rangle \\
 John \ seeks \ a \ \text{unicorn} & \rightsquigarrow & \langle John'(\wedge seek'(\lambda P^V P(x_0))), \langle a \text{ unicorn}', x_0 \rangle \rangle \\
 & & \text{retrieve from store, yielding} \\
 & & \langle a \text{ unicorn}'(\lambda x_0 (John'(\wedge seek(\lambda P^V P(x_0))))), \emptyset \rangle.
 \end{array}$$

Cooper is not very explicit about the details of his proposal, and therefore it is difficult to evaluate it. Nevertheless, I have serious doubts about the acceptability of his proposal in any approach which accepts the principle of compositionality of meaning. The reason for this is as follows. The phrase *seek a unicorn* has two parts: *seek* and *a unicorn*. The contribution of the latter part to the meaning of the whole phrase consists in three components, one of them being the variable x_0 . We have formalized meanings as abstract functions (intensions), and the symbol x_0 is not an element in this formalization. I assume that Cooper does not intend to define meanings as something which has the symbol x_0 as a component. So the mechanism does not build meanings from meanings, and therefore it violates the principle of compositionality of meaning. A more explicit description of a storage mechanism is given in PARTEE & BACH (1981); that

proposal is discussed in LANDMAN & MOERDIJK (1983), where it is shown that related objections apply.

The above discussion shows that we cannot get rid of trees like the one given in figure 3 by using Cooper storage. This has the following consequence. If one takes the tree representing the derivational history of a sentence in a Montague grammar to be the syntactic structure assigned to that sentence, then one has to conclude that in certain cases they are unacceptable as constituent structures. This is a practical reason against identifying the derivational histories with constituent structures. As will be explained below, there are also algebraic reasons against it.

2.2. Algebraic considerations

In our framework the syntax is an algebra, i.e. a collection of carriers with operations defined on them. An algebra can be defined in many ways. For instance, one can enumerate all the elements of each carrier, and state what the operators are. But we have developed a more efficient way of defining an algebra: state what the generators and the operators are. In this way with each element of the algebra (at least) one derivational history can be associated. Such derivational histories are important for the semantics, because this process is mirrored when building the corresponding meanings. We have met several examples where the choice of a certain generated algebra was determined by semantic considerations. If we consider only the syntactic side of the situation, the generation process is just some method to define the algebra. If we would replace a given definition by another definition of the same algebra, the elements and the operators would remain the same. More in particular, an element of an algebra by itself does not have a derivational history. Only if one has additional information concerning the way in which the algebra is defined, it becomes possible to associate with an element some derivational history, and with the algebra itself an algebra of derivational histories (a term algebra). The operators of a syntactic algebra are functions defined on the elements of that algebra, and since the information how the algebra is defined, cannot be read off from these elements, the operators of the syntactic algebra cannot interfere with derivational histories. In section 2 I argued that we need syntactic structures in order to design more sophisticated rules. As argued above, the syntactic rules are completely independent of such histories. Hence we cannot consider derivational histories to be the

structures we are looking for. This means that the only available trees cannot be used as a kind of syntactic structures. So the conclusion has to be that the PTQ grammar assigns *no structure at all* to the expressions it deals with.

If one wants the elements of an algebra to *have a structure*, then these elements *should be structures!* So in order to obtain a syntactic structure for the expression of a Montague grammar, this grammar should produce structures: trees, or, equivalently, labelled bracketings. This brings us to an approach dating from the first years of Montague grammar: PARTEE 1973. That proposal follows the sound approach to structure in Montague grammar. It distinguishes between the structure of the derivational history and the structure of the produced element itself. A remark about the relevance of distinguishing these two levels in a grammar for natural language can already be found in CURRY (1961), who calls the level of history 'tectogrammat-ics', and the level of produced expressions 'phenogrammat-ics'. DOWTY 1982 claims that rather different languages (such as Japanese and English) may have the same tectogrammatic structures, whereas the differences between the languages are due to phenogrammatical differences. This idea can also be found in LANDSBERGEN 1982, where it constitutes the basic idea for a computer program for automatic translation. In figure 5 the two kinds of structure are presented for the de-re reading of *John seeks a unicorn*: the trees within the squares are the constituent structures produced by the grammar, and the tree consisting of double lines with squares as nodes is the tree representing the derivational history.

2.3. Practical differences

Above I argued on algebraic grounds for distinguishing the structure an element has, from the derivational history assigned to it in some generative definition of the algebra. A practical aspect of this distinction is that there are completely different criteria for the design of these two kinds of structures. The derivational history is mapped homomorphically to the semantic algebra and determines the meaning of the expression. Semantic considerations play a decisive role in the design of the operators, and considerations concerning efficiency of definition determine the choice of the generators. The inherent constituent structure of the expressions is determined by syntactic considerations, e.g. the role such a structure has to play in the description of the syntactic effect of an operation. These




Figure 5 One derivational history containing many constituent structures

two different kinds of arguments may yield different kinds of structures. Below I will give some examples which show that the derivational history may sometimes differ considerably from what an acceptable constituent structure might be.

- a) The PTQ rule S_{14} produces e.g. *John runs out of John and He₁ runs*. This is not an acceptable syntactic structure since it contains at an end node a word that does not occur in the sentence (cf. the discussion concerning figure 3).
- b) In the grammar for questions by GROENENDIJK & STOKHOF (1981), there is a rule which substitutes a common noun into phrases of a certain kind. Thus *which man walks* is produced out of *man* and *which one walks*. Here the same argument applies as for the quantification rule of PTQ: it contains at an end node an argument that does not occur in the sentence.
- c) In HAUSSER (1979b) another variant is presented of the substitution of a common noun for an occurrence of *one* in some phrase. Here the same conclusion holds.
- d) BACH (1979a) presents rules which produce *persuade Bill to leave* out of *Bill* and *persuade to leave*. The operation which performs this task, called 'right-wrap', is a kind of substitution operation. It disturbs the sequence of words, and therefore it gives rise to a derivational history in which the order of the words does not correspond with the order of the words in the phrase. Therefore the derivational history is different from any possible syntactic structure.
- e) DOWTY (1978) gives a very elegant categorial treatment of phenomena which are traditionally treated by transformations. Examples are dative movement and object deletion. His rules shift *serve* from the category DTV (takes a dative and a term), to the category TTV (takes two terms), and next to TV and IV. This history is presented in figure 6. As far as I know, such a structure has not been proposed in transformational grammars, which is an indication that there is no syntactic motivation for this structural analysis. All steps in this production process are semantically relevant, and I consider it as a prime example of a semantically motivated elegant design of a derivational history.




Figure 6: History à la Dowty

3. TECHNICAL ASPECTS

3.1. Introduction

In this section I will sketch some tools which are useful in a version of Montague grammar in which the syntax produces structured expressions. The desire to provide handsome tools for a certain limited purpose leads to restricted tools (all-purpose tools are usually not very handsome: I would not like to describe a language by means of a Turing machine). So, whereas I do not have the aim of Partee ('defining as narrowly as possible the class of possible grammars of natural languages' (PARTEE 1979b, p.276)), the practical work is closely related. The tools I will use originate mainly from Partee (ibid); in the details there are some differences. It is not my aim to develop a complete theory about structured syntax, but I will use the opportunity to make some theoretical and practical remarks about the available techniques. For more ambitious proposals which use the same approach to structure, see BACH 1979b, PARTEE 1979a, 1979b, and LANDMAN & MOERDIJK 1981.

The basic change I will make here in comparison with previous chapters, is a change of the algebra on the background, which is always assumed when we define a generated algebra. In the previous chapters this was mostly the algebra consisting of all strings over the alphabet involved with concatenation as operator. In the present chapter this background algebra is replaced by one which consists of all trees, labelled in an appropriate way, and which has the basic operations which will be described in the sequel.

3.2. Operations on trees

It is not very convenient to describe operations on trees by means of English sentences. Following Partee, I describe such operations as the composition of a few basic ones. These are described in the sequel.

The operation *root* gives a new common root to the members of a list of trees. The new root is labelled with a given category name. Let α and β denote trees, and let (α, β) denote the list consisting of these two trees. The effect of $\text{root}((\alpha, \beta), \text{IV})$ is that the roots of the trees α and β are connected with a new root, labelled IV, see figure 7.




Figure 7: $\text{root}((\alpha, \beta), \text{IV})$

The operation *insert* substitutes a tree for a given node in some other tree. Let us accept the phrase 'first he_2 in (α) ' as a correct description of the node marked with x in tree α , see figure 8. Then the effect of $\text{insert}(\beta, \text{first } he_n \text{ in } (\alpha))$ is given in figure 9. A single word is considered as the denotation of a tree consisting of one node, labelled with that word. So the root operation can be applied to it. The effect of $\text{root}(\text{and}, \text{Con})$ is shown in figure 10.




Figure 8: situation

Figure 9: $\text{insert}(\alpha, \beta, x)$

Figure 10: $\text{root}(\text{and}, \text{Con})$

These two operations for tree manipulation, together with operations for feature manipulation and index manipulation, suffice for the treatment of the PTQ fragment. For larger fragments other operations might be required. An example is 'everywhere-substitution', which has the effect of substitution for all occurrences of a variable. This effect cannot be obtained by means of a repetition of the *insert* operator since one and the same tree cannot be at the same time the daughter of different nodes. So everywhere-substitution requires a copy operation which might be added as a primitive operation. PARTEE 1979b has no copy operation, but considers everywhere-substitution as a basic operation (we do not need everywhere-substitution since we deal with $S_{14,n}$ by means of an operation on features).

As an example I present the rule for verb-phrase conjunction. If we stay close to PTQ, it gets the following form, and yields the result given in figure 11.

$$S_{11}: IV \times IV \rightarrow IV$$

$$F_{11}: \text{root}((\alpha, \text{and}, \beta), IV).$$

One might prefer to give the connective *and* a categorial status in the syntactic structure; the status of a connective. Then the operation could read as follows, yielding the result given in figure 12.

$$S_{11}: IV \times IV \rightarrow IV$$

$$F_{11}: \text{root}((\alpha, \text{root}(\text{and}, \text{Con}), \beta), IV)$$





Figure 11: $\text{root}((\alpha, \text{and}, \beta), IV)$ Figure 12: $\text{root}((\alpha, \text{root}(\text{and}, \text{Con}), \beta), IV)$

3.3. Features and lexicon

Rule S_4 of PTQ tells that the subject-verb agreement in a sentence is obtained by replacing the first verb by its third person singular present. This is not an explicit formulation of what the effect of the rule is supposed to be. In an explicit form it would say that the verb *run* is to be replaced by *runs* and that *try to* (in PTQ a single word with a space inside)

is to be replaced by *tries to*. Rule S_4 can have its short readable form only since it is not explicit about such details. In order to obtain an explicit syntactic rule which is not full with details, we have to abstract from the inflection behaviour of the individual verbs. So it is useful to let the syntactic rules deal with more abstract lexical elements. By incorporating features in Montague grammar, rule S_4 may for the PTQ fragment simply attach the features like *present* and *third person singular* to an elementary form of the verb without being concerned with the effect of these details for every individual verb. The information about morphological behaviour of the verb can be given in the lexicon or in a separate morphological component.

Features originate from transformational grammar. They were used, as far as I know, for the first time in Montague grammar by GROENENDIJK & STOKHOF 1976 for a phenomenon like the one above. Features are also useful if one incorporates transformations into Montague grammar. PARTEE 1979b gives several examples of transformations which require features; an example is the Subject-Aux inversion process for questions which requires isolation of the tense morpheme.

As an example of the use of features I give a reformulation of rule S_4 using features. Of course, the rule has in other respects the same shortcomings as the original PTQ rule, but it is fully explicit now.

$$S_4: T \times IV \rightarrow S$$

$$F_4: \text{add features}((pres, sing\ 3), \text{first verb in } (\alpha)); \\ \text{root}((\alpha, \beta), S).$$

The details of the regular formation of the word forms can be given on a separate morphological component, whereas details about irregular word forms can be given in the lexicon. So the function *verbform* in the morphological component will be such that

$$\begin{aligned} \text{verbform}((pres, sing3), \alpha) &= as && (\text{e.g. } walks) \\ \text{verbform}((pst, sing3), \alpha) &= aed && (\text{e.g. } walked). \end{aligned}$$

The morphological component also contains a function *pronomen* such that

$$\begin{aligned} \text{pronomen}(sing3, acc, masc) &= him \\ \text{pronomen}(sing3, nom, neut) &= it. \end{aligned}$$

Besides morphological details, the lexicon also contains the information which features are inherent to the word (e.g. *John* always bears the feature

sing3) and information about kinds of features for which the word may be specified (e.g. *John* may not be specified for tense).

On the basis on the above considerations I define a lexical element as a list consisting of the following five components.

1. a string being the basic form of the word
2. a category symbol, being the category of the lexical element
3. a list of inherent features
4. a list of kinds of features for which the lexical element can be specified.
5. a description of the procedure for making derived forms of the word.

The above definition says that a lexical element is denoted by a list of five elements, of which some are lists themselves. We already introduced a notation for lists. Let furthermore () denote an empty list. The examples of lexical elements presented below are somewhat simplified with respect to PTQ since they only consider past and present tense.

```

("John",T, (masc,,sing3),(), wordform: "john")
("walk",IV,(),(tense,pres), wordform: verbform((tense,pres),"walk"), )
("run",IV,(),(tense,pres),

  if value of tense = past then wordform: "ran"
  else wordform: verbform((pres,sing3),"run") ).

```

Up till now we only considered kinds of features which are well known. But nothing in the feature definition prohibits us to define unusual ones. We might define a feature kind 'mainverb' with values # and -. The instructions for introduction or deletion of this feature can be the same as the instructions for Bennetts # mark which indicates the main verbs of a phrase (BENNETT 1976). In this way we can use the features as syntactic markers. Following Partee, I would not like to do so. Features are introduced for isolating morphological phenomena, not for syntactic marking. So I would like to restrict features to morphological relevant ones, just as PARTEE (1979b) proposed. This restriction requires, however, a formal definition of this notion (Partee gives no definition).

The notion 'morphologically relevant feature' is clearly word dependent. The case feature is relevant for *he* but not for *John*. So we might call a feature morphologically relevant if it is relevant for at least some word in the grammar. But what does this notion mean? Something like that the feature influences the form of the word? It is to be required further-

more that this influence can be observed in real sentences: it is not enough that it occurs in some morphological rule since this leaves open the possibility of a fake feature which influences the form of some 'external' word that does not occur in a produced sentence. We want a feature to create an opposition of wordforms in the produced language. Based upon these considerations I would define the notion as follows.

3.1. DEFINITION. A feature F is called *morphologically relevant* in grammar G if the following two conditions are satisfied.

1. There is a sentence $S_1 \in L(G)$ containing a lexical element W which bears feature F and which has wordform W_1 .
2. There is a sentence $S_2 \in L(G)$ containing an occurrence of W which does not bear feature F and which has wordform W_2 where W_2 is different from W_1 .

3.1. END

Note that this definition uses quantification over the sentences in the language $L(G)$. This quantification makes the notion 'morphologically relevant' to an undecidable notion. Suppose a list of syntactic rules, a lexicon containing features, and a list of morphological rules is given. Then one might try to show that a feature is not morphologically relevant by producing a lot of sentences and checking the conditions. However, one never reaches a stage that one can say for sure that such a feature is unacceptable. A formal proof is given in the following theorem.

3.2. THEOREM. *There exists no algorithm which decides for all grammars G and feature F whether F is morphologically relevant in G .*

PROOF. Suppose that such an algorithm would exist. Then this would give rise to a decision procedure for algebraic grammars, as will be shown below. Let G be an arbitrary algebraic grammar, with distinguished sort S , and suppose $L(G)$ is a language over the alphabet A . Let α be an arbitrary string over this alphabet, and let $w \in A$ be the first symbol of α . Let $w' \in A$ be a new symbol, and F a new feature not occurring in G . Define the wordform of w when bearing feature F as being w' . Extend now grammar G to G' by adding the following rule:

R: $S \rightarrow S$ is defined by

$R(\alpha) = \alpha'$ where α' is obtained from α by attaching F to w

$R(\phi) = \phi$ if ϕ is not equal to α .

The only way to introduce w' in some expression of $L(G')$ is by means of this new rule R . Hence F is morphologically relevant in G' if and only if $\alpha' \in L(G')$. From the definition of R it follows that $\alpha' \in L(G')$ iff $\alpha \in L(G)$. So if it would be decidable whether F is morphologically relevant, it would be decidable whether α is generated by grammar C . Since $L(G)$ can be any recursively enumerable language, this question is undecidable.

3.2. END

The undecidability of the notion 'morphologically relevant' has as a consequence that it can not be considered as a formal constraint, and that it cannot be incorporated in the definition of the notion 'grammar'. This does not mean that the property is worthless. It could play about the same role as the well-formedness constraint, being an important practical guideline for designing and evaluating grammars.

3.4. Queries for information

In syntax one often uses information about the grammatical function of words and groups of words. The grammatical tradition has constituted names for most of these functions, e.g. mainverb, subject and object. That the information for determining these functions is present in the syntactic structure assigned to them, has already been stated in CHOMSKY 1965. He defines the subject of a sentence as the NP which is immediately dominated by the main S node. In spite of this approach to grammatical functions, the tradition of transformational grammar never uses such information explicitly. PARTEE 1979b proposes to incorporate this information in Montague grammar and to make explicit use of it in the syntactic rules.

On the question what the main verbs of a sentence are, an answer like *run* is not good enough since that verb might occur more than once. An answer has to consist of a list of occurrences of verbs; or formulated otherwise a list of nodes of the tree which are labelled with a verb. Functions used to obtain syntactic information such as mainverb are functions from trees to lists of nodes of that tree. The first use of functions of this kind in Montague grammar is given in FRIEDMAN 1979. Such functions are called queries by KLEIN (1979); PARTEE (1979b) uses the name properties for a related kind of functions. The different name covers a different approach to such functions. It is a property of each individual occurrence of a verb to be a mainverb or not so: hence a property is a boolean valued

function, but a query is not. Since it is not convenient to use properties in syntactic rules, I use queries.

PARTEE (1979b) defines queries by means of rules parallel to the formation rules of the syntax. This has as a consequence that she in fact performs induction on the trees which represent derivational histories. Thus properties of derivational histories can be defined by means of her queries. It allows, for instance to define a query which tells us what the terms are which are introduced by means of a quantification rule. This query which I call 'substituted terms', can be defined as follows:

1. add to rule $S_{14,n}$ the clause
 substituted terms ($S_{14,n}(\alpha, \beta)$) = $\{\alpha\} \cup$ substituted terms (β)
2. do the same for the other quantification rules
3. add to the other rules the clause
 substituted terms ($S_i(\alpha, \beta)$) = substituted terms(α) \cup
 substituted terms(β)

Since we do not consider the derivational histories as representations of syntactic structures, we do not want information about the derivational history to be available in the syntax. Therefore I will not define queries in this way.

FRIEDMAN (1979) defines queries separately from the rules. She defines them for all constituent structures by means of a recursive definition with several clauses; each clause deals with a different configuration at the node under consideration. So Friedman performs induction on the constituent trees, and not on the derivational histories. Consequently, the query 'substituted terms' cannot be defined in Friedman's approach. In principle I will follow Friedman's method, but some modifications are useful.

Friedman's method has a minor practical disadvantage. If one adds a rule to the grammar, then at several places the grammar has to be changed: not only a rule is added, but all query definitions have to be adapted. In order to concentrate all these changes on one place in the grammar, I will mention the clauses of the definitions of a query within the operation which creates a new node, so within the operation root. In this way it is for each node determined how the answer to a query is built up from the answers to the query at its subnodes. If a root operation contains no specifications for a query, this is to be understood as that the answer to the query always consists of an empty list of nodes. As an example, I present rule S_{11a} , in which a clause of the query mainverb is defined.

$$S_{11}: IV \times IV \rightarrow IV$$

$$F_{11}: \text{root}((\alpha, \text{root}(\text{and}, \text{Con}), \beta), IV, \\ \text{mainverbs} = \text{mainverbs}(\alpha) \cup \text{mainverbs}(\beta)).$$

The basis of the recursion for a query is formed by its application to a node only dominating an end node of the tree. Then a query yields as result that end node. So is the query `mainverbs` is applied to the root of the tree in figure 13, then the result is that occurrence of *run*.



Figure 13: basis of recursion

4. PTQ SYNTAX

Below I will present a syntax for the PTQ fragment. The purpose of this section is to provide an explicit example of what a Montague grammar in which structures are used, might look like. It is not my aim to improve all syntactic shortcomings of PTQ; only some (concerning conjoined phrases) are corrected. For more ambitious proposals see PARTEE 1979b and BACH 1979.

In the formulation of the rules, several syntactic functions and operations will be used. Below the terminology for them is explained, thereafter they will be described.

1. *Queries*

Functions which have a tree as argument and yield a list of nodes in the tree. They are defined within the root operations.

2. *Primitive functions*

Functions of several types which yield information, but do not change anything.

3. *Primitive operation*

Operations of several types which perform some change of the tree or lists involved.

4. *Composed operations*

Like 3, but now built from other operations and functions.

QUERIES

Mainverbs

Yields a list of those occurrences of verbs in the tree which are the mainverbs of the construction.

Headnouns

Yields a list of those occurrences of nouns, pronouns and proper names which are the heads of the construction.

PRIMITIVE FUNCTIONS

Index of (w)

yields the index of the word w (provided that w is a variable)

First of (l)

yields the first element of list l.

All occurrences of (he_n , t)

yields a list of all occurrences of he_n in tree t.

Gender of (w)

yields the gender of word w, and of the first word of w if w is a list.

Is a variable (w)

determines whether term w is a variable (of the form he_n).

PRIMITIVE OPERATIONS

root((t_1, \dots, t_n), C, query:...).

Creates a new node which is the mother of the trees t_1, \dots, t_n .

This new node is labelled with category symbol C.

For all queries a clause of their recursive definitions is determined:

either explicitly, or implicitly (in case the query yields an empty list).

Add features (f, l)

Attaches to all elements of list l the features in feature list f.

Delete index (n, l)

Deletes index n from all elements in list l.

Replace index (l, m)

Replaces the index of all variables in list l by index m.

Insert (r, t)

Replaces node r by tree t, thus inserting a tree in another tree.

Union (l_1, l_2)

Yields one list, being the concatenation of lists l_1 and l_2 .

COMPOSED OPERATIONS

Termsubstitution (t,n,r)

Substitutes term t in tree r for the first occurrence of he_n . The operation is defined by:

if is a variable (t)

then replace index (all occurrences of (he_n, r) , index of (t))

else insert (t, first of (all occurrences of (he_n, r)))

define: list=all occurrences of (he_n, r)

delete index(list)

add features ((sing,pers3,gender of (first of (main nouns (t))), list).

End definition.

Below the rules for the PTQ-fragment are given with exception of the rules for tense. Their formulation resembles the formulation they would get in an ALGOL 68 computer program I once thought of.

S_2 : Det \times CN \rightarrow T

root $((\alpha, \beta), T,$

head nouns = head nouns (β)).

$S_{3,n}$: CN \times S \rightarrow CN

define: list = all occurrences of (he_n, β) ;

delete index (n, list);

add features (gender of (head nouns (α), list));

root $((\alpha, \text{root}(\text{such-that}, \text{Rel}), \beta), \text{CN},$

head nouns = head nouns (α)).

S_4 : T \times IV \rightarrow S

add features $((\text{pres}, \text{sing3}), \text{main verbs}(\beta));$

add features $(\text{nom}, \text{head nouns}(\alpha));$

root $((\alpha, \beta), S,$

main verbs = main verbs (β)).

S_5 : TV \times T \rightarrow IV

add features $(\text{acc}, \text{head nouns}(\beta));$

root $((\alpha, \beta), IV,$

main verbs = main verbs (α)).

S_6 : Prep \times T \rightarrow IAV

add features $(\text{acc}, \text{head nouns}(\beta));$

root $((\alpha, \beta), \text{IAV}).$


- $S_7: IV/S \times S \rightarrow IV$
 root $((\alpha, \beta), IV,$
 main verbs = main verbs (α)).
- $S_8: IV//IV \times IV \rightarrow IV$
 root $((\alpha, \beta), IV,$
 main verbs = main verbs (α)).
- $S_9: S/S \times S \rightarrow S$
 root $((\alpha, \beta), S,$
 main verbs = main verbs (β)).
- $S_{10}: IAV \times IV \rightarrow IV$
 root $((\beta, \alpha), IV,$
 main verbs = main verbs (β))
- $S_{11a}: S \times S \rightarrow S$
 root $((\alpha, \text{root}(\textit{and}, \text{Con}), \beta), S,$
 main verbs = union (main verbs (α) , main verbs (β)).
- $S_{11b}: S \times S \rightarrow S$
 root $((\alpha, \text{root}(\textit{or}, \text{Con}), \beta), S$
 main verbs = union (main verbs (α) , main verbs (β)).
- $S_{12a}: IV \times IV \rightarrow IV$
 root $((\alpha, \text{root}(\textit{and}, \text{Con}), \beta), IV,$
 main verbs = union (main verbs (α) , main verbs (β)).
- $S_{12b}: IV \times IV \rightarrow IV$
 root $((\alpha, \text{root}(\textit{or}, \text{Con}), \beta), IV,$
 main verbs = union (main verbs (α) , main verbs (β)).
- $S_{13}: T \times T \rightarrow T$
 root $((\alpha, \text{root}(\textit{or}, \text{Con}), \beta), T,$
 head noun = union (head nouns (α) , head nouns (β)).
- $S_{14,n}: T \times S \rightarrow S$
 termsubstitution (α, n, β) .
- $S_{15,n}: T \times CN \rightarrow CN$
 termsubstitution (α, n, β) .
- $S_{16,n}: T \times IV \rightarrow IV$
 termsubstitution (α, n, β) .

CHAPTER IX

RELATIVE CLAUSE FORMATION

ABSTRACT

Does the principle of compositionality compel us to a certain analysis of relative clause constructions? Answers given by Partee and Bach & Cooper will be investigated, and new arguments will be put forward. The question will be generalized and answered on the basis of algebraic properties of the framework. The investigations give rise to a restriction on the use of variables in Montague grammar: the variable principle.



1. INTRODUCTION

Our framework, which formalizes the principle of compositionality of meaning, says that the syntax and semantics are similar algebras, and that the meaning assignment function is a homomorphism. Now one may ask to what extent this organization of the grammar restricts the options we have in the syntax to describe a particular phenomenon. This question was raised by PARTEE (1973) with respect to relative clause constructions, and her answer was that we have to use a particular analysis. She concluded that the framework puts very strong constraints on the syntax, with the consequence that 'it is a serious open question whether natural language can be so described' (PARTEE 1973, p.55). Her argumentation is used by CHOMSKY (1975) to support his ideas of an autonomous syntax in transformational grammars. Partee's conclusion about relative clause formation has been disputed by BACH & COOPER (1978), who give an alternative construction.

In chapter 2 it has been proven that every recursively enumerable language can be described by means of a finite algebraic grammar. Hence Partee's question, as quoted above, has already been answered positively. But we will curtail it to the question whether the framework constrains the way in which natural language phenomena can be described. More in particular, we will investigate the thematic question: *does the framework of Montague grammar compel us to a particular syntactic analysis of restrictive relative clauses?* The arguments given in the literature will be considered, and new arguments will be put forward. In the course of the discussion positive and negative answers to the thematic question will alternate. An answer to the general version of the question is obtained as well. It will turn out that syntactic variables (like he_n) play an important role in relative clause constructions. This role is investigated, and this gives rise to the introduction of a new principle for Montague grammar: the variable principle. This chapter is a slightly revised version of JANSSEN (1981a).

2. THE CN-S ANALYSIS

2.1. The discussion by Partee

PARTEE (1973) considers three kinds of analyses of relative clause constructions which were proposed in the literature in the framework of transformational grammar. She investigates which of them constitutes a good

basis for a compositional semantics. The comparison is carried out in the way described in chapter 7, section 2.1: the derivational histories from Montague grammar are compared with the constituent structures proposed in transformational grammars. As was explained there, this is not the most felicitous way to compare the two approaches. Our thematic question, however, does not concern a comparison but is a question about the present framework itself: the structures from transformational grammar merely constitute a starting point. Hence all trees under discussion have to be taken as representing derivational histories, even in case they originate from transformational grammar as constituent structures. In the sequel I will use the categorial terminology from the previous chapters, and not the transformational terminology used in the proposals under discussion.

Below I summarize Partee's argumentation. She discusses three kinds of analysis for the restrictive relative clause construction. They are named after the configuration in which the relative clause is introduced. These analyses (of which the second was the most popular among transformational grammarians) are

1. CN-S : the Common Noun-Sentence analysis (Figure 1)
2. T-S : the Term-Sentence analysis (Figure 2)
2. Det-S: the Determiner-Sentence analysis (Figure 3).




Figure 1: CN-S




Figure 2: T-S




Figure 3: Det-S

In the analysis presented in Figure 1, the common noun *boy* can be interpreted as expressing the property of being a boy, and the phrase *who runs* as expressing the property of running. The conjunction of these properties is expressed by the noun phrase *boy who runs*. The determiner *the* expresses that there is one and only one individual which has these two properties. So the CN-S analysis provides a good basis for obtaining the desired meaning in a compositional way.

In the T-S analysis as presented in Figure 2, the term *the boy* is interpreted as expressing that there is one and only one individual with the property of being a boy. Then the information that the individual is running can only be additional. So in a compositional approach to semantics *who runs* has to be a non-restrictive relative clause. Therefore Partee's conclusion is that the T-S analysis does not provide a good basis for a compositional semantics of restrictive relative clauses.

The Det-S analysis from Figure 3 does not provide a good basis either. The phrase dominated by the uppermost Det-node (i.e. *the who runs*), expresses that there is one and only one individual with the property of running, and the information that this individual is a boy, can only be additional.

Of course, these arguments do not constitute a proof that it is impossible to obtain the desired meanings from the T-S and Det-S analyses. It is, in general, very difficult to prove that a given approach is not possible, because it is unlikely that one can be sure that all variants of a certain approach have been considered. This is noted by Partee when she says: 'I realize that negative arguments such as given against analyses 2. and 3 can never be fully conclusive. [...]' (PARTEE 1973, p.74 - numbers adapted T.J.). She proceeds: 'The argument against 3. is weaker than that against 2., since only in 2 the intermediate constituent is called a T.' (ibid.). Her carefully formulated conclusion is 'that a structure like 1, can provide a direct basis for the semantic interpretation in a way that 2 and 3 cannot' (ibid. p.54).

2.2. The PTQ-rules

Accepting the argumentation given in Section 2.1, is not sufficient to accept the claim that one should use the CN-S analysis. It remains to be shown that such an analysis is indeed possible, and this means providing explicit syntactic and semantic rules. Partee does not need to do so because in her discussion she assumes the rules for relative clause formation which are given in PTQ. Although these rules do not produce literally the same string as she discusses, the same argumentation applies to them.

I recall the rule for relative clause formation given in chapter 4.

$S_{3,n}$: $CN \times S \rightarrow CN$

$F_{3,n}$: Replace he_n in β by *he/she/it* and him_n by *him/her/it*, according to the gender of the first CN in α ; concatenate (α , *such that*, β)

$T_{3,n}$: $\lambda x_n [\alpha'(x_n) \wedge \beta']$.

According to this rule, the derivational history of *boy who runs* has the structure presented in figure 1. The phrase can be produced from *the* noun *boy* and the sentence *he₃ runs* by an application of instance $S_{3,3}$ of the above scheme. The corresponding translation reads

$$(1) \lambda x_3[\text{boy}(x_3) \wedge \text{run}(x_3)].$$

This expression is interpreted as the property which holds for an individual if it both is a boy and is running. This is completely in accordance with the interpretation sketched for figure 1.

I recall that $S_{3,n}$ can be applied two times in succession (or even more). Then sentences are obtained like (2) (due to Bresnan, see PARTEE 1975, p.263) and (3) (due to PARTEE - *ibid*).

(2) *Every girl who attended a women's college who made a large donation to it was included in the list.*

(3) *Every man who has lost a pen who does not find it will walk slowly.*

In these sentences two relative clauses are attached to a single head noun. This construction is known under the name stacking (of relative clauses). In Dutch and German stacking is not a grammatical construction.

Rules $S_{3,n}$ and $T_{3,n}$ do not give a correct treatment of all phenomena which arise in connection with relative clauses. Some examples are:

1. The rule produces the *such-that* form of relative clauses, and this is not their standard form. A rule which produces a form with relative pronouns cannot be obtained by means of a straightforward reformulation of $S_{3,n}$, since complications arise (see RODMAN 1976).
2. In certain circumstances $T_{3,n}$ may give rise to an, unintended, collision of variables. This problem was discussed in section 5.3 of chapter 6; see also section 6.1.
3. Some famous problematic sentences do not get a proper treatment with this rule. Examples are the so called 'Bach-Peters sentences' and the 'Donkey sentences'. There are several proposals for dealing with them. For instance HAUSSER (1979c) presents a treatment for the Bach-Peters sentence (4), and COOPER (1979) for the donkey sentence (5).

(4) *The man who deserves it gets the price he wants.*

(5) *Every man who owns a donkey beats it.*

For a large class of sentences, however, the PTQ rule yields correct results, and I will restrict the discussion to this class. The class

contains the relative clause constructions in the *such-that* form, the relative clause is a single (i.e. unconjoined) sentence, and stacking is allowed. Bach-Peters sentences and Donkey sentences are not included. For this class, the CN-S analysis gives a correct treatment in a compositional way, whereas for the T-S and Det-S analyses it is argued that this is not the case. So in this stage of our investigations, the answer to the thematic question has to be positive: the compositionality principle compels us to a certain analysis of relative clause constructions.

2.3. Fundamental problems

The PTQ rule for relative clause formation is essentially based on the use of variables in the syntax (he_n), and the use of unbound variables in the logic (x_n). This device gives rise to two problems which are of a more fundamental nature than the problems mentioned in Section 2.2. The latter concerned phenomena which were not described correctly by the given rule, but it is thinkable that some ingenious reformulation might deal with them. The fundamental problems I have in mind are problems which arise from the use of variables as such. It is essential for the entire approach to obtain a solution for these problems, since in case they are not solved satisfactorily we cannot use the tool at all. This aspect distinguishes them from the problems mentioned in Section 2.2. The problems also arise in connection with other rules dealing with variables ($S_{14,n}, \dots, S_{17,n}$). Note that the epithet 'fundamental' is not used to make a suggestion about the degree of difficulty of the problem, but to indicate the importance that some answer is given to it. The two fundamental problems are the following.

1) 'left-over'

The first problem is: what happens in case a variable is introduced that is never dealt with by $S_{3,n}$ or any other rule. On the syntactic side it means that we may end up with a sentence like he_γ runs. Since he_γ is not an English word, this is not a well-formed sentence, and something has to be done about it. On the semantic side it means that we may end up with an expression containing an unbound logical variable. From the discussion in Section 5 it will appear that it is not obvious how we should interpret the formulas thus obtained.

2) 'not-there'

The second problem is: what happens when a rule involving variables with a given index is applied in case such variables are not there. I give

two examples of such situations. The first is obtained if one applies $S_{3,1}$ to the common noun *man*, and the sentence *Mary talks*. Then the noun-phrase (6) is produced, which is ill-formed because there is no pronoun which is relativized.

(6) *man such that Mary talks.*

On the semantic side (6) gives rise to a lambda operator which does not bind a variable. The second example (GROENENDIJK & STOKHOF 1976b) is obtained by an application of $S_{3,1}$ to *man* and *he₂ walks*. Then the common noun phrase (7) is formed, out of which (8) can be obtained.

(7) *man such that he₂ walks.*

(8) *He₂ loves the man such that he₂ walks.*

By an application of $S_{14,2}$ we finally obtain

(9) *John loves the man such that he walks.*

This sentence has just one reading, viz. that John loves a walking man. The translation rules of PTQ however, yield (10) as reduced translation for (9).

(10) $\exists u[\forall v[[man_*(v) \wedge walk_*(john)] \leftrightarrow u = v] \wedge love_*(john,u)]$.

This formula expresses that the one who walks is John. THOMASON (1976) makes a related observation by counting the number of ambiguities of (11).

(11) *Bill tells his father that John resembles a man such that he shaves him.*

For the first problem it is evident that it is the use of variables which creates it, and that it are not the phenomena themselves: if there were no variables in the syntax, they could not be 'left-over', nor remain 'unbound' in their translation. For the second problem it is rather a matter of conviction that it is the use of variables that creates the problem. Even if (6) would be well-formed, I would consider its production in the way sketched above, as an undesirable side effect of the use of variables, because it does not exhibit a phenomenon for which variables are required.

In the literature there are some proposals for dealing with these two fundamental problems. One proposal (implicitly given in RODMAN 1976) is of a purely syntactic nature and simply says: the 'left-over' and 'not-there' constructions are not acceptable, and in case such a construction threatens to arise, it is filtered out. This approach is not considered here in detail, because it played no role in the discussion concerning our thematic

question. In the approach of COOPER (1975) the 'left-over' constructions are accepted, an answer is given to the semantic questions, and the 'not-there' constructions are dealt with in the semantics. In the next sections his proposal will be discussed in detail. A proposal combining syntactic and semantic aspects (JANSSEN 1980b) will be considered in Section 5.

3. THE T-S ANALYSIS

3.1. Cooper on Hittite

COOPER (1975) considers the construction in Hittite which corresponds to the relative clause construction in English. In Hittite the relative clause is a sentence which is adjoined to the left or the right of the main sentence. For this and other reasons, Cooper wishes to obtain such constructions by first producing two sentences and then concatenating them. A simplified example is the Hittite sentence which might be translated as (12), and has surface realization (13). The sentence is produced with the structure given in figure 4. For ease of discussion English lexical items are used instead of Hittite ones. 'Genitive' is abbreviated as 'gen', 'plural' as 'pl', 'particle' as 'ptc', and 'which' as 'wh'. The example is taken from BACH & COOPER (1978) (here and in the sequel category names are adapted).

(12) *And every hearth which is made of stones costs 1 shekel.*

(13) SA NA4 HI.A-ia kuies GUNNI.MES nu kuissa 1 GIN
gen.stone-pl.-and which hearth-pl. ptc. each(one) 1 shekel




Figure 4

Sentence (13) is assumed to have the same meaning as the corresponding English sentence (12). There seems to be a conflict between the arguments in favor of a CN-S analysis as given in section 2, and the wish to use the S-S analysis for Hittite. Cooper's solution is to allow the Term-phrase *each(one)* 'to denote the set of properties possessed by every entity having property R ' (BACH & COOPER 1978, p.147). Which property R is, is specified by the relative clause S_1 . The translations of S_2 and S_1 are (14) and (15), respectively (here and in the sequel \forall, \wedge and $*$ symbols are added).

(14) $\forall x[\forall R(x) \rightarrow \text{Cost-one-shekel}(x)]$

(15) $\text{Hearth}(z) \wedge \text{Made-of-stone}(z)$.

The syntactic rule which combines S_1 and S_2 to a phrase of the category \bar{S} , has as corresponding translation rule

$$\lambda R[S_2'](\wedge \lambda z[S_1']).$$

Here S_1' and S_2' are the translations of S_1 and S_2 , respectively. When this rule is applied to (14) and (15), we obtain (16) as reduced translation.

(16) $\forall x[\text{hearth}(x) \wedge \text{made-of-stone}(x) \rightarrow \text{cost-one-shekel}(x)]$.

Since \bar{S} is of another category than S_1 and S_2 , this production process does not allow for stacking, what is claimed to be correct for Hittite.

3.2. Bach & Cooper on English

BACH & COOPER (1978) argue that the treatment of COOPER (1975) of Hittite relative clauses can be used to obtain a T-S analysis for English relative clause constructions which is consistent with the compositionality principle. Terms are treated analogously to (the Hittite version of) *each(one)*. The term *every man* is assumed to denote, in addition to the PTQ interpretation, the set of properties possessed by *every man* which has the property R . Then the term-phrase *every man who loves Mary* is obtained from the structure given in figure 5.




Figure 5

The rule for combining the translation of the term and the relative clause is:

$$\lambda R[T'](\wedge S').$$

Here T' and S' are the translations of the term phrase and the relative clause, respectively. If we take (17) as translation of *every man*, and (18) as translation of the relative clause \bar{S} , then we obtain (19) as translation of the whole term (after reduction).

$$(17) \lambda P[\forall x[man(x) \wedge \forall R(x)] \rightarrow \forall P(x)]$$

$$(18) \lambda z[love_{*}(\forall z, mary)]$$

$$(19) \lambda P[\forall x[man(x) \wedge love_{*}(\forall x, mary)] \rightarrow \forall P(x)].$$

Thus a T-S analysis is obtained for relative clause constructions, of which the translation is equivalent to the translation in the case of a CN-S analysis.

As Bach and Cooper notice, if we follow this approach, a complication has to be solved, since English allows for indefinite stacking of relative clauses. The proposal sketched so far, provides for one relative clause for each T. The complication can be taken care of by allowing an alternative interpretation not only for Terms, but also for relative clauses. 'Thus, for example, the relative clause *who loves Mary* can denote not only the property of loving Mary but also the property of loving Mary and having property R' (BACH & COOPER 1978, p.149).

Bach and Cooper remark that their compositional treatment of the T-S analysis clearly is less elegant and simple than the alternative CN-S analysis. They conclude: 'Our results seem to indicate, however, that such an analysis cannot be ruled out in principle, since any constraint on the theory that would exclude the T-S analysis, would seem to exclude the Hittite analysis as well. [...] or the happy discovery of some as yet unknown principles will allow the one, but not other.' (ibid. p.149).

The conclusion which prompts itself in this stage of our investigations is that the answer to the thematic question is a negative one: the principle of compositionality does not compel us to a special analysis of English relative clauses.

3.3. Fundamental problems

As a matter of fact, the discussion in BACH & COOPER (1978) does not provide the evidence that a T-S analysis is indeed possible for English relative clauses. They do not present explicit rules, and neither is it immediately clear what the details would look like (e.g. what is the role of \bar{S} and COMP in the system of categories, and what is the translation rule which combines the translations of \bar{S} and COMP). Nevertheless, the main point of their approach has become clear from their exposition.

The kernel of the approach of Bach and Cooper is to let the translations of terms and relative clauses contain a free variable R . For this variable the translation of some relative clause will be substituted. However, this variable R gives rise to the same kind of problems as mentioned in section 1 with respect to the variables x_n .

1. 'Left-over'

We may select for a term the translation with free variable R , whereas we do not use in the remainder of the production a rule which deals with this variable. Since R has no syntactic counterpart, the produced sentences are not per se ill-formed, but the question concerning the interpretation of unbound variables remains to be answered.

2. 'Not-there'

There may be an occurrence of the term-phrase *every man* with the translation without R , nevertheless appearing in a structure where a relative clause is attached to it. Then an incorrect meaning is obtained.

Only when these fundamental problems are solved, we may hope that the idea of Bach and Cooper leads to rules for the T-S analysis. Notice that the proposal of RODMAN (1976) for solving the two fundamental problems by filtering them out, cannot be followed here because in the syntactic expressions there is no variable which may control the filter. A solution has to be found on the semantic side. These problems for the Bach-Cooper idea, are signalized for the case of Hittite by COOPER (1975). He has proposed some solutions which are assumed by Bach and Cooper. In order to obtain further justification for the answer to the thematic question given in Section 3.2, we have to check the details of Cooper's proposals for these problems. This will be done in the next section.

4. THE PROPOSALS OF COOPER

4.1. Not-there

A translation rule which usually binds a certain variable, may be used in a situation where no occurrences of such a variable are present. To avoid problems, Cooper proposes to give no semantic interpretation to expressions of intensional logic which contain a vacuous abstraction. According to his proposal the interpretation of $\lambda R\alpha$ is undefined in case α has no occurrences of R .

Let us first consider in which way this idea might be formalized. At first glance it seems easy to obtain the desired effect. One just has to look into the expression α in order to decide whether $\lambda R\alpha$ is defined or not. However, this is not acceptable. Such an approach would disturb the homomorphic interpretation of intensional logic: for each construction of the logical language there is a corresponding interpretation instruction. To obtain the interpretation of a compound logical expression, the interpretations of the parts of that compound are relevant, but not their actual forms. An important consequence of this is that two semantically equivalent expressions are interchangeable in all contexts. If we would have a condition like 'look into α ' in the definition of interpretation, this basic property of logic would no longer be valid. Two IL-expressions α and β might be semantically equivalent, whereas α satisfies the 'look into'-condition, and β not. Consequently, the interpretation of just one of $\lambda R\alpha$ and $\lambda R\beta$ would be defined. Such a violation of the fundamental law of substitution of equivalents is of course not acceptable, and therefore, a 'look into' clause has to be rejected. One has to respect the homomorphic interpretation of logic, and therefore, the situations in which $\lambda R\alpha$ should receive no interpretation have to be characterized in terms of the semantic properties of α (i.e. in terms of the interpretation of α with respect to a point of reference and a variable assignment). Cooper follows this strategy.

Cooper's first step towards a characterization consists of adding a restriction to the usual definition of the interpretation of $\lambda u\alpha$. '[...] the function denoted by the abstraction expression $\lambda u\alpha$ is only defined for entities within its domain if a different assignment to the variable u will yield a different denotation for α ' (COOPER 1975, p.246). As he notes, this definition has as a consequence that $\lambda u\alpha$ is 'undefined' not only if α does not contain a free occurrence of u , but also if α is

a tautology. Thus for instance, according to this definition $\lambda u[u=u]$ represents a function which is undefined for any entity. However, the technique of supervaluation [...] will show these expressions to be defined but not those where α is not a tautology' (ibid.). This definition is Cooper's final one, but it is not the one we need. It implies that now $\lambda R[x=x]$ is defined. This has the following consequence for relative clause formation. One might produce some sentence expressing a tautology, while its translation does not contain an occurrence of the variable R . Syntactically there needs not, in Cooper's approach, be anything which can prevent us from using this sentence in a relative clause construction, whereas, contrary to his intention, the interpretation of the translation is defined. So Cooper's definition does not provide a solution to the 'not-there' problem.

Cooper's aim was to give a semantic characterization of the IL-syntactic property 'contains an occurrence of the variable R '. I expect that there is no semantic property coinciding with the syntactic one. This is suggested by the observation that almost always a semantic irrelevant occurrence of a certain variable can be added to a given IL-expression. (ϕ and $R=R \wedge \phi$ are semantically indiscernable). Therefore, I expect that no solution in this direction can be found. Moreover, I consider the whole idea underlying Cooper's approach to be unsound. The standard interpretation of $\lambda R\alpha$ is, in case α does not contain an occurrence of R , a function that delivers for any argument of the right type, the interpretation of α as value. So $\lambda R\alpha$ denotes a constant function. Following Cooper's idea, one would lose this part of the expressive power of IL, a consequence I consider to be undesirable.

4.2. Left-over, proposal 1

The translation of a completed syntactic production of a sentence may contain an occurrence of a free variable. The second fundamental problem was what to do with variables that are 'left over'. Cooper proposes to assign no interpretation to such an expression, and to follow this approach for special variables only. Let z be such a variable (of the type of individuals). As was the case with the first problem, discussed in Section 4.1, one has to respect the homomorphic interpretation of IL. The desired effect should not be obtained by looking into the formula, but by changing the definition of interpretation. Cooper claims that the desired effect is obtained 'by restricting the assignments to variables so that z is always

assigned some particular non-entity for which no predicate is defined' (COOPER 1975, p.257). This proposal gives rise to a considerable deviation from the model for IL as it is defined in PTQ. In that model, there are for every entity predicates which hold for it, e.g. the predicate of being equal to itself (viz. $\lambda u[u=u]$). This property is lost in Cooper's approach. He does not define a model which has the desired properties, nor does he give other details. For the discussion concerning the thematic question, this point is not that relevant, because BACH & COOPER (1978) do not propose to follow this proposal in the case of English relative clause constructions, but another one, which will be discussed in Section 4.3.

4.3. Left-over, Proposal 2

A second proposal of COOPER (1975) for the treatment of unbound variables which occur in the translation of a completed production of a sentence is to let the unbound variables be interpreted by the variable assignment function, and to give some linguistic explanation of how to understand the results thus obtained. This approach assumes that in complete sentences indices of variables can be neglected, or that there is some final 'cleaning-up' rule which deletes the indices. For our discussion of relative clause formation the syntactic details of this proposal are irrelevant because the variable R leaves no trace in the syntax.

The unbound relative clause variable R only occurs in subexpressions of the form $R(x)$. These subexpressions are understood by Cooper as 'a way of representing pragmatic limitations on the scope of the quantifier [binding x].[...]. Thus assigning a value to R in this case has the same effect as adding an unexpressed relative clause to show which particular set we are quantifying over' (COOPER 1975, p.258-259). The same strategy is employed in COOPER (1979a,b) for indexed pronouns. A pronoun he_n that has not been dealt with by a relative clause formation rule or some other rule, is considered as a personal pronoun referring to some contextually determined individual. Its translation has introduced a variable x_n , which remains unbound, and is interpreted by the variable assignment. This idea for dealing with free variables is also employed in GROENENDIJK & STOKHOF (1976b). In one respect the idea leads to a deviation from PTQ. There, an expression of type t is defined to be true in case it denotes 1 for every variable assignment (MONTAGUE 1973, p.259). So, $run(x)$ would mean the same as its universal closure. In the proposal under discussion this definition

has to be dropped, but this does not cause any difficulties.

I have several objections against this proposal of Cooper. The first one is that it yields incorrect results; the other three argue that the whole approach is unsound. My objections are explained below.

1. If the translation of a phrase contains two occurrences of R , and a relative clause is combined with that phrase, then the translation of the relative clause is, by λ -conversion, substituted for both occurrences of R . As Cooper mentions, this phenomenon arises in his grammar for Hittite for (the Hittite variant of):

(20) *That(one) adorns that(one)*.

Here the translation of both occurrences of *that(one)* contains an occurrence of the variable R . If this sentence is combined with a sentence containing two occurrences of a *wh*-phrase, semantically strange things happen. Cooper notes this problem and he says: 'My intuition is, however, that if there were such sentences, they would not receive the interpretation assigned in this fragment. [...] As it is not clear to me what exactly the facts of Hittite are here I shall make no suggestions for improving the strange predictions of the fragment as it is.' (COOPER 1975, p.260).

Unfortunately, the proposal for English of BACH & COOPER (1978) runs into a related problem. Consider the structure for the term phrase given in Figure 6. It is an example taken from their article, and exhibits stacking of relative clauses (the structure is simplified by omitting Comp's).




Figure 6

The translation of *every man* has to contain a variable for the relative clause. Recall that, in the conception of Bach & Cooper, the proposal discussed in Section 4.1 deals with the situation that we have the translation not containing R . Let us assume that we have taken the translation (21), which contains an unbound variable R .

$$(21) \lambda P \forall x [man(x) \wedge \forall R(x) \rightarrow \forall P(x)].$$

Suppose now that the referent of a *girl* is to be contextually determined (this possibility is not considered by Bach & Cooper). Then the translation of a girl has to contain the variable R . Besides this variable the translation of (22) has to contain a variable R for the second relative clause. So the translation of (22) has to be (23).

(22) *who loves a girl*

$$(23) \lambda z \exists y [girl(y) \wedge \forall R(y) \wedge love_{*}(\forall z, \forall y) \wedge \forall R(z)].$$

Consequently, the translation of (24) has to be (25).

(24) *every man who loves a girl*

$$(25) \lambda P \forall x [man(x) \wedge \exists y [girl(y) \wedge \forall R(y) \wedge love_{*}(\forall x, \forall y) \wedge \forall R(x)] \rightarrow \forall P(x)].$$

The translation of *who lives in Amherst* roughly is indicated in (26).

$$(26) \lambda z [live-in-Amherst(z)].$$

The translation of the entire term-phrase in figure 6 is described by

$$(27) \lambda R [every\ man\ who\ loves\ a\ girl'] (who\ lives\ in\ Amherst').$$

This yields a logical expression which says that both the man and the girl live in Amherst, which is not the intended reading of the construction with stacked relative clauses.

These incorrect predictions are not restricted to stacking. The same problems arise in case a relative clause like *who runs* is combined with a disjoined term phrase like *the man or the woman*. Then semantically both terms are restricted, whereas syntactically only the second one is. The source of all these problems is that a single variable is used for relative clauses and for contextual restrictions. These two functions should, in my opinion, be separated. But then the left-over/not-there problem for relative clause variables arises with full force again.

2. As a motivation for interpreting the R 's as contextual restrictions, the argument was given that when we speak about every man, we in fact intend every man from a contextually determined set. But this argument applies with the same force in case we speak about every man who runs. It is not true that terms sometimes are contextually determined, and sometimes not. If one wishes to formalize contextual influence, then every term should be restricted. This suggests (as under 1) a system of variables for context

restrictions which is independent of the system of variables for relative clauses.

3. Variables of which the interpretation is derived from the context have to receive a very special treatment. This can be shown most clearly by considering a sentence which has as translation a formula containing an occurrence of an unbound variable of the type of individuals or individual concepts: *he runs*, obtained from the sentence he_n runs. These sentences have as translation $run(x_n)$. For every variable assignment this translation gets an interpretation. One of the possible assignments is that x_n is the person spoken to, so *He runs* would have the same truth conditions as *You run*. Some female person might be assigned to x_n , so the sentence may have the same truth-conditions as *she runs*. These are incorrect results, so there has to be some restriction on the variable assignments for x_n . There are also semantic arguments for such a restriction. A pronoun *he* usually refers to individuals from a rather small group (e.g. the person mentioned in the last sentence, or the person pointed at by the speaker). So again some restriction has to be given. These two sources of inadequacy can be dealt with by not evaluating a complete sentence with respect to all variable assignments, but only to a subset thereof. In the light of the arguments given above, this subset is rather small. So the contextually determined variables are not so variable at all; they behave more like constants.

4. A rather fundamental argument against the use of variables for formalizing contextual influence is the following. In PTQ the contextual factor of the reference point under consideration (a time world pair), is formalized by means of the so called indices I and J. Several authors have proposed to incorporate other factors in the indices. LEWIS (1970), for instance, mentions as possible indices: speaker, audience, segment of surrounding discourse, and things capable of being pointed at. These indices constitute an obvious way to formalize contextual influence. In the light of this, it is very important to realize that in IL the interpretation of constants is 'index dependent', whereas variables have an 'index independent' interpretation:

$$c^{A,i,j,g} = F(c)(i,j), \quad x^{A,i,j,g} = g(x).$$

This means that in IL it is very strange to use logical variables for the purpose of encoding contextual restrictions. The obvious method is by means

of constants. This is precisely the method employed in MONTAGUE (1968) and BENNETT (1978).

4.4. Conclusion

We considered Cooper's proposals concerning the solution of the 'not-there/left-over' problems. His idea to give a semantic treatment of the 'not-there' problem was not successfully formalized. His treatment of the variables 'left-over' led to incorrect results for English sentences. We have to conclude that the technical details of the Bach & Cooper proposal are such that their approach does not work correctly. This means that at the present stage of our investigations concerning the thematic question we are back at the situation of the end of Section 2: only the CN-S analysis seems to be possible.

I have not formally proved that it is impossible to find some treatment in accordance with Cooper's aims. As I said in Section 2, such a proof is, in general, difficult to give. But I have not only showed that the proposals by Bach & Cooper do not work correctly, I have also argued that they have to be considered as unsound. They constitute a very unnatural approach, and in my opinion one should not try to correct the proposals, but rather give up the idea underlying them altogether. Since I consider such proposals as unsound, I will in the next section put forward a principle which prohibits proposals of these kinds.

5. THE VARIABLE PRINCIPLE

In the previous section we have considered some attempts to deal with the 'not-there/left-over' problems. These attempts do not give me the impression that the considered situations they deal with are welcome; rather they seem to be escapes from situations one would prefer not to encounter at all. In my opinion these attempts arise from a neglect of the special character of syntactic variables. Syntactic variables differ from other words in the lexicon since they are introduced for a special purpose: viz. to deal with coreferentiality and scope. In this respect they are like logical variables, and in fact they can be considered as their syntactic counterpart. One would like to encounter syntactic variables only if they are used for such purposes. This special character of syntactic variables is expressed by the variable principle, of which a first tentative version is given in (29).

(29) *Syntactic variables correspond closely to logical variables.*

The intuition behind this statement is not completely new. THOMASON (1976) draws attention to the analogy between 'that-complement' constructions in English, and the λ -abstraction operator in logic. PARTEE (1979b) proposes the constraint that any syntactic variable must be translated into an expression of the logic containing an unbound logical variable. Partee does not accept this constraint the other way around, precisely because she does not want to disallow Cooper's treatment of free variables.

The formulation of the principle given in (29) is vague, and one might be tempted to strengthen it to (30).

(30) *An expression contains a syntactic variable if and only if its unreduced translation contains a corresponding unbound logical variable.*

This is intuitively an attractive formulation. However, a major drawback is that it does not fit into the framework of Montague grammar. It would give the unreduced translation of an expression a special status which it does not have in the framework as it is. The unreduced translation, would no longer be just one representation among others, all freely interchangeable. It would become an essential stage since the principle would have to function as a filter on it. It would no longer be allowed to reduce the intermediate steps in the translation process since then a semantically irrelevant occurrence of a logical variable might disappear, and thereby a translation that had to be rejected, might become acceptable. Therefore, I will give a formulation which turns the principle into a restriction on possible Montague grammars. The formulation below has the same consequences for the unreduced translation as (30), but it is not a filter on the unreduced translations and it leaves the framework untouched. This formulation is slightly more restrictive than (30), and than the formulation in JANSSEN (1980b).

The VARIABLE PRINCIPLE is defined as consisting of the following 6 requirements:

- 1a) *A syntactic variable translates into an expression which contains a free occurrence of a logical variable, and which does not contain occurrences of constants.*
- 1b) *This is the only way to introduce a free occurrence of a logical variable.*

- 2a) *If a syntactic rule removes all occurrences of a certain syntactic variable in one of its arguments, then the corresponding translation rule binds all occurrences of the corresponding logical variable in the translation of that argument.*
- 2b) *If a translation rule places one of its arguments within the scope of a binder for a certain variable, then its corresponding syntactic rule removes all the occurrences of the corresponding syntactic variable from the syntactic counterpart of that argument.*
- 3a) *The production of a sentence is only considered as completed if each syntactic variable has been removed by some syntactic rule.*
- 3b) *If a syntactic rule is used which contains instructions which have the effect of removing all occurrences of a certain variable from one of its arguments, then there indeed have to be such occurrences.*

This formulation of the variable principle is not what I would like to call 'simple and elegant'. I hope that such a formulation will be possible when the algebraic theory of the organization of the syntax is further developed. Suppose that we have found which basic operations on strings are required in the syntax (following PARTEE (1979a,b, see chapter 8)), and that a syntactic rule can be described as a polynomial over these basic operations. Then we may hope to formulate the variable principle as a restriction on the relation between the syntactic and semantic polynomials. We might then require that these polynomials are isomorphic with respect to operations removing/binding variables.

Requirement 1a) is a restriction on the translation of lexical elements. It can easily be checked whether a given grammar satisfies the requirement. It is met by all proposals in the field of Montague grammar that I know of; e.g. the PTQ translation of he_n is $\lambda P[{}^V P(x_n)]$, and the translation of the common noun variable one_n (HAUSSER 1979c) is the variable P_n .

For reasons of elegance, one might like to have formulation 1a') instead of formulation 1a).

1a') *A syntactic variable translates into a logical variable.*

In order to meet 1a') in the PTQ fragment, one could introduce a category of Proper Names containing *John, Mary, he₁, he₂, ...* (with translations *john, mary, x₁, x₂*, respectively). Out of these Proper Names, Terms could be produced which obtain the standard translation ($\lambda P[{}^V P(john)]$, etc.). Since I do not know of a phenomenon, the treatment of which would be simplified using this approach, and since the variable principle then still

would not have a simple formulation anyhow, I will not use it here. Requirement 1a) has as a consequence that the translation of a syntactic variable is logically *equivalent* to a logical variable. If constants are allowed to occur, then this would no longer be true (e.g. it is not true that for every c the formula $\exists x[x=c]$ is valid).

Requirement 1b) is a restriction both on the translation of lexical elements, and on the translation rules. This requirement is met by PTQ. It is not met by the proposals of BACH & COOPER (1978) which allow free variables to occur which do not have a syntactic counterpart. Since they do not present explicit rules, I do not know at which stage the context variable R is introduced, as a lexical ambiguity of the noun, or by means of some syntactic rule.

Requirements 2a) and 2b) are conditions on the possible combinations of a syntactic rule with a translation rule. Whether a grammar actually meets them is easily checked by inspection (PTQ does). Requirement 2b) is not met by the Bach & Cooper proposal since their approach in some cases gives rise to the introduction and binding of logical variables without any visible syntactic effect.

Requirements 3a) and 3b) we have already mentioned in chapter 6. They are not met by PTQ, nor by Bach & Cooper. In a certain sense they constitute the kernel of the principle. They express that certain configurations (described with respect to occurrences of variables) should not arise. When these requirements are met, the fundamental problems described in Section 1 disappear. As such, the two requirements are closely related to two instructions in JANSSEN (1980a, p.366), and to two conventions in RODMAN (1976, one mentioned there on p.176, and one implicitly used on p.170). Requirements 3a) and 3b) alone, i.e. without 1) and 2), would suffice to eliminate the syntactic side of the two fundamental problems, but then the close relationship between syntactic and logical variables would not be enforced. That freedom would give us the possibility to abuse syntactic variables for other purposes than coreferentiality and scope. An extreme case is given in JANSSEN (1980b), where some rules which obey 3a) and 3b), but violate 1) and 2), are defined in such a way that the information that a rule is obligatory is encoded in the syntactic variables. I intend to prohibit this and other kinds of abuse of variables by combining the third requirement with the first and second. In chapter 6, section 5.3, it is discussed how we might incorporate requirements 3a) and 3b) by filters

and partial rules or by total rules (using a refined system of categories) For the present discussion it is irrelevant how these requirements are exactly incorporated in the system. Since we are primarily interested in the effects of the principle, it suffices to know that it can be done in some way.

Let me emphasize that the principle is intended to apply to the standard variables of intensional logic and their corresponding syntactic variables. For instance, the argument concerning the use of unbound variables for contextual influence does not apply if we do not translate into IL but into Ty2. If Ty2 is used, the variable principle does not simply apply to all the variables of type *s*. Neither does the principle apply to so called 'context variables' of HAUSSER (1979c), or the 'context expressions' of GROENENDIJK & STOKHOF (1979), which both are added to IL for the special purpose of dealing with contextual influence.

The principle eliminates the basic problems from section 2 and disallows the treatment of variables aimed at in COOPER (1975), and COOPER (1979a,b). Another example of a treatment which is disallowed is the proposal of OH (1977). For a sentence without discourse or deictic pronouns he gives a translation containing a unbound variable! A consequence of the principle is that the denotation of a sentence is determined completely by the choice of the model and the index with respect to which we determine its denotation. In other words, the denotation is completely determined by the choice of the set of basic entities, the meaning postulates, the index, and the interpretation function for constants (i.e. the interpretations of the lexical elements in the sentence). In determining the denotation the non-linguistic aspect of an assignment to logical variables plays no role. This I consider to be an attractive aspect of the principle. What the impact of the principle is for the answer on the thematic question will be investigated in the next section.

6. MANY ANALYSES

6.1. The CN-S analysis for English

Do the rules for the CN-S analysis of relative clauses obey the variable principle?

Recall the PTQ rules from Section 2.1.

$S_{3,n}$ CN \times S \rightarrow CN

$F_{3,n}$ Replace he_n in β by *he/she/it* and him_n by *him/her/it*, according to the gender of the first CN in α ; concatenate (α , *such that*, β).

$T_{3,n}$ (PTQ) $\lambda x_n [\alpha'(x_n) \wedge \beta']$.

This combination of $S_{3,n}$ and $T_{3,n}$ does not obey the variable principle since possible occurrences of x_n in α' are, by λx_n , bound in the translation, whereas the occurrences of the corresponding syntactic variable he_n in α are *not* removed. This aspect is the source of the 'collision of variables' mentioned in Section 3.1. (for details see section 3.4 of chapter 5). A reformulation of $T_{3,n}$ which avoids such a collision is given by THOMASON (1974, p.261).

$T_{3,n}$ (THOMASON)

$\lambda x_m [\alpha'(x_m) \wedge \tilde{\beta}']$

where $\tilde{\beta}'$ is the result of replacing all occurrences of x_n in β' by occurrences of x_m , where m is the least even number such that x_m has no occurrences in either α' or β' .

The syntactic rule $S_{3,n}$ removes the occurrences of he_n in β . Thomason's reformulation has the effect that the unbound logical variables x_n in β' do not occur free in the translation of the whole construction, whereas the same variables in α remain unbound. Nevertheless, Thomason's reformulation does not obey the variable principle since in the syntax occurrences of he_n in β are removed, whereas in the translation the occurrences of the corresponding variable (i.e. x_n) are not bound, but of a variable x_m (where $n \neq m$).

Another kind of objection against Thomason's rule is that it is not a polynomial over IL. This objection was considered in chapter 5, section 3.4. The formulation proposed there for the translation rule is the following.

$T_{3,n}$ $\lambda P[\lambda x_n [{}^V P(x_n) \wedge \beta'](\wedge \alpha')]$.

This formulation has as a consequence that only those occurrences of x_n are bound, of which the syntactic counterparts are removed in $S_{3,n}$.

6.2. The S-S analysis for Hittite

Is an analysis of Hittite relative clause constructions possible which on the one hand satisfies the variable principle, and on the other hand produces such a construction out of two sentences?

Below I will describe an analysis which shows that the answer is affirmative. I will only deal with the example discussed in Section 3, and not with all other cases of Hittite relative clauses which are treated by COOPER (1975). My analysis is intended mainly as an illustration of the kinds of technique which are available if one obeys the variable principle.

The treatment described in Section 3 violates the variable principle because both subsentences in Figure 4 have a translation which contains an unbound variable, whereas the sentences themselves do not contain a syntactic variable. Given the principle, in both sentences there has to be an occurrence of a syntactic variable as well. The English variant of sentence S_2 gives a hint on how to do this. It contains in a CN-position the word (*one*) – probably added for explanatory reasons. This word suggests the introduction in the syntax of CN variables one_1, one_2, \dots , which are translated into logical variables P_1, P_2, \dots , respectively (such CN-variables are discussed in HAUSSER (1979c)). The rule which combines S_1 with S_2 will then give rise to a translation in which (by λ -conversion) the relevant property is substituted for P_n . In case one prefers not to introduce a new constituent one_n , a new variable of category T might be introduced alternatively: (31), translating as (32).

(31) $each_n$

(32) $\lambda_Q[\forall x[\overset{V}{P}_n(x) \rightarrow \overset{V}{Q}(x)]]$.

The variable in the translation of the relative clause can be introduced by the translation of the determiner *wh*. Therefore, the category of determiners (which contains the Hittite version of *every*, etc.) is extended with a variable (33), translating as (34).

(33) wh_n

(34) $\lambda_Q\lambda P[\overset{V}{Q}(z_n) \wedge \overset{V}{P}(z_n)]$.

We have to combine a relative clause containing a free variable z_n with a main sentence containing a free variable P_n . This can be done by means of a single rule binding both logical variables and performing the relevant operations on both syntactic variables, or by means of two rules, each dealing with one variable at a time. The former method would yield the tree from figure 4, but it would implicate that a new kind of rules is introduced (rules with two indices). I will follow the two-rules approach.

First the relative clause is transformed into an expression of the new

category Prop (=t//e), being a set of expressions denoting properties.
 We do this by means of the following rule (the numbers in the 800-series
 are numbers of newly proposed rules).

$S_{801,n}$ $S \rightarrow \text{Prop}$

$F_{801,n}$ Replace wh_n in α by wh

$T_{801,n}$ $\lambda z_n[\alpha']$.

The rule combining a property with a sentence is

$S_{802,n}$ $\text{Prop} \times S \rightarrow S$

$F_{802,n}$ delete all occurrences of one_n from β ;
 concatenate (α, β)

$T_{802,n}$ $[\lambda P_n \beta'](\wedge \alpha')$.

Using these rules, the Bach & Cooper example is obtained in the way
 indicated in figure 7. Its translation is equivalent to the one given in
 Section 3 for figure 4. Since we assume that it is guaranteed that the vari-
 able principle is obeyed, no problems arise with the syntactic variables.
 The principle guarantees that rule $S_{802,1}$ is applied only in case the main
 sentence contains an occurrence of one_1 and that rule $S_{801,2}$ is applied
 only when the sentence contains an occurrence of the variable wh_2 . Further-
 more, it guarantees that all syntactic variables finally will have disap-
 peared.




Figure 7

6.3. The T-S analysis for English

As shown in Section 6.2, an S-S analysis can be obtained simply by
 introducing a variable in the syntax, when such a variable is required in

the translation. The same idea can be used to obtain a T-S analysis for relative clauses. In this case, we need a variable of the category Prop, written as *of-kind_n*. It translates into the variable K_n .

A property and a common noun phrase combine to a new common noun phrase as follows:

S₈₀₃ CN × Prop → CN

F₈₀₃ concatenate (α, β)

T₈₀₃ λy[α'(y) ∧ β'(y)].

A category RC of relative clauses (RC = t//e) is introduced because RC's and Prop's will occur in different positions. The expressions of the category RC are made out of sentences as follows:

S_{804,n} S → RC

F_{804,n} delete the index *n* from all pronouns in α;
concatenate (*such that*, α)

T_{804,n} λx_n[α'].

A relative clause may be quantified into a term phrase by substituting the relative clause for a property variable:

S_{805,n} T × RC → T

F_{805,n} substitute β for *of-kind_n* in α

T_{805,n} λK_n[α']([^]β').

An example of a production using these rules is given in figure 8.




Figure 8

The translation of the lower term phrase in figure 8 is (35), the translation of the RC phrase (36), and of the upper term phrase (after

reduction) is (37).

$$(35) \lambda Q \forall x [boy(x) \wedge \forall K_3(x) \rightarrow \forall Q(x)]$$

$$(36) \lambda x_2 [run(x_2)]$$

$$(37) \lambda Q \forall x [boy(x) \wedge run(x) \rightarrow \forall Q(x)].$$

Note that the intermediate stage of an RC is not required if S_{805} is a double indexed rule, dealing both with he_n and $of-kind_m$.

6.4. The Det-S analysis for English

Is a Det-S analysis possible which obeys the variable principle? Recalling the pattern underlying the S-S and T-S analyses, one might try to find such an analysis as a variant of the CN-S analysis by introducing new variables. It appeared, to my surprise, that it is possible to obtain a Det-S analysis which is not a variant of the CN-S analysis, but which is a pure Det-S analysis (recall the proviso by Partee for her argumentation concerning the Det-S analysis). I will not discuss the heuristics of this analysis, but present the rules immediately.

$$S_{806,n} \quad Det \times S \rightarrow Det$$

$$F_{806,n} \quad \text{remove all indices } n \text{ from pronouns in } \beta;$$

$$\text{concatenate } (\alpha, \text{such that}, \beta)$$

$$T_{806,n} \quad \lambda R [\alpha' (\wedge \lambda y [\forall R(y) \wedge \lambda x_n [\beta'](y)])].$$

Maybe the following explanation of the translation is useful. A determiner δ is, semantically, a function which takes as argument the property η expressed by a noun and delivers a collection of properties which have a certain relation with η . S_{806} produces a determiner which takes a noun property η and delivers a set of properties which has that relation with the conjunction of η and the property expressed by the relative clause.

The combination of a CN with a Det-phrase, requires that the CN is placed at a suitable position in the determiner phrase. In the present fragment this position is the second position (if we had determiners like *all the*, then also other positions might under certain circumstances be suitable). The rule for this reads as follows:

$$S_{807} \quad Det \times CN \rightarrow CN$$

$$F_{807} \quad \text{insert } \beta \text{ after the first word of } \alpha$$

$$T_{807} \quad \alpha' (\wedge \beta').$$

The combination of the determiner *every* with the sentence *he₂ runs* yields determiner (38), with (39) as unreduced, and (40) as reduced translation.

(38) *every such that he₂ runs*

(39) $\lambda R[\lambda Q\lambda P[\forall x[{}^V Q(x) \rightarrow {}^V P(x)]](\lambda y[{}^V R(y) \wedge \lambda x_2[run(x_2)](y)])]$

(40) $\lambda R\lambda P\forall x[{}^V R(x) \wedge run(x) \rightarrow {}^V P(x)]$.

The combination of (38) the common noun *man* yields the term phrase (41), which has the (usual) reduced translation (42).

(41) *every man such that he runs*

(42) $\lambda P\forall x[man(x) \wedge run(x) \rightarrow {}^V P(x)]$.

The techniques which are used to obtain a T-S analysis from a CN-S analysis can be used as well to obtain a T-S analysis which is a variant of the Det-S analysis: introduce in the Det-S analysis the variable *of-kind_n*, but now within the determiner. This means that at least two kinds of T-S analyses are available.

6.5. Conclusion

In Section 5 a new principle was introduced: the variable principle. Obeying this principle we designed rules for relative clause constructions. It turned out that for English besides the CN-S analysis both the T-S and the Det-S analysis are possible in at least two essentially different variants. And for Hittite an S-S analysis is possible. So at the present stage of our investigations a negative answer to the thematic question has to be given: several analyses of relative clauses are possible.

Consider the T-S analysis of 6.3 again. Is it the kind of T-S analysis meant by Partee? I do not think so. At a certain level we indeed have a T-S analysis, but on another level in the production tree there is a CN-Prop analysis which is nothing but a variant of the CN-S analysis. The opposition between the two analyses was, however, the main point in the discussion of PARTEE (1973). So one could say that her conclusion that the pure T-S analysis cannot be used, in a certain sense still holds. For the case of Hittite however, the discussion primarily aimed at obtaining an S-S analysis at some level, rather than at avoiding the CN-S analysis on all levels. In Section 2 I quoted Bach & Cooper who expressed the hope for the 'happy discovery of yet unknown principles' which exclude the

T-S-analysis, but allow for the S-S-analysis. It seems reasonable to interpret this as the desire for a principle which prohibits the pure T-S analysis, but allows some variant of the S-S analysis. The variable principle has such an effect. But if it is interpreted as the hope for a principle which excludes all kinds of T-S analyses, or which allows a pure S-S analysis, then the variable principle is not such a principle. So the answer to the thematic question I gave above, has to be relativized: although *several* analyses are available, not *all* analyses are possible.

The answer to the thematic question obtained in this section, was based upon an investigation of the relative clause construction as such. Interaction with other phenomena was not taken into consideration. In the next section I will leave this isolation and consider the interaction of relative clause formation with some other phenomena.

7. OTHER ARGUMENTS

7.1. Syntax: gender agreement

The relative pronoun has to agree in gender with the antecedent noun-phrase. In the Det-S analysis, this poses a problem. The rule which combines a determiner with a relative clause has to specify what is to be done with the syntactic variable. The formulation I gave of rule $S_{806,n}$ just deletes the index, so it gives a correct result if the noun has male gender. But in the same way as we produced *every boy such that he runs*, we may produce *every girl such that he runs*. It is not possible to formulate S_{806} in such a way that this kind of ill-formedness is avoided, because the information which gender the noun has, is not available at the stage at which the determiner and the relative clause are combined. Not removing the index would, according to the variable principle, require a free variable in the translation of the term phrase; but I do not see how this approach might work.

The T-S analysis gives rise to a similar problem. The rule which makes the relative clause (RC) out of a sentence (S), has to specify what has to be done with he_n . The formulation I gave of S_{804} works correctly for masculine nouns only. Again, information about the gender of the noun is not yet available, and not removing the index would constitute a break with the principle. This argument does not apply to the T-S analysis in which a double indexed rule is used. In the CN-S analysis, no problems arise from

gender agreement, since at the stage at which the index has to be removed, the gender of the noun is known.

One should not conclude from this discussion that it is impossible to obtain correct gender agreement in case of the Det-S or T-S analysis under discussion. I expect that it can be done by means of further subcategorization. One has to distinguish feminine, masculine, and neuter relative clauses, and feminine, masculine, and neuter determiners, and probably one needs to make similar distinctions in other categories. Then the subcategory system provides the information needed to obtain precisely the correct combinations of relative clause, determiner and noun.

There is the hidden assumption in this discussion that gender agreement has to be handled within the syntax. If we do not assume this, then a phrase as *a girl such that he runs*, is no longer considered to be syntactically ill-formed. COOPER (1975) argues in favor of dealing with gender in the semantics (at least for English). Others might prefer to handle gender in pragmatics (Karttunen, according to PARTEE (1979a)). Then the arguments given here are no longer relevant. But in languages with grammatical gender (e.g. Dutch, German), this escape is not available. Here one might adopt one of the solutions I mentioned: refined subcategorization, a T-S analysis with a double indexed rule, or simply the CN-S analysis for relative clauses.

7.2. Semantics: scope

Consider the following sentence (exhibiting stacking on the head *man*):

(43) *Every man such that he loves a girl such that he kisses her is happy.*

This sentence has a possible reading in which *every* has wider scope than *a*. In a PTQ like approach (so with the CN-S construction for relative clauses), this reading is obtained by quantification of *a girl* into the CN phrase

(44) *man such that he loves him_n such that he kisses him_n.*

The corresponding translation of the sentence (44) reduces to

(45) $\forall y[\exists x[\text{girl}(x) \wedge \text{man}(y) \wedge \text{love}_x^V(y, x) \wedge \text{kiss}_x^V(y, x)] \rightarrow \text{happy}(y)].$

Can this reading be obtained in other analyses of relative clauses?

In the T-S analysis this stacking of relative clauses can be obtained by means of a process indicated in figure 9. In order to obtain coreferentiality between both occurrences of the term *him_n*, the term *a girl* has

to be substituted at a stage in which both relative clauses are present. The earliest moment at which this is the case, is immediately after the uppermost term has been formed. Using a rule analogous to the standard quantification rules would assign the existential quantifier wider scope than the universal quantifier, thus not yielding the desired reading. So it seems to be impossible to obtain in such a T-S analysis coreferentiality and correct scope at the same time.




Figure 9

In the Det-S analysis the earliest stage at which the coreferentiality of *she* and *a girl* can be accounted for, is when the determiner phrase (46) has been formed.

(46) *every such that he loves him₃ such that he kisses him₃.*

Some later stage (e.g. the term level), might be selected as well. But in all these options, the quantification rule would give wider scope to *a* than to *every*, thus not yielding the desired reading.

Underlying this discussion is the assumption that there is something like stacking of relative clauses. If there is stacking, then the rule for quantification into a CN is essential for the PTQ fragment (FRIEDMAN & WARREN (1979b)). But is stacking indeed a phenomenon of natural language? As for Hittite, BACH & COOPER (1975) inform us that no stacking occurs, and in Dutch and German stacking is not possible. As for English, no author expresses doubts, except for PARTEE (1979b). She states that the evidence for stacking is spurious. If we accept this, it would leave a rather small basis for our argumentation concerning an answer on the thematic question.

There is another phenomenon, however, that requires quantification

into CN's. It might be the kind of examples meant by PARTEE (1975, p.236). Example (47) assumes that there are common nouns in the fragment of the form *friend of*.

(47) *Every picture of a woman which is owned by a man who loves her is a valuable object.*

Here the intended reading is the one in which *every* has wider scope than *a*, and in which there is coreferentiality between *a woman* and *her*. This reading can easily be obtained by means of substitution of *a woman* into the CN-phrase (48).

(48) *picture of he₁ such that it is owned by a man such that he loves him₁.*

So even if we do not accept stacking as a phenomenon of English, a CN-S analysis would be required.

It is remarkable to notice that the variable principle plays no role in the discussion concerning scope. The occurrences of the Prop variables, which form a practical consequence of the principle, were not relevant. If they were omitted, which would bring us back to the original Bach & Cooper approach, then still the same problems would arise with respect to scope. So even without the variable principle a CN-S analysis appears to be required. This conclusion has to be relativized immediately. I have not given a formal proof that it is impossible to obtain a correct treatment of scope in the other analyses. I just showed that the CN-S analysis provides a direct basis for a semantic treatment of scope phenomena in a way that the considered T-S and Det-S analyses can not. This conclusion mentions another argument for relativizing. We only considered the three analyses which had our main interest. A lot more analyses are possible, and for some a correct treatment of scope may be possible. For instance, a correct treatment of scope might be possible if the category of determiners contains variables for which a determiner can be substituted in a later stage.

7.3. Conclusion

In the previous section we observed that the framework of Montague grammar hardly restricts the possible syntactic analyses of relative clauses. In this section we investigated the possibilities for incorporating the available options in a somewhat larger fragment. It turned out that from the three main options only one was suitable. From this we learn that it is important to consider phenomena not only in isolation, but to design

grammars for larger fragments. The fact that for each isolated phenomenon there are many syntactic options available, gives us a firm basis for the hope that it is indeed possible to find a combination of syntactic constructions that fits together in a system yielding the correct semantics for the constructions involved. Thus we see that extending fragments is a fruitful step which has impact on the description of isolated phenomena. This can be considered as a reaction to be a remark Van BENTHEM (1981, p.31) who denies the use of generalization and the combination of partial theories.

8. THE GENERAL QUESTION

In this section I answer the general version of the thematic question. We employ a framework in which the syntax and semantics have to be algebras, and in which meaning assignment is a homomorphism. The general version of the thematic question was to what extent this organization of the grammar restricts the options we have available for describing a particular phenomenon in the syntax.

For the special case of relative clause formation we obtained in section 6 the answer that any kind of analysis can be obtained, but that certain kinds of analysis cannot be avoided. This practical result will be explained below on the basis of the algebraic properties of the framework, and the result will be generalized to an answer on the general question.

Let us suppose that we have found a semantic operation T_{888} which takes two arguments, and delivers the meaning of a certain construction. So in the semantics we have the construction step $T_{888}(\alpha', \beta')$. Due to the homomorphism relation, there has to be a corresponding operation $F_{888}(\alpha, \beta)$ in the syntax, and the two semantic arguments have to correspond with the two syntactic arguments. Instead of the semantic step $T_{888}(\alpha', \beta')$, several variants are possible, each with its own consequences for the syntax. These variants amount to a construction process with two stages. We may first have $T_{888}(\alpha', R)$, where R is a variable, and introduce in a later stage a λ -operator for R taking β' as argument:

$$\lambda R[\dots T_{888}(\alpha', R)\dots](\beta').$$

This means that the syntactic expression β can be introduced in an arbitrary later stage of the syntactic production process. Consequently, a lot of variants of the original syntactic construction can be formed. These variants

are based on the use of the construction step $T_{888}(\alpha', R)$ in the logic. Due to the variable principle, the variable R has to be introduced by the translation of some syntactic variable. Let us suppose that V is such a variable. Due to the homomorphic relation between syntax and semantics, this means that in the syntax there has to be a step $F_{888}(\alpha, V)$. So whereas we have gained the freedom to introduce β in a later stage of the syntactic construction process, step F_{888} is not avoided. The same argumentation applies when the first argument of T_{888} is replaced by a variable. It is even possible to replace both arguments by a variable, thus obtaining a large freedom in the syntax concerning the stage at which α and β are introduced. But in all these variants F_{888} is not avoided. Application of this argumentation to the case of relative clauses (where two basic constructions are found) means that we cannot avoid both the CN-S and the Det-S construction at the same time.

So on the basis of the compositionality principle, formalized in an algebraic way, many relative clause constructions are possible. This is due to the power of λ -abstraction. This operation makes it possible that on the semantic side the effect is obtained of substituting the translation of one argument on a suitable position within the other argument, whereas in the syntax a completely different operation is performed. Referring to this power Partee once said 'Lambdas really changed my life' (Lecture for the Dutch Association for Logic, Amsterdam, 1980).

The above argumentation is not completely compelling: there is (at least) one exception to the claim that it is not possible to make a variant of a given semantic construction which avoids the corresponding syntactic construction step. An example of such an exception arose in the S-S analysis for Hittite. In the main sentence we had the Det-CN construction *each one_n*, where *one_n* was a variable. We obtained a variant in which there is no Det-CN construction: the logical variable introduced by *one_n*, could be introduced by a new variable *each_n* (see (34)). The algebraic description of this method is as follows. Consider again $T_{888}(\alpha', R)$. The variable R might, under certain circumstances, be introduced by the translation of α , thus allowing to replace T_{888} by a related semantic operation which takes only one argument. That the translation of α introduced the variable R , means that in the syntax α is to be replaced by some variable, say an indexed variant of α . Its translation is then a compound expression (being a combination of the old translation α' with the variable R). This process,

which avoids to have F_{888} in the syntax, is possible only if α is a single word with a translation which does not contain a constant (e.g. if α is a determiner). If the translation of α would contain a constant, then requirement 1a) of the variable principle would prohibit that its translation introduces a variable. If α is not a single word, then it cannot be replaced by a syntactic variable (maybe one of its parts can then be indexed). This method of creating exceptions would be prohibited when requirement 1a) of the variable principle would be replaced by the more restrictive version 1a'). In order to prove that the exception described here is the only one by which a given analysis can be avoided, the details of the relation between operations in the semantics or in the syntax have to be formalized algebraically (see also Section 3).


These algebraic considerations explain the results of our practical work. On the basis of these considerations it would be possible to explain that a Det-S analysis which is variant of the CN-S analysis, is not to be expected (in any case the described method for obtaining variants does not work). The algebraic considerations also answer the general question whether the principle of compositionality restricts the options available for descriptive work. On the basis of a given construction step, a lot of variants are possible, but due to the variable principle and the homomorphic relation between syntax and semantics, this construction step cannot be avoided in these variants. So the answer to the general question is that there are indeed restrictions on the syntactic possibilities, but only in the sense that a basic step cannot, generally speaking, be avoided. But these restrictions are not that strong that only a single analysis is possible. Formal proofs for these considerations would require, as I said before, a further algebraization of the syntax.

CHAPTER X

SCOPE AMBIGUITIES OF TENSE, ASPECT AND NEGATION

ABSTRACT

In this chapter verbal constructions with *will*, *have*, with negation, and with the past tense are considered. The meanings of these syntactic constructions are expressed by semantic operators. These operators have a certain scope, and differences in scope give rise to semantic ambiguities. These scope ambiguities are investigated, and a grammar dealing with these phenomena is presented. In this grammar features and queries are used, and the grammar produces labeled bracketings.



1. INTRODUCTION

Verbal constructions with *will*, *have*, with negation, or with the past tense, give rise to semantic operators: negation, tense operators and aspect operators. The syntactic counterparts of such operators I will call 'verb-modifiers'. So a basic verb modifier consists sometimes of a single word (*will*, *have*), sometimes of two words (*do not*), and sometimes of a verb affix (for the past). Compound verb modifiers are combinations of basic modifiers; they may consist of several words (*will not have*).

The semantic operators which correspond with basic verb modifiers have a certain scope, and a sentence can be ambiguous with respect to the scope of such an operator. The aim of the present chapter is to present a treatment of scope phenomena involving terms and verb modifiers. Examples of such ambiguities are provided by sentences (1) and (2). Both are ambiguous; each sentence may concern either the present president or the future president.

- (1) *The president will talk*
 (2) *John will hate the president.*

It is not my aim to analyse in detail the semantic interpretation of operators corresponding with verb modifiers. I will not present proposals for the formal semantics of tense or aspect; there is, in my treatment, no semantic difference between past and perfect (there is a syntactic difference). It is my aim to investigate only scope phenomena of operators and not to consider the operators themselves.

The main conclusion concerning the treatment of scope will be that the introduction of verb modifiers has to be possible both on the level of verb phrases and on the level of sentences. Another conclusion will be that compound verb modifiers have to be introduced in a step by step process: each step introducing one semantical operator. The treatment that I will present does not deal with all scope phenomena correctly (see section 5).

2. THE PTQ APPROACH

2.1. Introduction

As starting point for my investigations I take the treatment of verb modifiers as presented in MONTAGUE 1973 (henceforth PTQ). I will discuss F. Heny and B Richards (eds.), *Linguistic Categories: Auxiliaries and Related Puzzles*, Vol. Two, 55-76.

Copyright © 1983 by D. Reidel Publishing Company.

syntactic and semantic aspects of that proposal and compare these with related aspects of my approach.

2.2. Syntax of PTQ

The grammar of PTQ has six operations for the treatment of verb modifiers: rules for present, perfect and future, and for the negated variants of these tenses. Some examples:

F₁₄ (*John, walk*) = *John has walked*

F₁₅ (*John, walk*) = *John has not walked*

F₁₁ (*John, walk*) = *John does not walk.*

These operations are completely independent. The operation 'make a sentence in the perfect tense' is independent of the operation 'make a sentence in the negative perfect tense'. One would like to have here another situation. My treatment aims at a so-called 'orthogonal' syntax: each phenomenon will be treated by its own collection of rules (e.g. 'negating' will be treated by means of a rules which just deal with negation), and all such collections of rules will have the same structure as much as possible.

The PTQ rules do not treat conjoined verb phrases correctly since only the first verb is conjugated. So the PTQ syntax produces (3) instead of (4).

(3) *John has walked and talk*

(4) *John has walked and talked*

FRIEDMAN (1979) has given a treatment of this kind of error, and the treatment in this chapter of these problems will be about the same as hers.

The rules of PTQ deal with only three verb-modifiers: future, perfect and negation. Compound modifiers such as past perfect (in *had walked*) are not treated, nor the simple past (*walked*). These modifiers will be incorporated in the fragment of the present chapter. Furthermore, compound verb phrases will be incorporated of which the conjuncts (disjuncts) may be modified in different ways (*has walked and will talk*).

2.3. Ambiguities

The grammar of PTQ deals with several scope ambiguities. I will recall two of them because variants of them will return in the discussion. The most famous example is (5). This sentence has a de-re reading (6) and a de-dicto reading (7).

(5) *John seeks a unicorn*

(6) $\exists u[\text{unicorn}_*(u) \wedge \text{seek}_*(\text{john}, u)]$

(7) $\text{seek}(\wedge \text{john}, \wedge \lambda P \exists u[\text{unicorn}_*(u) \wedge \vee P(u)])$.

Another example is the scope ambiguity in (8); this sentence has readings (9) and (10)

(8) *Every man loves a woman*

(9) $\forall u[\text{man}_*(u) \rightarrow \exists v[\text{woman}_*(v) \wedge \text{love}_*(u, v)]]$

(10) $\exists v[\text{woman}_*(v) \wedge \forall u[\text{man}_*(u) \rightarrow \text{love}_*(u, v)]]$.

The readings of (8) have a remarkable property. Reading (10) logically implies (9). This means that there is no situation in which (10) is true, while (9) is false. For this reason one might doubt whether this scope ambiguity is an ambiguity we have to account for: reading (9) seems to be always acceptable. I will give two arguments explaining why (8) is considered ambiguous. Both arguments are due to Martin Stokhof (personal communication); see also chapter 4, section 6.

The first argument is that for slight variants of (8) the two readings are logically independent. Consider sentence (11), in which we understand *one* as *precisely one*.

(11) *Every man loves one woman.*

This sentence has readings (12) and (13), where neither (12) follows from (13), nor (13) from (12).

(12) $\forall u[\text{man}_*(u) \rightarrow \exists! v \text{woman}_*(v) \wedge \text{love}_*(u, v)]$

(13) $\exists! v[\text{woman}_*(v) \wedge \forall u[\text{man}_*(u) \rightarrow \text{love}_*(u, v)]]$.

A more well-known variant of the scope ambiguities of (8) and (11) is sentence (14). Also here the two readings are independent.

(14) *Every man in this room speaks two languages.*

These considerations show that sentences closely resembling (8) exhibit independent ambiguities.

The second argument is that in certain contexts the weaker reading of (8) is required. Consider (15) or (16).

(15) *It is not the case that every man loves a woman*

(16) *John does not believe that every man loves a woman.*

Sentence (15) can be used in situations in which it means (17), as well as in situations where it means (18).

(17) $\neg [\forall u[man_*(u) \rightarrow \exists v[woman_*(v) \wedge love_*(u,v)]]]$

(18) $\neg \exists v[woman_*(v) \wedge \forall u[man_*(u) \rightarrow love_*(u,v)]]$.

Here the implication goes in the other direction: reading (17) implies (18). So if we preferred to have only one reading for (15), it would have to be (18). It is very likely that (15) is obtained by building (8), and next negating it. This construction requires that reading (18) of (15) be produced from reading (10) of (8). So sentences like (15) require that reading (10) is available. Hence (8) should get both reading (9) and (10).

2.4. Model

In several recent proposals arguments are put forward in favor of another model for time than the one used in PTQ. Such proposals are based upon a model with an interval semantics for time, rather than one with time point semantics (DOWTY 1979b, many contributions in ROHRER 1980). I will not incorporate these innovations, but follow the PTQ logic and semantics since it was not my aim to improve the PTQ interpretation of modifiers. This means that the logic does not provide for the tools to discriminate semantically between simple past and perfect, and therefore I will assign the same meanings to them. The use of the PTQ model has as a consequence that, formally spoken, I only deal with a limited use of tense: the reportive use (see BENNETT 1977).

Using such a 'primitive' semantics is, for the present purposes, not a great drawback. The scope phenomena under discussion will arise within any semantic treatment of tenses, no matter what kind of a model is used. I expect that my treatment can be adopted for another model (by taking the same derivational history, but changing the translations or their interpretations).

3. BASIC VERB MODIFIERS

In this section sentences will be considered which contain basic verb modifiers. First such sentences will be considered from a syntactic point of view. The PTQ rules produce such modifiers in few contexts only, but there are more situations in which they may occur. Next we will consider

such sentences from a semantic point of view and investigate their scope ambiguities. Finally we will consider which kind of rules might be used to produce such sentences and to obtain the desired meanings.

The first kind of situation we will consider are the complements of verbs like *try*, *assert* and *regret*. The rules of PTQ allow for unmodified verb phrases as complement. An example is (19).

(19) *John tries to run.*

PTQ does not allow for negated verbs as complement. Such complements are possible as is pointed out by BENNETT (1976); see example (20). The sentence is intended in the reading that what John tries, is not to run.

(20) *John tries not to run.*

As sentence (21) shows, complements in the perfect are also possible (unlike the PTQ predictions).

(21) *John hopes to have finished.*

Future is not possible in these complements (but in Dutch it is possible).

The second kind of situations where the PTQ rules are inadequate is provided by sentences with conjoined verb phrases. The PTQ syntax states that the first verb has to be conjugated. If we assume that the rule is changed to mark all relevant verbs, then sentences like (22) are produced.

(22) *John has walked and talked.*

In the PTQ approach it is not possible to obtain differently modified verbs in the conjuncts to the verb phrase; yet it was noticed by BACH (1980) and JANSSEN (1980b) that they can be combined freely. Some examples, due to Bach (op. cit.) are (23) and (24).

(23) *Harry left at three and is here now.*

(24) *John lives in New York and has always lived there.*

These examples can easily be adapted for other verb modifiers. In (25) negation occurs and in (26) future.

(25) *Harry left at three but is not here now.*

(26) *John has always lived in New York and will always stay there.*

So the PTQ syntax has to be extended for complements and conjuncts.

Now we come to the semantic aspect. Sentences which contain a modifier exhibit scope ambiguities with respect to the corresponding operator. An

example is (27).

(27) *The president will talk.*

This sentence has a reading which says that the present president will speak at a moment in the future (maybe after his presidency). It also has a reading which says that on a future moment the then president will speak. So sentence (27) has readings (28) and (29).

(28) $\exists u[\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge W[\textit{talk}_*(u)]]$

(29) $W\exists u[\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge \textit{talk}_*(u)].$

Notice that I consider *president* a predicate which may apply for different reference points to different persons. In some cases an index independent interpretation of an in principle index-dependent noun seems to be required. The American hostages in Iran will probably always be called *hostages* although they are no longer hostages. This means that this noun in sentence (30) is used with an index independent interpretation.

(30) *The hostages were received by the president.*

I assume that even *the president* can be used in an index independent way; in a biography about Eisenhower one might say

(31) *The president studied at West-Point.*

With an index independent interpretation of *president* formulas (28) and (29) are equivalent. An example of a term for which only an index-dependent interpretation is possible is *70-years-old-man*. Sentence (32) only has readings (33) and (34).

(32) *A 70 years old man will visit China.*

(33) $W\exists u[70\text{-years}_*(u) \wedge \textit{man}_*(u) \wedge \textit{visit}_*(u, \textit{China})]$

(34) $\exists u[70\text{-years}_*(u) \wedge \textit{man}_*(u) \wedge W[\textit{visit}_*(u, \textit{China})]].$

For past tense and for negation ambiguities arise which are related to the ambiguities for future discussed above. For perfect the opinions vary. Some native speakers have claimed that perfect can only have narrow scope, whereas others have no problems with two readings for sentence (35).

(35) *The president has talked.*

The grammar I will present, assigns two readings to (35), but a slight modification would give only one reading.

For sentences with differently modified verb phrases there is no scope ambiguity. Sentence (36) only has reading (37), see BACH 1980.

(36) *A woman has walked and will run*

(37) $\exists u[\text{woman}_*(u) \wedge H[\text{walk}_*(u)] \wedge W[\text{run}_*(u)]]$.

The above examples concerning embeddings and conjunctions suggest that it is useful to have rules which produce modified verb phrases. This is the approach that will be followed in this article. But the examples do not *prove* that it is impossible to design a system in which only sentences with verb modifiers are produced, and no modified verb phrases. I will sketch below some problematical aspects of such approaches.

One might think of introducing the perfect on the level of sentences, thus obtaining (39) from (38). Combination with (40) then yields (41).

(38) *Harry leaves at three*

(39) *Harry has left at three*

(40) *Harry is here now*

(41) *Harry has left at three and Harry is here now.*

From (41) we obtain (42) by means of a deletion rule.

(42) *Harry has left at three and is here now.*

For these sentences there arise no problems with this approach. But for

(43) it is problematic since (43) does not have the same meaning as (44).

(43) *A man left at three and is here now*

(44) *A men left at three and a man is here now.*

Our framework requires that there be a semantic operation which corresponds with the rule that produces (42) from (41) and (43) from (44). I do not know of a semantic operator which has the desired effect, and therefore it is questionable whether this approach can be followed.

A variant of this method, due to Van Benthem (personal communication) is to produce (42) from (45).

(45) *He₁ has left at three and is here now.*

Sentence (45) is produced in the way sketched above, so obtained from (46) by means of a deletion rule.

(46) *He₁ has left at three and he₁ is here now.*

The semantic problem mentioned above does not arise because (45) and (46) are equivalent. I expect that an approach like this will require rules which are far more complex than the rules which produce modified verb phrases in this chapter.

If we have rules introducing verb modifiers at the level of verb phrases do we then still need rules introducing them at the level of sentences? The answer of BACH (1980) seems to be that only rules for verb phrases are needed. He presents a new translation rule corresponding with the syntactic rule which produces a sentence from a term and a verb phrase. His translation rule has the effect that in the translation of the sentence the operator in the IV-translation gets wider scope than the subject. So the basic situation is that subjects obtain narrow scope, and subjects can obtain wide scope by quantifying in. In this way the two readings of (47) are obtained.

(47) *The president will walk.*

An exception to this pattern is the conjunction (disjunction) of differently modified verb phrases. As we observed above, the subject can only have wide scope. Recall (36)

(36) *A woman has walked and will run.*

In order to deal with such constructions, Bach presents translation rules for conjunction and disjunction of verb phrases which have the effect that for such constructions the subject gets wide scope.

Bach's approach is insufficient because there are examples where one wishes to quantify a term in, but where nevertheless this term should be within the scope of the tense operator. I will give three examples. Each exhibits in the future tense a phenomenon for which quantification rules are commonly used in the present tense. The first example concerns scope ambiguity of quantifiers: sentence (48) with reading (49).

(48) *Every catholic will follow a man*

(49) $\forall u[\text{man}_*(u) \wedge \forall v[\text{catholic}_*(v) \rightarrow \text{follow}_*(v,u)]]$.

In order to obtain reading (49) one wishes to quantify *a man* into *Every catholic follows him* and only after that, assign the tense. The second example concerns the de-dicto/de-re ambiguity: sentence (50) with reading (51).

(50) *John will seek a unicorn*

(51) $W\exists u[\text{unicorn}_*(u) \wedge \text{seek}_*(\text{john},u)]$.

Here John seeks a future 'de-re unicorn'. Again one wishes to quantify in, and then assign tense. The third example concerns coreferentiality of terms inside the scope of the tense operator: sentence (52) with reading (53).

(52) *The president will love a woman who kisses him*

(53) $W\exists u[\forall v[\text{president}_*(v) \leftrightarrow u = v] \wedge \exists w[\text{woman}_*(w) \wedge \text{kiss}_*(w,u) \wedge \text{love}_*(u,w)]]$.

This translation represents the reading in which the loving and kissing happen on the same moment in the future. Again one wishes to produce this sentence by means of first quantifying in at the sentence level, followed by tense assignment on that level. Related examples can be given for other tenses and aspects.

For the introduction of negation on the sentence level related examples can be given: situations where one wished to quantify in, but where negation has wide scope. Examples are the wide quantifier scope in (54), the de-re reading of (55) and the coreferentiality in (56).

(54) *Every woman does not love a man*

(55) *John does not seek a unicorn*

(56) *The president does not love the woman who kisses him.*

The main conclusion of this section is that rules are needed which introduce modifiers on the level of verb phrases, but that also rules are needed which do so on the level of sentences. This aspect constitutes the fundamental difference between the present approach and the approach of BACH (1980). Notice that an important part of the argumentation is based upon the fact that phenomena like scope of terms, de-dicto/de-re ambiguity and coreferentiality, are dealt with by means of quantification rules.

The last part of this section consists of two examples of sentences which are produced according to the ideas I have just sketched. The details of the rules will not be given here, but the sequence of stages of the process (and the respective translations) are the same as the ones obtained by using the rules of the grammar from section 7.

The first example is sentence (57), with reading (58).

(57) *John will seek a unicorn*

(58) $W[\exists u[\textit{unicorn}_*(u) \wedge \textit{seek}_*(\textit{john},u)]]$.

First sentence (59) is produced, it has (60) as translation.

(59) *John seeks him₁*

(60) $\textit{seek}(\wedge \textit{john}, \wedge \lambda P \wedge P(x_1))$.

The next step is to quantify in the term *a unicorn*. Then sentence (61) is obtained, with translation (62).

(61) *John seeks a unicorn*

(62) $\exists u[\textit{unicorn}_*(u) \wedge \textit{seek}_*(\textit{john},u)]$.

The last step is the introduction of future tense in (61). This gives us sentence (57), with (58) as translation.

The second example concerns the sentence *John tries not to run*. This sentence contains the verb phrase *not to run*, and this raises the question which kind of translation we use for verb phrases. BACH (1980) has given several arguments for considering verb phrases as functions operating on subject terms. This approach has as a consequence that a new, somewhat complex translation rule has to be used for S4 (the rule which combines a T and an IV to make an S). One of Bach's arguments in favor of considering verb phrases as functions was his treatment of tense and aspect. As we concluded, his proposal is in this respect not satisfactory. His other arguments in favor of verb phrases as functions, concern phenomena I do not deal with in this article (such as 'Montague phonology' and constructions like *A unicorn seems to be approaching*). Since in our fragment we will not have any of the advantages of that approach, I will use the simpler PTQ translation. But no matter which translation is chosen, the conclusion that modifiers have to be introduced on two levels remains valid.

Let us return to the example, sentence (63) with translation (64).

(63) *John tries not to run*

(64) $\textit{try to}(\wedge \textit{john}, \lambda x \neg [(\textit{run}(\vee x))])$.

The first stage in the production of this sentence is to produce verb phrase (65). Its translation as explained above, is (66).

(65) *do not run*

(66) $\lambda x \neg [(\textit{run}_*(\vee x))]$.

The next step is the addition of *try to*, yielding (67), with (68) as translation.

(67) *try not to run*

(68) $try\ to(\lambda x, \neg [run_*^V(x)])$.

Combination with the term *John* gives sentence (63), with translation (64).

4. COMPOUND VERB MODIFIERS

In this section I will consider sentences in which verbs occur which are accompanied by compound modifiers: constructions with *will not*, *will have*, *had*, *would*, etc. The sentences exhibit ambiguities which give us suggestions as to how to deal with compound modifiers.

The first example concerns the combination of negation and future. Sentence (69) has three readings, viz. (70), (71) and (72).

(69) *Every woman will not talk*

(70) $\forall u[woman_*(u) \rightarrow \neg W[talk_*(u)]]$

(71) $\neg W\forall u[woman_*(u) \rightarrow talk_*(u)]$

(72) $\neg \forall u[woman_*(u) \rightarrow W\ talk_*(u)]$.

Notice that in all readings negation has wider scope than future. The first two readings are the most likely ones. A situation in which the relative scope of the third reading seems to be intended arises in HOPKINS (1972, p.789). Hopkins argues that it is not necessary to always design elegant computer programs because

(73) *Every program will not be published. Many will be used only once.*

In the PTQ approach only readings (70) and (71) can be obtained. This is due to the fact that in PTQ tense and negation are treated as an indivisible unit which is introduced by one single step.

For sentence (74) related ambiguities arise. The sentence is three ways ambiguous.

(74) *The president will have talked.*

This sentence may concern

- (i) An action of the present president (maybe after his presidency).
- (ii) An action of some president during his presidency (maybe a future president).

(iii) An action of a future president (maybe before his presidency).

This readings are presented in (75), (76) and (77) respectively.

(75) $\exists u[\forall v[\textit{president}_*(v) \wedge u = v] \wedge \textit{WH}[\textit{talk}_*(u)]]$

(76) $\textit{WH}\exists u[\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge \textit{talk}_*(u)]$

(77) $\textit{W}\exists u[\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge \textit{H}[\textit{talk}_*(u)]]$.

I assume that (75) is the most likely reading of (74). The relative scope of the tense operators and president as indicated in (76) is, however, the most likely reading of (78)

(78) *{In 2000 the political situation will be different since}*

A USA president will have visited Cuba.

The relative scope as indicated in (77) is the most likely reading of (79).

(79) *The president will have learned Montague grammar at high school.*

These examples show that the two tense operators corresponding with the compound modifier 'future perfect' may have different scope. For the other compound modifiers related examples can be given. I will give some examples of the reading in which the scope of the two operators is not the same. Sentence (80) with reading (81) can be said about Eisenhower.

(80) *The president had been a general. {Therefore he knew about the power of the military-industrial complex}*

(81) $\textit{H}\exists u[\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge \textit{H}[\textit{general}_*(u)]]$.

Sentence (82) gives information about the former Dutch queen Wilhelmina.

(82) *{In May 1940 Wilhelmina had to leave her country but}*

The queen would return to Holland.

(83) $\textit{H}\exists u[\forall v[\textit{queen}_*(v) \leftrightarrow u = v] \wedge \textit{W}[\textit{return}_*(u)]]$.

Notice that in sentence (82) *would* is used to indicate a certain temporal sequence. At the moment in the past under consideration, the return was still in the future. Also the construction *would have* can be used to describe a certain temporal sequence. The information about queen Wilhelmina given above can be extended by (84).

(84) *At her departure she was just the queen, at the moment of her return she would have become a symbol of patriotism.*

The use of *would* and *would have* described above is somewhat exceptional. More frequently they are used in constructions like (85) and (86).

(85) *John would come, but he is not here.*

(86) *If John had come, we would have won the game.*

I do not intend to deal with constructions like (85) and (86), but only with the 'temporal' constructions.

For simple modifiers ambiguities of the kind considered above do *not* arise: (87) does *not* have reading (88), which would express the fact that the action may take place after the presidency of a future president.

(87) *The president will visit Holland*

(88) $W\exists u[\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge W[\textit{visit}_*(u, \textit{Holland})]]$.

The ambiguities considered in this section suggest that the compound modifiers have, for semantic reasons, to be introduced in a process with several stages, each stage introducing one operator in the translation. For instance, a sentence containing *will have* is obtained by first introducing perfect and next introducing future. Analogously *had* is analyzed as past + perfect and *would* as past + future. The semantic ambiguities can easily be accounted for since for an operator we have the options of introducing it on the level of verb phrases and of introducing it on the level of sentences.

Besides the semantic arguments there is also syntactic evidence for the introduction of compound modifiers by means of a process with several stages. Some compound modifiers can be split up when they occur in connection with a conjoined verb phrase. An example is (89).

(89) *The president will have talked and have walked.*

In (89) the verb phrases *have talked* and *have walked* share the auxiliary verb *will*.

The main conclusion of this section is that compound modifiers have to be introduced by a process with several stages, each stage introducing a new operator in the translation. An example illustrating this process is sentence (90) with reading (91).

(90) *The president will have talked*

(91) $W\exists u[\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge H[\textit{talk}_*(u)]]$.

The first step is the production of the verb phrase (92), which has

translation (93).

(92) *have talked*

(93) $\lambda x H[\text{talk}_*(x)]$.

Next sentence (94) is formed with translation (95).

(94) *The president has talked.*

(95) $\exists u[\forall v \text{ president}_*(v) \leftrightarrow u = v] \wedge H[\text{talk}_*(u)]$.

The last step is the introduction of the future, this yields sentence (90) with translation (91).

5. COMPLEX CONSTRUCTIONS

5.1. Introduction

In the previous sections we considered scope phenomena of simple and of compound verb modifiers. In this section scope phenomena will be considered in connection with more complex constructions than considered before. The most important ones are conjoined and disjoined phrases and combinations of them. I will use the name conjoined phrases to cover such conjunctions and disjunctions except where the difference is relevant. This section has a somewhat different character than the previous two, because conjoined constructions give rise to phenomena which do not constitute a clear and simple pattern. The acceptability of the sentences or interpretations is sometimes marginal and the judgements may have to be changed in some cases. The present discussion is intended primarily to point out some interesting phenomena.

5.2. Conjoined verb phrases with positive verbs

Conjoined verb phrases which consist of unmodified verbs give rise to the same phenomena as single verbs. The conjoined phrases can be modified as if they were simple verbs and they exhibit the same ambiguities. An example is sentence (96), which has readings (97) and (98).

(96) *The president will walk and talk*

(97) $W\exists u[\forall v[\text{president}_*(v) \leftrightarrow u = v] \wedge [\text{walk}_*(u) \wedge \text{talk}_*(u)]]$.

(98) $\exists u[\forall v[\text{president}_*(v) \leftrightarrow u = v] \wedge W[\text{walk}_*(u) \wedge \text{talk}_*(u)]]$.

The formulas (97) and (98) present the possible readings as far as the

position of *president* with respect to the future operator is concerned. Both readings, however, say that on a moment on the future a certain person will both walk and talk. Probably this is too precise, and a better interpretation would be that there is a future interval of time in which both the walking and the talking are performed, possibly on different moments in that interval. So this kind of objections against (97) and (98) might be solved by using another model relative to which the formulas are interpreted. But concerning the scope aspect, the formulas seem correct, and therefore the rules will produce only (97) and (98) as translations for (96).

Conjoined verb phrases with verbs which are modified differently only have a reading in which both operators have narrow scope. We have already met example (99) with reading (100).

(99) *A woman has walked and will run.*

(100) $\exists u[\text{woman}_*(u) \wedge H[\text{walk}_*(u)] \wedge W[\text{run}_*(u)]]$.

If the verbs of the conjoined phrase are modified in the same way, there is a reading which corresponds with the above example: sentence (101) has a reading (102)

(101) *The president will walk and will talk*

(102) $\exists u[\forall v[\text{president}_*(v) \leftrightarrow u = v] \wedge W[\text{walk}_*(u)] \wedge W[\text{talk}_*(u)]]$.

Sentence (101) can, however, be considered as dealing with a future president, so it also has reading (103).

(103) $W\exists u[\forall v[\text{president}_*(v) \leftrightarrow u = v] \wedge \text{walk}_*(u)] \wedge \text{talk}_*(u)$.

The possibility that the walking and talking are performed on the same moment in the future can be dealt with in the same way as I suggested for sentence (96). The fact that sentence (101) has reading (103) (= 97!) suggests us that we consider sentence (101) as a syntactic variant of (96) and assign it, too, reading (98). The same treatment will be given of the perfect.

For the past tense the same pattern applies: sentence (104) not only has reading (105) but also readings (106) and (107).

(104) *The president walked and talked.*

(105) $\exists u[\forall v[\text{president}_*(v) \leftrightarrow u = v] \wedge H[\text{walk}_*(u)] \wedge H[\text{talk}_*(u)]]$

(106) $\exists u[\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge \textit{walk}_*(u) \wedge \textit{talk}_*(u)]$

(107) $\exists u\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge H[\textit{walk}_*(u) \wedge \textit{talk}_*(u)]$.

Conjoined negated verbs do not follow this pattern. Sentence (108) has reading (109), but it has no reading with only one negation sign.

(108) *The president does not walk and does not talk.*

(109) $\exists u[\forall v[\textit{president}_*(v) \leftrightarrow u = v] \wedge \neg [\textit{walk}_*(u)] \wedge \neg [\textit{talk}_*(u)]]$.

A conjoined verb phrase which consists of equally modified verbs can, in some cases, be modified further. An example is (110), where a modifier is applied to a conjunction of verbs in the perfect.

(110) *The president will have visited Rome or have visited Tokyo.*

Conjoined verb phrases with equally modified verbs cannot be negated, as (111) illustrates. That example cannot be interpreted as a negation of a conjunction of perfect verb phrases, but only as a negated verb phrase conjoined with a non-negated one.

(111) *The president has not visited Rome or has visited Tokyo.*

If another modifier is applied first, the phrase behaves as a simple construction and can be negated, see (112).

(112) *The president will not have visited Rome or have visited Tokyo.*

5.3. Conjoined verb phrases with negated verbs

If in a conjoined verb phrase the first verb is not modified and the other verbs are negated, then the whole construction behaves as a verb phrase with unmodified verbs. This means that such a construction can be modified further; an example is (113) with reading (114).

(113) *John will walk and not talk*

(114) $W[\textit{walk}_*(john) \wedge \neg [\textit{talk}_*(john)]]$.

Note that sentence (113) is not ambiguous with respect to the scope of *W* because the interpretation of *John* is index independent. Were *John* be replaced by *the president*, then ambiguities would arise of the kind we have discussed before.

If in a conjoined verb phrase the first phrase is negated and the others are not negated, then the situation is different. A modifier

'absorbs' the negation: sentence (115) only has reading (116).

(115) *John will not walk and talk*

(116) $\neg W[\text{walk}_*(\text{john}) \wedge \text{talk}_*(\text{john})]$.

If all the verbs in a conjoined verb phrase are negated, then the two patterns give rise to an ambiguity. Sentence (117) has both reading (118) and (119).

(117) *John will not walk and not talk*

(118) $W[\neg \text{walk}_*(\text{john}) \wedge \neg \text{talk}_*(\text{john})]$

(119) $\neg W[\text{walk}_*(\text{john}) \wedge \text{talk}_*(\text{john})]$.

For conjoined verb phrases with verbs in the perfect a related situation arises. Sentences (120) and (121) seem to have one reading, whereas (122) has two readings.

(120) *John will not have walked and have talked*

(121) *John will have walked and not have talked*

(122) *John will not have walked and not have talked.*

Corresponding with the above sentences there are sentences with the contracted forms like *won't*. Sentence (123) has the same reading as its uncontracted variant (120).

(123) *John won't have walked and have talked.*

Sentence (124), however, is not equivalent with the corresponding uncontracted form (117): it only has reading (119), but not reading (118).

(124) *John won't walk and not talk.*

The way in which we may treat contracted forms depends on the organization of the morphological component. Suppose that one decides that the input of the morphological component has to consist of a string of words (where the words may bear features). Then the contraction of *will not* to *won't* cannot be dealt with in the morphological component because sentence (118) gives no syntactic information about the intended reading. This means that the contraction has to be described within the rules: the rule introducing negation should have the option of producing contracted forms like *won't*. If one has the opinion that the input of the morphological component has to be a syntactic structure, then the situation is different. I assume

that sentence (117) will have a structure in which *will* is directly connected with *not* and a structure in which *walk* is directly connected with *not*. This structural information desambiguates sentence (117) and provides sufficient information to deal with contracted forms: *will not* only reduces in case it is a constituent.

5.4. Terms

The PTQ fragment only has terms which require a third person singular of the finite verb. This is probably caused by the desire to keep the syntax simple. Incorporating pronouns for the first and second person singular would not be interesting in the light of our investigations for the following reason. The pronouns *I* and *you* get an index independent interpretation and therefore (125) and (126) are not ambiguous.

(125) *I will have visited China*

(126) *You have discovered the solution.*

In what follows we will only consider 'third-person' terms.

Disjoined terms give rise to the same scope phenomena as simple terms. Sentence (127) has a reading that says that the present president or the present vice president will go, and a reading that says that the future president or future vice-president will go.

(127) *The president or the vice-president will visit Holland.*

A complication may arise from quantifying in. One might first produce (128) and obtain (127) from this by means of quantifying in.

(128) *The president or he₁ will visit Holland.*

This might result in a reading in which the present vice-president or the future president will visit Holland. Such mixed readings are not possible for sentence (127). This means that we have to constrain the possible applications of the quantification rule. I have not investigated these matters and I will therefore simply assume the (ad hoc) restriction that there are no terms of the form T_1 or T_2 in the fragment, where one or both terms are indexed pronouns.

In the examples above the determiners *the* and *a* are most frequent. For the term *every president* corresponding results are obtained: sentence (129) gets readings (130) and (131).

(129) *Every president will talk*

(130) $W[\forall u \text{ president}_*(u) \rightarrow \text{talk}_*(u)]$

(131) $\forall u[\text{president}_*(u) \rightarrow W[\text{talk}_*(u)]]$.

If (129) concerns future presidents, it is unlikely that they have to be presidents at the same moment. One might try to represent such a meaning by formula (132).

(132) $\forall uW[\text{president}_*(u) \rightarrow \text{talk}_*(u)]$.

This is, however, not correct, since (132) would (vacuously) be true in situations such that for everyone there is a future moment at which he is *not* a president. I expect that the desired reading can be obtained by interpreting formula (130) in some model with interval semantics for time. Then (131) might get the interpretation that there is an interval in the future during which all individuals who are president in that interval will talk during that interval. The scope aspect of the meaning of (129) is then adequately represented by formulas (130) and (131).

For conjoined terms the same ambiguities will be obtained as for disjoined terms. Sentence (133) has a reading about present statesmen and one about future statesmen.

(133) *The president and the vice president will visit Cuba.*

The problem of 'mixed' readings, noticed with respect to disjunctions, also arises here, and for conjoined terms a corresponding (ad hoc) restriction on quantifying in is required. Furthermore there is the same difficulty as for the term *every president*. It is not necessary that the two statesmen of sentence (133) will visit Cuba together. A solution might be found following the suggestions concerning the interpretation of (129).

5.5. Embeddings

An important source of scope phenomena are the embedded sentences (in verb complements and in relative clauses). LADUSAW (1974) and EJERHED (1981) point out several sentences that are not treated correctly in PTQ. A variant of an example of Ladusaw is (133).

(133) *Mary has found the unicorn that walks.*

The rules produce the possible reading in which the unicorn presently walks. But they also produce a reading in which the unicorn walks on the moment of

discovery (which is not a possible reading). For the future tense this ambiguity seems to be correct, see sentence (134).

(134) *Mary will find the unicorn that walks.*

An example from EJERHED (1981) is

(135) *Bill will assert that John loves Mary.*

She argues that this sentence is ambiguous. On the one reading John loves Mary at the moment of asserting, and on the other reading he loves her now. PTQ cannot distinguish between these readings, nor can the present treatment.

In order to deal with embeddings, Ladusaw makes his syntactic rules rather complex (using e.g. dominance relations) and his success is partial. I would try to find a solution in the logic. Priorian tense logic is not a suitable logical language to deal with the semantics of embedded sentences. This is illustrated by example (136).

(136) *A child was born that will become ruler of the world.*

The *will* of the embedded sentence takes as 'starting point' the reference point used for the interpretation of the whole sentence, and not the reference point introduced by the past tense. Sentence (136) was one of Kamp's arguments for introducing the 'Now'-operator (KAMP 1971). However, more power is required. The 'now'-operator keeps trace of the first point of reference one encounters during the evaluation of the sentence: the point of utterance. SAARINEN (1978) gives examples which show that one needs to be able to keep trace of all points of reference one encounters in evaluating the sentence. One of his examples is (137).

(137) *Bob mentioned that Joe has said that a child had been born who would become ruler of the world.*

Saarinen argues that the *would* can have as starting point for its evaluation any of the reference points introduced by the previous past tense operators. So each operator introduces its own variant of 'now'. This means that considerable expressive power has to be added to the logic we use for representing meanings. Since I use the logic of PTQ, with its Priorian tense operators, it is not surprising that embedded constructions in general are not treated correctly by my grammar.

6. ONE OF THE RULES

Most of the scope phenomena discussed in the previous sections will be treated explicitly: in section 7 by providing a grammar. That grammar is in some respects different from the grammar used for the PTQ fragment. The differences have already been introduced in chapter 8: words may bear features, information provided by queries is used, the rules produce labeled bracketings (or, equivalently, labeled trees), and in the formulation of the rules certain basic operations can be mentioned. These aspects of the grammar will be considered below, thereafter one of the rules will be discussed extensively.

The features are used mainly to facilitate the explicit formulation of the rules. It is, for instance, shorter to write formulation (A) instead of (B), and probably easier to understand.

(A) add features ((past, sing 3), δ)

(B) replace δ by its third person singular past tense form.

The features are not intended as a part of a general theory about features, and therefore I will only introduce those features which I find useful for the treatment of the present fragment. These are: *past*, *pc* (for participles) and *sing 3* (for the third person singular). Other features are not needed (e.g. there is no feature *pres* since $walk_{sing3,past} \rightsquigarrow walked$, and $walk_{sing3} \rightsquigarrow walks$).

The most important query that will be used is *Fin*. The Fins of a sentence or verb phrase are its finite verbs, i.e. the verbs which agree (in person and number) with the subject of the sentence. So it is about the same as the query *Mainverb* introduced in chapter 7. I prefer to avoid the name *mainverb* in the present context, because auxiliary verbs (such as *will* and *do*) can be used as finite verbs, and maybe not everyone would be happy to call those auxiliary verbs 'mainverbs'. The other query that will be used is *Verbphrase*. It gives the information what the verbphrase of a given sentence is. For the present fragment it turned out to be the most simple to define the queries directly on all trees, and not within the root operation (as was the method employed in chapter 7).

The labeled bracketing are used mainly to give a correct treatment of conjoined phrases. FRIEDMAN (1979) has shown that for dealing correctly with the conjoined and embedded phrases of the PTQ fragment, it is sufficient to have the bracketing available: the labels are not needed.

For the fragment under discussion the same holds: no rule needs the information provided by the labels. The choice of the labels is, for our purposes, arbitrary. Which labels actually have to be chosen, can only be decided if larger fragments of English are considered, then we might decide which rules need which information. The decision to call *will* in *John will run* an auxiliary is arbitrary from my point of view, I have no arguments pro or contra this choice. Since labels play no essential role in the discussion, I will simplify the presentation of the grammar by omitting the labels, e.g. in the presentation of the produced expressions and in the formulation of the root operations. Furthermore, I will omit brackets around simple lexical items. (e.g. using *run* instead of $[run]$). These simplifications allow me to write (139) instead of (138).

(138) $[[John]_T [love_{sing3}]_{TV} [Mary]_T]_{IV} S$

(139) $[John [love_{sing3} Mary]]$.

The basic operations we will use are *root* and *adjoin*. The operation *root* takes a sequence of trees as argument, and connects them with a new common root, labeled with a given category (see chapter 7). The operation *adjoin* takes two trees as argument, and connects them with a new root, which bears the same label as the root of the second argument.

As introduction to the presentation of the grammar I will extensively consider one of the rules. It is the rule which has the effect that the modifier for future tense is introduced into sentence (140), thus obtaining (141).

(140) $John walk_{sing3} and talk_{sing3}$

(141) $John will_{sing3} walk and talk$.

Every sentence cannot be used as input for this rule; for instance a sentence in the future tense cannot be futurized again. There have to be restrictions on the possible applications of a rule introducing future. For other modifiers the same holds: not every sentence can be modified. One might wish to have in the grammar a single rule for the introduction on sentence level of all modifiers. This rule has to mention under which circumstances a future may be introduced, and the same for other modifiers. Moreover for each modifier it has to describe precisely in which way it has to be introduced. In this way we would obtain one great rule with a lot of subclauses. For reasons of elegance and understandability I prefer to have for each modifier a separate rule.

We have to characterize the sentences to which a certain modifier can be added. There is a hierarchy which accounts for the relative scopes of modifiers as we observed this in sections 3 and 4 (conjoined phrases give rise to complications). The hierarchy is

[neg[past[fut[perf]]]].

This hierarchy claims, for instance, that negation always has wider scope than the perfect. It also says that future can be added to a positive perfect sentence and to a positive sentence in the present tense. It says moreover that future cannot be added to a negated sentence because that would give rise to an incorrect order of scope.

The hierarchy suggest to us how the possible applications of the rule introducing future has to be restricted. It can only be applied to sentences in the positive present perfect and in the positive present tense. The rule introducing future then has to contain a specification of what such sentences look like. One might expect as characterization of sentences in the positive present perfect that the sentence has as finite verb the auxiliary *have*, and as characterization of a present tensed sentence that its finite verb is in the present tense. Such a description is not sufficient. In the description of the present tensed sentences one has to exclude finite verbs which are modifiers themselves (*has, will, do*). Furthermore we have to exclude negations. If conjoined verb phrases are involved, further caution is required. These considerations show that the desired characterizations will become rather complex. I do not doubt that such characterizations are possible, but I prefer to use a somewhat different method.

The method I prefer consists of subcategorization of the sentences and verb phrases. This subcategorization is not obtained by describing explicitly which sentences belong to a certain subcategory, but indirectly by means of the rules. The rule which introduces the perfect gives the information that the sentence obtained belongs to the subcategory of sentences in the perfect tense and the rule which introduces negation gives the information that if the rule is applied to a perfect sentence the resulting sentence is the subcategory of negated perfect sentences. In this approach the rules take expressions of specified subcategories and produce expressions of specified subcategories. In this way we avoid complex conditions in the rules: the grammar does the job.

I have already mentioned two subcategories of expressions to which

future tense can be added: the positive sentences in the present tense and those in the present perfect. For these subcategories I will use the names *perf S* and *S* respectively. For the category of all sentences I will use the name *full S*, so *S* in this chapter has a different meaning than in PTQ. In the rules many more subcategories are relevant than the ones mentioned here. The names of almost all subcategories that will be used, are indicated by the following scheme of names:

(neg) (past) (fut) (perf) S.

The names of subcategories are obtained from this scheme by replacing each subexpression of the form (α) by the expression α or by the empty string. Some examples of subcategories are as follows:

<u>name</u>	<u>intuitive characterization</u>
S	sentences in the positive present tense
neg past S	negated sentences in the past tense
past perf S	unnegated sentence in the past perfect

For verb phrases a related system of subcategories will be used. The system is somewhat larger because there are some categories for conjoined phrases, e.g. the category of conjoined phrases consisting of verbs in the perfect. The names which can be used are given by the following scheme

(conj) (neg) (past) (fut) (perf) IV.

Whether a conjoined phrase belongs to a subcategory of conjoined phrases is determined by the rules. This might have as a consequence, however, that the subcategorization of a phrase and the intuitive expectation about this do not always coincide. One might, for instance, expect that *will have walked and have talked* is a conjoined phrase. Since it behaves as a single verb in the future tense it is considered as an expression of the subcategory fut IV. For the set of all verb phrases we use the name full IV, the subcategory IV consists (in principle) of unmodified verbs.

Now I return to the rule under discussion: the one which introduces future tense. One might design a two-place rule which combines the modifier *will* with a sentence of the subcategory *S* or *perf S*. Then the rule yields a sentence of (respectively) the subcategory *fut S* or *fut perf S*. The translation of *will* introduced on the level of sentences has to be $\lambda p\mathcal{W}[\overset{\vee}{p}]$, where p is a variable of type $\langle s, t \rangle$, and the translation rule corresponding with this syntactic rule could then be $\text{MOD}'(\wedge'S')$. Such a rule exhibits a

a remarkable property: there is just one expression which can be used as first argument of the rule. Since only one argument is possible one could as well incorporate all information about this argument in the rule. In this way the rule with two arguments is replaced by a rule with one argument. I consider such a one-place rule as simpler and therefore I will follow this approach.

A one-place rule which introduces future in a given sentence has to contain some syntactic operation which has the effect of introducing *will*. In this way *will* becomes a syncategorematic symbol. This *will*, when considered in isolation, does not get a translation. But this does not mean that its introduction has no semantic effect: its effect is accounted for by the translation rule (which introduces the future tense operator *W*). Nor does the syncategorematic introduction of *will* mean that it has no syntactic status. The role of *will* in the syntax can be accounted for in the surface structure which is produced by the rule. There it can be given the position it should get on syntactic grounds and there it can get the label it should bear.

For other verb modifiers the same approach will be followed. There is no semantic or syntactic reason to have essentially different derivational histories for past sentences and sentences with future. Both verb modifiers can be introduced by means of one-place rules. That there is a great syntactic difference (in English) between past and future can be accounted for in the produced bracketing: there the introduction of past has the effect of the introduction of an affix and the introduction of future the effect of introducing an (auxiliary) verb. Also the difference between future tense in French (where it is affix) and in English can be accounted for in the labeled bracketing. Notice that the decision to introduce verb modifiers syncategorematically is not made for principled reasons, but just because it gives rise to a more elegant grammar.

Next I will consider the formulation of the rule introducing future on the level of sentences. This rule can be considered as consisting of two rules: one producing expressions of subcategory fut S (from expressions in the subcategory S) and one producing expressions of the subcategory fut perf S (from perf S expressions). The subcategorical information is combined in the following scheme (or hyperrule, see the discussion on Van Wijngaarden grammars in chapter 6, section 5):

$$R_{\text{fut}}: (\text{perf})S \rightarrow \text{fut}(\text{perf})S.$$

From this scheme we obtain information about actual rules by replacing (perf) on both sides of the arrow consistently either by perf or by the empty string. The scheme says that there is a rule (function) from the subcategory S to the subcategory fut S and a function from the subcategory perf S to the subcategory fut perf S. Which particular rules there are is determined by the syntactic operation F_{fut} . It consists of two syntactic subfunctions which have to be performed consecutively.

F_{fut} : delete (sing 3, Fin(S));
 adjoin (*will*_{sing3}, verb phrase(S)).

Agreement is dealt with in a primitive way: the rule is correct only for subjects which require the third person singular form of the verb. This is sufficient because our fragment contains only such terms. Notice that there is for both rules indicated in the scheme, one single syntactic operation. For the corresponding translation rule the same holds: there is one translation rule which reads as follows:

T_{fut} : W[α³].

7. THE GRAMMAR

7.1. Introduction

Now we come to the kernel of the proposal: the rules. Presenting an discussion on how to treat a certain phenomenon is one step, but providing for explicit rules is another important step. The rules presented here are not just a formalization of the previous discussion. They contain more information because I have to be explicit about details I did not discuss (see also section 7.5). The rules do not deal with all phenomena mentioned in section 2 (simple modifiers) and in section 3 (compound modifiers). Furthermore the rules deal with all phenomena concerning conjoined verb phrases discussed in 4.2 and 4.3, except for the contracted forms. As for 4.4, the fragment contains disjuncted terms, but no conjoined ones. Although embedded constructions are in the fragment, the predictions of the rules are in several cases incorrect.

The fragment described by the rules is an extension and variation of the PTQ fragment. The lexical elements are supposed to be the same as in PTQ, except for verbs like *try to*, which loose their *to*. The rules (schemes) presented below, replace the PTQ rules S_3 (relative clauses), S_4 (IV+T),

S_8 (IV/IV+IV), S_9 (S/S), S_{10} (IV//IV+IV), S_{14} (quantification of T into S), S_{15} (quantification into IV), and S_{17} (variants of S_4). Other rules are assumed to be as in PTQ, with the change that now bracketings are produced.

The rules will be presented in the form described in the previous section; i.e. by presenting their S,F, and T component. Furthermore, some of the rules are accompanied by comments or examples. In the examples the subcategory of the produced expression is mentioned between braces. The rules are divided into six groups. Each rule bears an index in the 900-series.

7.2. Rules

I. Rules modifying verb phrases

S_{901} : (conj)IV \rightarrow perf IV

F_{901} : if *do* is among $\text{Fin}(\alpha)$ then delete this *do*;
add feat (pc,Fin(α)); adjoin (*have*, α)

T_{901} : $\lambda xH[\alpha'(x)]$

example: F_{901} (*walk and*[[*do not*]*talk*]) = [*have*_{pc}[*walk*_{pc} and[*not talk*_{pc}]]] =
have walked and not talked {perf IV}.

comment 1: The subcategory indication *conj* is not mentioned in the output subcategory because the resulting phrase behaves as an simplex verbphrase in the perfect.

S_{902} : (conj)(perf)IV \rightarrow fut(perf)IV

F_{902} : if *do* occurs in $\text{Fin}(\alpha)$, then delete this *do*; adjoin (*will*, α)

T_{902} : $\lambda xW[\alpha'(x)]$

example: F_{902} (*walk and*[[*do not*]*talk*]) = [*will*_{pc}[*walk and*[*not talk*]]]
{fut IV}

S_{903} : (fut)(perf)IV \rightarrow past(fut)(perf)IV

F_{903} : add features (past,Fin(α))

T_{903} : $\lambda xH[\alpha'(x)]$

examples: F_{903} (*walk and*[[*do not*]*talk*]) = [*walk*_{past} and[[*do*_{past} not]*talk*]] =
walked and did not talk {past IV}

F_{903} (*will walk*) = [*will*_{past} *walk*] = *would walk* {past fut IV}

comment: Notice that this rule has the same translation rule as the rule introducing perfect (S_{901}). In case we use a logic which allows for dealing with the semantic differences between past and perfect, the translation rules would be different.

II. Rules producing tensed sentences

S₉₀₄ : S → perf S
 F₉₀₄ : delete features (sing 3,Fin(α)); F₉₀₁ (verb phrase(α));
 add feat (sing 3,Fin(α))

T₉₀₄ : H[α']

example : F₉₀₄ ([John walk_{sing3}]) = [John have_{sing3} walk_{pc}] =
 John has walked {perf S}

comment : If one decided that *have* cannot have wide scope (see section 3),
 then this rule would have to be removed from the syntax.

S₉₀₅ : (perf)S → fut(perf)S

F₉₀₅ : delete features (sing 3,Fin(α)); F₉₀₂ (verbphrase(α));
 add features (sing 3,Fin(α))

T₉₀₅ : W[α']

S₉₀₆ : (fut)(perf)S → past(fut)(perf)S

F₉₀₆ : add features (past,Fin(α))

T₉₀₆ : H[α'].

III. Rules for negation

S₉₀₇ : (conj)(past)(fut)(perf)IV → neg(past)(fut)(perf)IV

F₉₀₇ : case 1 there is one verb in Fin(α):
 let *f* be the list of features of Fin(α)
 if Fin(α) is *be*, *will* or *have* then replace it by [*be_f* not],
 [*will_f* not] or [*have_f* not] respectively;
 otherwise adjoin (root(*do_f*,not),α).
case 2 there is more than one verb in Fin(α).
 if *do* is in Fin(α) then delete this *do*;
 adjoin (root(*do*,not),α).

T₉₀₇ : λx ⊃ [α'(x)]

examples : F₉₀₇ ([will walk]) = [[will not]walk] = will not walk {neg fut IV}
 F₉₀₇ ([try[not[to walk]]) = [[do not][try[not[to walk]]]] =
 do not try not to walk {neg IV}
 F₉₀₇ ([walk and[[do not]talk]) = [[do not][walk and[not talk]]] =
 do not walk and not talk {neg IV}

S₉₀₈ : (past)(fut)(perf)S → neg(past)(fut)(perf)S

F₉₀₈ : F₉₀₇ (verb phrase(α))

T₉₀₈ : ⊃ α'.

IV. *IV-complements and adverbs*

S_{909} : $IV//IV \times (\text{conj})(\text{neg})(\text{perf})IV \rightarrow IV$

F_{909} : if *do* is the only element of $\text{Fin}(\beta)$ then produce
 $\text{root}(\alpha, \text{root}(\text{not}, \text{root}(to, \tilde{\beta})))$, where $\tilde{\beta}$ is obtained from β by
 deleting *do not*
 otherwise
 if there are occurrences of *do* in $\text{Fin}(\beta)$ then delete these *do*'s;
 $\text{root}(\alpha, \text{root}(to, \beta))$

T_{909} : $\alpha'(\hat{\beta}')$

examples: F_{909} (*try*, [[*do not*]*run*]) = [*try*[*not*[*to run*]]] {IV}
 F_{909} (*hope*, [*have talk*_{pc}]) = [*hope*[*to*[*have talk*_{pc}]]] {IV}
 F_{909} (*wish*, [*walk and*[[*do not*]*talk*]]) = [*wish*[*to*[*walk and*[*not talk*]]]]
 {IV}.

comment: The resulting phrases are of the subcategory IV because all verb modifiers can be added to them. The possible inputs of the rule are characterized as $(\text{conj})(\text{neg})(\text{perf})IV$, predicting that all verb phrases of the corresponding categories can be input for the rule. This prediction is incorrect witness (142).

(142) *John regrets to have talked.*

Further investigations are required in order to decide which verbs take which modified complements.

S_{910} : $IIV \times (\text{neg})(\text{conj})IV \rightarrow IV$

F_{910} : $\text{root}(\beta, \alpha)$

T_{910} : $\alpha'(\hat{\beta}')$

examples: F_{910} (*slowly, talk*) = *talk slowly*
 F_{910} (*voluntarily, [do[not[talk]]]*) = [[*do[not talk]*]*voluntarily*]
 {IV}.

V. *Rules for conjoined phrases*

In section 5 we observed that conjoined phrases behave in various ways. This means that they are in various subcategories and that they have to be produced by several rules. The first two rules mentioned below do not create a conjoined phrase, but say that all modified verb phrases and sentences are members of the categories full IV and full S respectively. Most conjunction and disjunction rules are defined on these categories.

S_{911} : (conj)(neg)(past)(fut)(perf)IV \rightarrow full IV
 F_{911} : no change of the expression
 T_{911} : α'
 S_{913} : full IV \times full IV \rightarrow full IV
 F_{913} : root(α , *and*, β)
 T_{913} : $\lambda x[\alpha'(x) \wedge \beta'(x)]$
 S_{914} : full S \times full S \rightarrow full S
 F_{914} : root(α , *and*, β)
 T_{914} : $\alpha' \wedge \beta'$
 S_{915} : as S_{913} but now for disjunction
 S_{916} : as S_{914} but not for disjunction.

The following two rules produce verb phrases which can be modified further.

S_{917} : IV \times (neg)IV \rightarrow conj IV
 F_{917} : root(α , *and*, β)
 T_{917} : $\lambda x[\alpha'(x) \wedge \beta'(x)]$
 S_{918} : as S_{917} but now for disjunction.

The following rules produce constructions with an exceptional character.

S_{919} : neg IV \times neg IV \rightarrow conj perf IV
 F_{919} : delete *do* from α ; add feature(pc, Fin(α));
 delete *do* from β ; add feature(pc, Fin(β));
 root(*have*, root(α , *and*, β))
 T_{919} : $\lambda xH[\alpha'(x) \wedge \beta'(x)]$
 example : S_{919} ([[*do not*]walk], [[*do not*]talk]) =
 [*have*[[*not walk*_{pc}] and [*not talk*]]] {conj perf IV}
 The corresponding translation is
 $\lambda xH[\text{walk}(x) \wedge \text{talk}(x)]$.
 Note that the output of S_{919} can be used as input for S_{902} , i.e.
 future tense can be added to the output of S_{919} .

S_{920} : perf IV \times (neg)perf IV \rightarrow fut perf IV
 F_{920} : delete *do* from β ; adjoin (*have*, root(α , *and*, β))
 T_{920} : $\lambda xW[\alpha'(x) \wedge \beta'(x)]$
 example : F_{920} (*have walk*_{pc}, [[*do not*][*have talk*_{pc}]]) =
 [*will*[*have walk*_{pc}] and [*not have talk*_{pc}]] =
 will have walked and not have talked {fut perf IV}

S_{921} : neg perf IV \times neg perf IV \rightarrow neg fut perf IV

F_{921} : delete *do* from α ; delete *do* from β ;
adjoin(*will*, root(α , and, β))

T_{921} : $\lambda xW[\alpha'(x) \wedge \beta'(x)]$

example: F_{921} ([[do not][have walk_{pc}]], [[do not][have talk_{pc}]]) =
[will[[not[have walk_{pc}]] and [not[have talk_{pc}]]]] =
will not have walked and not have talked {neg fut perf IV}.

comment: If the example given with rule S_{920} is negated, the resulting phrase is identical with the example given for rule S_{921} . The respective translations are different, thus accounting for the ambiguity noted in section 5.

$S_{922}, S_{923}, S_{924}$ as $S_{919}, S_{920}, S_{921}$, but now for disjunction.

VI. Other rules

S_{925} : T \times full IV \rightarrow full S

F_{925} : add feature (sing3, Fin(β))
root(α, β)

T_{925} : $\alpha'(\wedge\beta')$

S_{926} : T \times (neg) (past) (fut) (perf)IV \rightarrow (neg) (past) (fut) (perf)S

F_{926} : $F_{925}(\alpha, \beta)$

T_{926} : $\alpha'(\wedge\beta')$

$S_{927,n}$: CN \times full S \rightarrow CN

$F_{927,n}$: see $F_{3,n}$ in PTQ

$T_{927,n}$: see $T_{3,n}$ in PTQ

S_{928} : S/S \times full S \rightarrow full S

F_{928} : adjoin (α, β)

T_{928} : see T_7 in PTQ

comment: The requirement that the sentence is an element of the category full S prevents the introduction of a verb modifier after application of S_{928} . Hence negation cannot have wide scope in:

(143) *Necessarily John does not run.*

$S_{929,n}$: T \times (neg) (past) (perf)S \rightarrow (neg) (past) (fut) (perf)S

$F_{929,n}$: see $F_{10,n}$ in PTQ

$T_{929,n}$: see $T_{14,n}$ in PTQ

$S_{930,n} : T \times (\text{neg})(\text{past})(\text{fut})(\text{perf})IV \rightarrow (\text{neg})(\text{past})(\text{fut})(\text{perf})IV$

$F_{930,n} : \text{see } F_{10,n} \text{ in PTQ}$

$T_{930,n} : \text{see } T_{15,n} \text{ in PTQ}$

S_{931}, S_{932} as S_{929} and S_{930} , but now for the categories full S and full IV respectively.

7.3. Morphology

I will not explicitly describe a morphological component since that would be an ad hoc version. I have already sketched (section 4) two views on what the input for this component could be: either the whole surface structure or only the string of lexical items (with features). In both approaches it cannot be determined whether a certain occurrence of *will* was introduced on sentence level or on verb phrase level. There is for the morphological component just one *will*. Analogously there is just one *have*, whether it was introduced as auxiliary at some stage, or as a main verb describing the relation between owner and property.

7.4. Fins and verb phrase

In the rules the queries *verb phrase* and *Fin* are used. Below I will give a definition of these queries. Although I have described the framework as one which produces labeled bracketings, I did not specify labels because they are not needed in the rules. In the definition of *Fin* the labels are useful and I will refer to them. (If the reader has objections against this situation - not introducing the labels explicitly, but still using them - then he should neglect the labels. It does not lead to different predictions of the grammar.)

In the definition below *V* is a parameter which stands for all verbal categories, i.e. *V* has to be replaced by *IV*, *IV*, *IV/IV*, or *Aux* (or whatever the label is of *will*, *have* and *do*). The *X*, *Y* and *S* stand for arbitrary labels, and \emptyset for the empty set.

$\text{Fin}(\alpha)$	= α	if α is a verb
$\text{Fin}([\alpha]_V \text{ and } [\beta]_V)$	= $\text{Fin}(\alpha) \cup \text{Fin}(\beta)$	
$\text{Fin}([\alpha]_V \text{ or } [\beta]_V)$	= $\text{Fin}(\alpha) \cup \text{Fin}(\beta)$	
$\text{Fin}([\alpha]_V [\beta]_X)_V)$	= $\text{Fin}(\alpha)$	
$\text{Fin}([\alpha]_X [\beta]_S)_Y)$	= $\text{Fin}(\alpha)$	if <i>X</i> is not a verbal category
$\text{Fin}(\alpha)$	= \emptyset	if α does not satisfy one of the above clauses.

Verb phrase is defined analogously.

Verb phrase(α) = α if α is a verb

Verb phrase($[\alpha]_S$ and $[\beta]_S$) = verb phrase(α) \cup verb phrase(β)

Verb phrase($[\alpha]_{S/S}[\beta]$) = verb phrase(β)

Verb phrase($[[\alpha]_T[\beta]_{IV}]_S$) = β

Verb phrase(α) = \emptyset if α does not satisfy one of the above clauses.

7.5. Final remarks

I would like to end by saying something about the methodology. I fully agree with the final remark of PARTEE 1979a (p.94): 'It can be very frustrating to try to specify frameworks and fragments explicitly; this project has not been entirely rewarding. I would not recommend that one always work within the constraint of full explicitness. But I feel strongly that it is important to do so periodically because otherwise it is extremely easy to think that you have a solution to a problem when in fact you don't.'

Some remarks about my experiences in formulating the rules.

1. The project was not entirely successful. It was too difficult to do everything correctly at once. By providing explicit rules, I am also explicit in cases where I know the proposals to be incorrect (see section 5), or to be ad hoc (e.g. agreement).
2. The rules are explicit about borderline cases in which it is not evident that the produced sentences or the obtained readings are possible (e.g. verb phrase complements with a verb modifier).
3. The rules describe a rather large system and they make predictions about a lot of kinds of sentences I never thought of (for instance because they do not resemble the phenomena I thought of when designing the rules). I would feel safer about the correctness of the rules if I had a computer program producing hundreds of sentences of the fragment, together with their reduced translations.
4. Writing explicit rules forced me to consider the 'irrelevant' details. It turned out for instance that of the three methods for defining Fin's mentioned in JANSSEN 1980, in fact only one was applicable.
5. Considering some larger fragment explicitly gave me suggestions for finding arguments. I have presented a related treatment of verb modifiers in JANSSEN 1980 as well, but most of the arguments given in sections 2, 3 and 4 of this chapter are new, and these were found when I extended

the fragment with the quantification rules and conjunction rules.

Although the first three points are not really a recommendation for the rules presented here, I would not like to call these negative consequences of working explicitly. They are inherent to such a method of working, and constitute, in my opinion, rather an advantage. Shortcomings of a proposal with explicit rules are easier found than of a proposal without. Therefore such an approach is, generally speaking, a better starting point for further research and improvements.

APPENDIX 1

INDIVIDUAL CONCEPTS IN PTQ

In chapter 4 the syntax and semantics of the PTQ fragment were presented. The common nouns and intransitive verbs of the fragment were translated into constants which denote predicates on individual concepts. Reduction rules allowed us to replace them by predicates on individuals. What is then the benefit of using such concepts? The argument given in chapter 4 was based upon the artificial name *Bigboss*. In PTQ two less artificial examples are given as justification for the translation into predicates on individual concepts.

Consider the sentences (1) and (2).

(1) *The temperature is ninety*

(2) *The temperature is rising.*

A naive analysis of (1) and (2), using standard logic, might allow to conclude for (3).

(3) *Ninety rises.*

This would not be correct since intuitively sentence (3) does not follow from (1) and (2). So we have to provide some analysis not having this consequence. This example is known as the *temperature paradox*.

Montague's second example is a variation of the temperature paradox. Sentence (6) does not follow from sentences (4) and (5), whereas a naive analysis might implicate this.

(4) *Every price is a number*

(5) *A price rises*

(6) *A number rises.*

The solution of these problems is based upon the use of individual concepts. The idea of the solution is explained as follows. Imagine the situation that the price of oil is \$ 40, and becomes \$ 50. In this situation one might say:

(7) *The price of oil changes.*

By uttering (7) one does not intend to say that \$ 40 changes, or that \$ 50 changes. It is intended to express a property of the oil price, considered

as a function from moments of time to amounts of dollars. Therefore (7) could be translated into a formula expressing a property of an individual concept: the oil price concept. Formally spoken, prices are considered as functions from indices to numbers, and the same for temperatures. Numbers are considered as elements in D_e , so prices and temperatures are of type $\langle s, e \rangle$ they are individual concepts.

The technical details of the solution of the temperature paradox can be illustrated by the treatment of sentence (1). The first step of its production is application of S_5 to *be* and *ninety*. This yields (8); the corresponding translation reduces to (9).

(8) *be ninety*

(9) $\lambda x[\forall x = \textit{ninety}]$.

The next step is to combine (8) with term (10), which has (11) as translation.

(10) *the temperature*

(11) $\lambda P[\exists x \forall y[\textit{temperature}(y) \leftrightarrow x = y] \wedge \forall P(x)]$.

Since the meaning postulate for common nouns (MP2) does not hold for *temperature*, its translation (11) cannot be reduced to a formula with quantification over individuals. Combination of (8) with (11) according to S_4 yields sentence (1); the corresponding translation reduces to (12).

(12) $\exists x[\forall y[\textit{temperature}(y) \leftrightarrow x = y] \wedge \forall x = \textit{ninety}]$.

The translation of sentences (2) and (3) are respectively (13) and (14).

(13) $\exists x[\forall y[\textit{temperature}(y) \leftrightarrow x = y] \wedge \textit{rise}(x)]$

(14) $\textit{rise}(\wedge \textit{ninety})$.

From (12) and (13) it does not follow that (14) is true.

Montague's treatment of the temperature paradox has been criticized for his analysis of the notion temperature. But there are examples of the same phenomenon which are not based upon temperatures (or prices). Several examples are given by LINK (1979) and LOEBNER (1976). One of their examples is the German version of (15).

(15) *The trainer changes.*

On the reading that a certain club gets another trainer, it would not be correct to translate (15) by a formula which states that the property of

changing holds for a certain individual.

The temperature paradox (and related phenomena) explain why individual concepts are useful. But in most circumstances we want to reduce them to individuals. In the remainder of this appendix it will be investigated when such a reduction is allowed. First we will do so for translations of intransitive verbs, then for other verbs, and finally for translation of common nouns.

The only intransitive verbs in the PTQ fragment which do not express a property of an individual, but of an individual concept, are *rise* and *change*. Therefore we restrict our attention to those models of IL in which the constants corresponding with the other intransitive verbs are interpreted as expressing properties of individuals. This is expressed by the following meaning postulate.

1. Meaning Postulate 3

$$\exists M \forall x \square [\delta(x) \leftrightarrow [{}^V M]({}^V x)] \quad (M \in \text{VAR}_{\langle s, \langle e, t \rangle \rangle})$$

where $M \in \text{VAR}_{\langle s, \langle e, t \rangle \rangle}$ and δ translates any member of B_{IV} other than *rise* or *change*.

1. END

This meaning postulate states that for all involved predicates on individual concepts there is for each index an equivalent predicate on individuals. This predicate is index dependent: the set of walkers now may differ from the set of walkers yesterday. MP3 expresses the existence of such an equivalent predicate by the existential quantification $\exists M$. This M is of type $\langle s, \langle e, t \rangle \rangle$ because variables get an index independent interpretation, and as argued before, the predicate on individuals corresponding with δ has to be index dependent.

In chapter 4 section 2, the δ_* notation was introduced as an abbreviation for $\lambda u[\delta(\hat{u})]$, so as an abbreviation for those cases where it could be said that δ was applied to an individual. The above meaning postulate says that for certain constants δ the argument always is an individual, even if this is not apparent from the formula. Therefore it might be expected that MP3 allows us to introduce the δ_* notation for those constants in all contexts. The following theorems allow us to replace a formula with an occurrence of δ (where MP3 holds for δ), by a formula with an occurrence of δ_* .

2. THEOREM. MP3 is equivalent with

$$\models \Box \delta(x) \leftrightarrow \delta_*(^V x).$$

PROOF part 1. Suppose that MP3 holds, so $\models \exists M \forall x \Box [\delta(x) \leftrightarrow [^V M](^V x)]$.

Then there is a g such that $g \models \forall x \Box [\delta(x) \leftrightarrow [^V M](^V x)]$.

Now for all $g' \underset{V}{\sim} g$ $g' \models \delta_*(v) = \lambda u [\delta(^A u)](v) = \delta(^A v) = [^V M](^V v) = [^V M](v)$.

Consequently $g' \models \delta_* = ^V M$.

So there is a $g: g \models \forall x \Box [\delta(x) \leftrightarrow \delta_*(^V x)]$.

Since there are no free variables in this formula, we have

$$\models \Box [\delta(x) \leftrightarrow \delta_*(^V x)].$$

REMARK. The following more direct approach is incorrect because the conditions for λ -conversion are not satisfied.

$$g \models \delta_*(^V x) = \lambda u \delta(^A u)(^V x) = \delta(^A ^V x) = \delta(x) = [^V M](^V x).$$

PROOF part 2. Suppose $\models \Box [\delta(x) \leftrightarrow \delta_*(^V x)]$.

Let g, i be arbitrary and define $g' \underset{M}{\sim} g$ by $g'(M) = [^A \lambda u \delta(^A u)]^{A, i, g}$.

Then $i, g' \models \delta(x) \leftrightarrow [\lambda u \delta(^A u)](^V x) \leftrightarrow [^V \lambda u \delta(^V x)] \leftrightarrow [^V M](^V x)$.

Since g, i were arbitrary, MP2 follows.

2. END

On the basis of this theorem, we have besides RR_3 , another reduction rule introducing the $*$.

3. Reduction rule 11

Let be given an expression of the form $\delta(x)$, where δ is the translation of an intransitive verb other than *rise* or *change*.

Then replace $\delta(x)$ by $\delta_*(^V x)$.

CORRECTNESS PROOF

Apply theorem 2.

3. END

Now we have two rules for the introduction of δ_* : RR_3 and RR_{11} . The one requires that the argument is of a certain form, the other that the function is of a certain nature. They have different conditions for

application, and none makes the other superfluous. In case both reduction rules are applicable, they yield the same result. It is not clear to me why MP3 is formulated as it is, and not directly in the form given in theorem 2.

For verbs of other categories there are related meaning postulates. For instance the transitive verb *find* should be interpreted as a relation between individuals. The meaning postulate for the transitive verbs were already given in chapter 4 (MP4). Exceptions to that meaning postulate were *seek* and *conceive* because these verbs do not express a relation between individuals. But also about these verbs something can be said in this respect. The first arguments have to be (intensions of) individuals: it is an individual that seeks, and not an individual concept. This is expressed by meaning postulate 5, that will be given below. For verbs of other categories a related postulate expresses that their subjects are not individual concepts, but individuals.

4. Meaning postulate 5

$$\forall P \exists M \forall x \square [\delta(x, P) \leftrightarrow [{}^V M]({}^V x)]$$

where $\delta \in \{\textit{seek}, \textit{conceive}\}$.

5. Meaning postulate 6

$$\forall P \exists M \forall x \square [\delta(x, P) \leftrightarrow [{}^V M]({}^V x)]$$

where $\delta \in \{\textit{believe that}, \textit{assert that}\}$.

6. Meaning postulate 7

$$\forall P \exists M \forall x \square [\delta(x, P) \leftrightarrow [{}^V M]({}^V x)]$$

where $\delta \in \{\textit{try to}, \textit{wish to}\}$.

6. END

These three meaning postulates do not give rise to new reduction rules because there are no generally accepted notations for the corresponding predicates with an individual as first argument.

The treatment of the temperature paradox was essentially based on the use of individual concepts.

This explains why all common nouns are translated into constants denoting predicates on individual concepts. Most common nouns express a predicate on individuals. This is formulated in a meaning postulate which I recall from chapter 4.

7. Meaning postulate 2

$$\Box [\delta(x) \rightarrow \exists u[x = \hat{u}]]$$

where $\delta \in \{\text{man, woman, park, fish, pen, unicorn}\}$.

7. END

The meaning postulates for nouns and for verbs have a related aim: they both aim at excluding arbitrary individual concepts as argument and guaranteeing an individual as argument. So one might expect that there is a close relation between the consequences of the two meaning postulates. One might for instance expect that for nouns something holds like the formula in MP3. This is not the case, as is expressed in the following theorem.

8. THEOREM. Let $\delta \in \text{CON}_{\langle\langle s, e \rangle, t \rangle}$.

Let (I) be the formula $\Box [\delta(x) \rightarrow \exists u[x = \hat{u}]]$

and (II) the formula $\Box [\delta(x) \leftrightarrow \delta_*(x)]$.

Then (i) (I) $\not\Rightarrow$ (II)

and (ii) (II) $\not\Rightarrow$ (I).

PROOF. (i) In chapter 3 we introduced the constant *bigboss*, which will be used here. Suppose that

$$i_1 \models \text{bigboss} = \hat{\text{nixon}} \text{ and } i_2 \models \text{bigboss} = \hat{\text{bresjnev}}.$$

Then

$$[\text{bigboss}]^{A, i_1, g}(i_2) = \lambda i [\text{bigboss}]^{A, i_1, g}(i)(i_2) = \text{bigboss}^{A, i_2, g}(i_2) = \text{bresjnev}$$

and

$$[\text{bigboss}]^{A, i_1, g}(i_1) = \lambda i [\text{bigboss}]^{A, i_1, g}(i)(i_1) = \text{bigboss}^{A, i_1, g}(i_1) = \text{nixon}.$$

This means that ${}^{\wedge} \text{bigboss}$ is an expression of type $\langle s, e \rangle$ which does not denote a constant function. Since $[\text{bigboss}]^{A, i_1, g}(i_2) = [\text{bigboss}]^{A, i_1, g}(i_1) = \lambda_i g(i)$

we have that for no $g: g, i_1 \models \wedge u = \wedge^V \text{bigboss}$. Suppose furthermore that δ is a constant for which MP_2 holds, say man .

Then (I) is satisfied.

So $g, i_1 \models man(\wedge^V \text{bigboss}) \rightarrow \exists u[\wedge^V \text{bigboss} = \wedge u]$.

Due to the just proved property of $\wedge^V \text{bigboss}$, the consequence is never true.

So for no $g: g, i_1 \models man(\wedge^V \text{bigboss})$.

Suppose moreover that the predicate man_* holds for Nixon.

So $g, i_1 \models man_*(nixon)$.

Since $g, i_1 \models \vee \text{bigboss} = nixon$

we have $g, i_1 \models man_*(\vee \text{bigboss})$. Consequently

for no $g: g, i_1 \models man(\wedge^V \text{bigboss}) \leftrightarrow man_*(\vee \text{bigboss})$.

So if $g(x) = [\wedge^V \text{bigboss}]^A, i_1, g$ statement (II) is not true. Finally, note

that it is easy to design a model in which *bigboss* and *man* have the assumed properties. Hence we have proven (i).

PROOF. (ii) Let *bigboss* be as above. Assume now that δ is a constant for which MP_3 holds, say *walk*. Then (II) holds for δ .

Suppose now $i_1 \models walk_*(nixon)$.

So $i_2 \models walk_*(\vee \text{bigboss})$.

Let $g(x) = [\wedge^V \text{bigboss}]^A, i, g$.

Then it is not true that

$$i \models walk_*(x) \rightarrow \exists u[x = \wedge u]$$

because the antecedence is true, whereas the consequence is false. A model in which *walk* and *bigboss* have the desired properties can easily be defined and that is a counterexample to the implication.

8. END

Consequences of the above theorem are:

1. The formulations of the meaning postulates for Common Nouns and for Intransitive Verbs cannot be transformed into each other.
2. The following statement from PTQ (MONTAGUE 1973, p.265,+19) is incorrect:
 $\square [\delta(x) \leftrightarrow \delta_*(\vee x)]$ if δ translates a basic common noun other than *price* or *temperature*.
3. The meaning postulate for common nouns does not allow for replacing in all contexts an individual concept variable by the extension of this variable. This result was independently found by LINK (1979, p.224).

Next I will prove that in certain contexts the meaning postulate for common nouns does allow us to replace bound variables of type $\langle s, e \rangle$ by variables of type e . It are contexts created by these translation rules of the PTQ-fragment: the translation rule for the determiner CN and the determiner-CN-rel.clause constructions. In the sequel δ stands for the translation of a CN for which MP_2 holds, and ϕ for the translation of a relative clause. This ϕ may be omitted in the formulation of the theorems.

9. LEMMA. If $A, i, g \models \forall x \psi$ then $A, i, g \models \forall u [\wedge u/x] \psi$
 iff $A, i, g \models \exists u [\wedge u/x] \psi$ then $A, i, g \models \exists x \psi$.

PROOF. $\{m \in D_{\langle s, e \rangle} \mid m = [\wedge u]^{A, i, g}\} \subset \{m \in D_{\langle s, e \rangle} \mid m = x^{A, i, g}\}$.
 9. END

The next theorem deals with terms in which the determiner is a .

10. THEOREM. $A, i, g \models \exists x [\delta(x) \wedge \phi \wedge \forall P(x)]$
 iff $A, i, g \models \exists u [\delta(\wedge u) \wedge [\wedge u/x] \phi \wedge \forall P(\wedge u)]$.

PROOF. Suppose that

$$(1) \quad A, i, g \models \exists x [\delta(x) \wedge \phi \wedge \forall P(x)].$$

Then there is a $m \in D_{\langle s, e \rangle}$ such that

$$(2) \quad A, i, [x \rightarrow m]g \models \delta(x) \wedge \phi \wedge \forall P(x).$$

From MP_2 and (2) follows

$$(3) \quad A, i, [x \rightarrow m]g \models \exists u [x = \wedge u].$$

So there is a $a \in D_e$ such that

$$(4) \quad A, i, [x \rightarrow m, u \rightarrow a]g \models x = \wedge u.$$

From (4) and (2) follows

$$(5) \quad A, i, [x \rightarrow m, u \rightarrow a]g \models \delta(\wedge u) \wedge [\wedge u/x] \phi \wedge \forall P(\wedge u).$$

So

$$(6) \quad A, i, g \models \exists u[\delta(\wedge u) \wedge [\wedge u/x]\phi \wedge \vee P(\wedge u)].$$

Reversely (6) implies (1), as follows from the above lemma.

10. END

The terms with determiner *every* are dealt with in theorem 11.

$$11. \text{ THEOREM. } A, i, g \models \forall x[\delta(x) \wedge \phi \rightarrow \vee P(x)]$$

$$\text{iff } A, i, g \models \forall u[\delta(\wedge u) \wedge [\wedge u/x]\phi \rightarrow \vee P(\wedge u)].$$

PROOF. One direction of the theorem follows immediately from Lemma 9.

The other direction is proved by contra-position. Assume that was *not* true that

$$(1) \quad A, i, g \models \forall x[\delta(x) \wedge \phi \rightarrow \vee P(x)].$$

This means

$$(2) \quad A, i, g \models \neg \forall x[\delta(x) \wedge \phi \rightarrow \vee P(x)].$$

This is equivalent with

$$(3) \quad A, i, g \models \exists x[\delta(x) \wedge \phi \wedge \neg \vee P(x)].$$

Application of the argumentation of theorem 10 gives

$$(4) \quad A, i, g \models \exists u[\delta(\wedge u) \wedge [\wedge u/x]\phi \wedge \neg \vee P(\wedge u)].$$

Therefore it is not true that

$$(5) \quad A, i, g \models \forall u[\delta(\wedge u) \wedge [\wedge u/x]\phi \rightarrow \vee P(\wedge u)].$$

So (5) implicates (1).

11. END

The next two theorems deal with terms with determiner *the*.

12. **THEOREM.** If $A, i, g \models \exists y[\forall x[\delta(x) \wedge \phi] \leftrightarrow x=y] \wedge \forall P(y)$.
 Then $A, i, g \models \exists u[\forall v[\delta(\wedge v) \wedge [\wedge v/x]\phi] \leftrightarrow u=v] \wedge \forall P(\wedge u)$.

PROOF. Suppose

$$(1) \quad A, i, g \models \exists y[\forall x[\delta(x) \wedge \phi \leftrightarrow x=y] \wedge \forall P(y)].$$

This means that there is an $m \in D_{\langle s, e \rangle}$ such that (2) and (3) hold

$$(2) \quad A, i, [y \rightarrow m]g \models \forall x[\delta(x) \wedge \phi \leftrightarrow x=y]$$

$$(3) \quad A, i, [y \rightarrow m]g \models \forall P(y).$$

From (2) follows (4), and therefore (5) holds.

$$(4) \quad A, i, [y \rightarrow m]g \models \delta(y) \wedge [y/x]\phi \leftrightarrow y = y$$

$$(5) \quad A, i, [y \rightarrow m]g \models \delta(y) \wedge [y/x]\phi.$$

From (5) and MP_2 follows that there is an $a \in D_e$ such that (6)

$$(6) \quad A, i, [y \rightarrow m, u \rightarrow a]g \models y = \wedge u.$$

From (3) and (6) follows (7)

$$(7) \quad A, i, [y \rightarrow m, u \rightarrow a]g \models \forall P(\wedge u).$$

Apply lemma 9 to (2) and substitute $\wedge v$ for y . Since (6) holds it follows that (8) holds

$$(8) \quad A, i, [y \rightarrow m, u \rightarrow a] \models \forall v[\delta(\wedge v) \wedge [\wedge v/y]\phi \leftrightarrow \wedge v = \wedge u].$$

Since $[u=v]^{A, i, g}$ equals $[\wedge u = \wedge v]^{A, i, g}$, we may replace in (8) $\wedge v = \wedge u$ by $v = u$. Combination of (8) with (7) yields (9)

$$(9) \quad A, i, [y \rightarrow m, u \rightarrow a] \models \forall v[\delta(\wedge v) \wedge [\wedge v/y]\phi \leftrightarrow v = u] \wedge \forall P(\wedge u).$$

From this the theorem follows.

12. END

13. THEOREM. If $A, i, g \models \exists v[\forall u[[\delta(\wedge u) \wedge [\wedge u/x]\phi] \leftrightarrow u=v] \wedge \forall P(\wedge v)]$
then $A, i, g \models \exists y[\forall x[[\delta(x) \wedge \phi] \leftrightarrow x=y] \wedge \forall P(x)]$.

PROOF. Suppose

$$(1) \quad A, i, g \models \exists v[\forall u[\delta(\wedge u) \wedge [\wedge u/x]\phi \leftrightarrow u=v] \wedge \forall P(\wedge v)].$$

Then there is an $a \in D_e$ such that (2) and (3) hold

$$(2) \quad A, i, [v \rightarrow a]g \models \forall u[\delta(\wedge u) \wedge [\wedge u/x]\phi \leftrightarrow u=v]$$

$$(3) \quad A, i, [v \rightarrow a]g \models \forall P(\wedge v).$$

Let $m \in D_{\langle s, e \rangle}$ be such that (4) holds.

$$(4) \quad A, i, [x \rightarrow m]g \models \delta(x) \wedge \phi.$$

Then from MP_2 follows that there is a b such that

$$(5) \quad A, i, [x \rightarrow m, u \rightarrow b]g \models x = \wedge u.$$

From (5) and (2) follows (6)

$$(6) \quad A, i, [x \rightarrow m, v \rightarrow a, u \rightarrow b]g \models \delta(x) \wedge \phi \leftrightarrow x = \wedge v.$$

Since (4) holds, it follows from (6).

$$(7) \quad A, i, [x \rightarrow m, v \rightarrow a]g \models x = \wedge v.$$

It follows from (4) and (7) that (8) holds

$$(8) \quad A, i, [v \rightarrow a]g \models \forall x[\delta(x) \wedge \phi \rightarrow x = \wedge v].$$

Let now $m \in D_{\langle s, e \rangle}$ be such that (9) holds

$$(9) \quad A, i, [v \rightarrow a, x \rightarrow m]g \models x = \hat{\wedge} v.$$

From (2) it then follows that (10) holds

$$(10) \quad A, i, [v \rightarrow a, x \rightarrow m]g \models \delta(x) \wedge \phi.$$

From (9) and (10) follows (11)

$$(11) \quad A, i, [v \rightarrow a]g \models \forall x [x = \hat{\wedge} v \rightarrow \delta(x) \wedge \phi].$$

From (3), (8) and (11) the theorem follows.

13. END

The above theorems constitute the justification for the following reduction rule.

14. REDUCTION RULE 12

Let δ be the translation of a common noun for which meaning postulate MP_2 holds. Let be given a formula of one of the following forms,

$$\exists x [\delta(x) \wedge \phi \wedge \forall P(x)]$$

$$\forall x [\delta(x) \wedge \phi \rightarrow \forall P(x)]$$

$$\exists y [\forall x [\delta(x) \wedge \phi \leftrightarrow x=y] \wedge \forall P(y)].$$

Then replace this formula by respectively

$$\exists u [\delta(\hat{\wedge} u) \wedge [\hat{\wedge} u/x] \phi \wedge \forall P(\hat{\wedge} u)]$$

$$\forall u [\delta(\hat{\wedge} u) \wedge [\hat{\wedge} u/x] \phi \rightarrow \forall P(\hat{\wedge} u)]$$

$$\exists v [\forall u [\delta(\hat{\wedge} u) \wedge [\hat{\wedge} u/x] \phi \leftrightarrow u=v] \wedge \forall P(\hat{\wedge} v)]$$

(provided that ϕ does not contain a free occurrence of u or v).

CORRECTNESS PROOF

See the theorems.

14. END

The theorems mentioned above, allow us to change the types of bound variables in a lot of contexts which arise if one deals with sentences from the PTQ-fragment. But they do *not* cover all contexts arising in this fragment. If the rule of quantification into a CN phrase (i.e. $S_{15,n}$) is used, then no reduction rule is applicable. An example is (14) in the reading in which *every* has wider scope than *a*. The corresponding translation is (15), and although none of the reduction rules is applicable, it is equivalent with (16).

(14) *Every man such that he loses a pen such that he finds it, runs.*

(15) $\forall x[\exists u[\text{pen}_*(u) \wedge \text{man}(x) \wedge \text{lose}_*(x,u) \wedge \text{find}_*(x,u)] \rightarrow \text{run}_*(x)]$

(16) $\forall v[\exists u[\text{pen}_*(u) \wedge \text{man}_*(v) \wedge \text{lose}_*(v,u) \wedge \text{find}_*(v,u)] \rightarrow \text{run}_*(v)].$

One would like to have a reduction rule which is applicable to constructions in which quantification into a CN is used. However, not in all such contexts reduction is possible. This was discovered by FRIEDMAN & WARREN (1980a). Consider sentence (17)

(17) *A unicorn such that every woman loves it changes.*

Suppose that 17 is obtained by quantification of *every woman* into *unicorn such that he₁ loves it*. Then the translation of (17) reduces to (18); Friedman & Warren call this 'a rather unusual reading'.

(18) $\exists x[\forall u[\text{woman}_*(u) \rightarrow \text{unicorn}(x) \wedge \text{love}_*(u,x) \wedge \text{change}(x)]]$.

This translation is, however not equivalent with (19).

(19) $\exists v[\forall u[\text{woman}_*(u) \rightarrow \text{unicorn}_*(v) \wedge \text{love}_*(u,v) \wedge \text{change}_*(v)]]$.

This situation might rise doubts about rule $S_{15,n}$. However, see chapter 9, section 7.2 for an example where the rule is needed.

APPENDIX 2

SET MANIPULATION IN SYNTAX

In chapter 6 I provided a system of categories for dealing with syntactic variables. The rules given there implicitly assume that the reader knows what sets are, and what \cup , with and $-$ mean. This is set theoretical knowledge, and not knowledge of the grammatical system. In the present section I will formulate syntactic rules which allow for replacing expressions like $\{1,2\} - 1$ by $\{2\}$ and $\{1,2\} \cup 3$ by $\{1,2,3\}$. So we aim at rules which remove the symbols \cup , with and $-$ from the formulation of the rules. The collection of rules performing this task is rather complex. I wish to emphasize that the rules do not arise from the requirement of using total rules, but from grammatical formalism. A related situation would arise when using partial rules. Such rules would mention a condition like 'contains an occurrence of he_n '. Since 'containing' is not a notion defined by grammatical means, a formalist might wish to do so. Then rules are needed which are related to the rules below since they have to perform related tasks. Once it is shown that the set-theoretical notions \cup , with and $-$ are definable by means of grammatical tools, there is no objection against using them in the grammar even when not explicitly defined in this way.

Let G be a grammar with a collection hyperrules H , and let the elements of H contain expressions like $set - n$. Then the actual rules of the grammar are defined as the result of performing the following actions in order.

1. replace the metavariables in the hyperrules by some terminal metaproduction of the meta-grammar
2. replace subexpressions in the rules by other expressions, according to the rules given below, until there are no occurrences of the non-acceptable symbols (\cup , with, $-$).

The rules eliminating the non acceptable symbols introduce some non-acceptable symbols themselves. These are $+$, is, unless, true and false; these symbols have to be added to those mentioned in point 2 above. The collection of rules performing the task of eliminating these symbols is infinite, and will be defined by means of a two-level grammar. The hyperrules describing the elimination of the unacceptable symbols are unrestricted rewriting rules with metavariables. These variables are mentioned below, together with some examples of their terminal productions. Different examples are

separated by a bar symbol: /, and ϵ denotes the empty string.

set : {1,2} / {3,1} / \emptyset .
seq : 1,2 / 3,1 .
lseq: 1, / 3,1, / ϵ .
rseq: ,2 / ,1,5 / ϵ .
n : 1 / 5.

The metarules for these metavariables are as follows (again a bar / separates alternatives, the non-terminal symbols are in italics).

set \rightarrow *seq* / \emptyset .
seq \rightarrow *n* / *n*, *seq*.
n \rightarrow 1/2/3/4/5/6/7/8/9/*m*/*n*0.
lseq \rightarrow *seq*, / ϵ .
rseq \rightarrow ,*seq* / ϵ .

The rules for with have to allow for replacing {1,2} with 1 by {1,2}, whereas they should not allow for replacing {2,3} with 1 by {2,3}. The hyperrule describing such replacements is

{*lseq*,*n*,*rseq*} with *n* \rightarrow {*lseq*,*n*,*rseq*}.

An example of a rule derived from this hyperrule is

{1,3,5} with 3 \rightarrow {1,3,5}.

Thus the expression with 3 is eliminated. In case one meets the subexpression {2,3} with 1 there is no rule which can be obtained from this hyperrule and which can be applied to this subexpression. So we cannot get rid of the non-acceptable symbol with, as was required in point 2. So we do not obtain an actual rule and the derivation cannot continue. This 'blind alley' technique is due to SINTZOFF (1967).

The rules for eliminating the - sign have to replace {1,2} - 2 by {1}, and {1,2} - 3 by {1,2}. The rules have to check whether the number preceded by the - sign occurs in the set mentioned before the sign. For this purpose, we need grammatical means to check whether two numbers are equal or different. It is easy to design a rule which can be applied only if two numbers are equal: a hyperrule with two occurrences of the meta-variable *n* can be transformed into a real rule only by substituting for both occurrences

the same number. If a hyperrule contains metavariables n_1 and n_2 , then it can be transformed into a real rule by substituting for n_1 and n_2 different numbers. But nothing prevents us to substitute the same number. It is difficult to guarantee that two numbers are different, but we need such rules. The rules which do so use the blind alley technique again, now on the symbol unless. The hyperrules are as follows.

$$\begin{array}{lll}
 0 \text{ is } 0 \rightarrow \text{true} & 1 \text{ is } 0 \rightarrow \text{false} & 2 \text{ is } 0 \rightarrow \text{false} \quad \dots \\
 0 \text{ is } 1 \rightarrow \text{false} & 1 \text{ is } 1 \rightarrow \text{true} & 2 \text{ is } 1 \rightarrow \text{false} \\
 \vdots & \vdots & \vdots \\
 0 \text{ is } 9 \rightarrow \text{false} & 1 \text{ is } 9 \rightarrow \text{false} & \\
 n_1 \text{ is } n_2 \rightarrow n_1 \text{ is } n_2 & & 2n_1 \text{ is } n_2 \rightarrow \text{false} \\
 n_1 \text{ is } 2n_2 \rightarrow \text{false} & & 2n_1 \text{ is } 2n_2 \rightarrow n_1 \text{ is } n_2 \\
 \vdots & & \\
 n_1 \text{ is } 9n_2 \rightarrow \text{false} & \dots & \\
 \text{unless true} \rightarrow \text{false} & & \\
 \text{unless false} \rightarrow \text{true} & & \\
 \text{true} \rightarrow \epsilon. & &
 \end{array}$$

The above rules have the effect that an expression of the form unless a is b reduces to unless true and next to unless in case a equals b. This unless constitutes a blind alley. If a is not equal to b, the expression reduces to unless false and through true to the empty string. Then the test is eliminated, and the production may proceed.

The rules for the - sign have to check for all elements of the mentioned set whether they are equal to the element that has to be removed. The element for which equality is tested (in a step of the testing process) is the last element of the sequence describing the set. If equality is found, the element is removed. If a check shows that the numbers are different, then the element which has been checked, is put at the beginnings of the sequence, and the new 'last element' is checked. By means of the * sign the numbers are separated which are already checked from the numbers which are not yet checked. This rotation technique is due to Van WIJNGAARDEN (1974). The hyperrules introducing and removing the * sign are as follows.

$$\begin{aligned}
\{seq\} - n &\rightarrow \{*, seq\} - n \\
\emptyset - n &\rightarrow \emptyset \\
\{seq, *\} - n &\rightarrow \{seq\} \\
\{*\} - n &\rightarrow \emptyset.
\end{aligned}$$

The hyperrule removing an element is

$$\{lseq * rseq, n\} - n \rightarrow \{lseq * rseq\} - n.$$

The rule rotating the sequence is

$$\{lseq * rseq, n_1\} - n_2 \rightarrow \{n_1, lseq * rseq\} - n_2 \text{ unless } n_1 \text{ is } n_2.$$

We use the unless phrase to guarantee that n_1 and n_2 are different. If the numbers are different, then the phrase reduces to the empty string. If they are equal the unless phrase reduces to the expression unless, and we cannot get rid of this phrase. This means that we are in a blind alley: we do not get an actual rule.

The rules for \cup use the $-$ sign. It would be easy to reduce $\{1,2\} \cup \{2\}$ to $\{1,2,2\}$ but, in order to avoid this repetition of elements, I first remove the 2 from the leftmost set and then add 2 to the set thus obtained. The rules are as follows.

$$\begin{aligned}
set \cup \{n, rseq\} &\rightarrow set - n + n \cup \{rseq\} \\
\{seq\} + n &\rightarrow \{seq, n\} \\
\emptyset + n &\rightarrow n \\
set \cup \{\emptyset\} &\rightarrow set \\
set \cup \{ \} &\rightarrow set.
\end{aligned}$$

This completes the description of the set of rules needed for dealing with set-theoretical notions by grammatical means.

INDEX OF NAMES

Andreka	87
Bach	119, 120, 122, 124, 125, 133, 138, 144-147, 150, 151, 157, 164, 167, 168, 178, 180-183
Bartsch	43, 54, 61, 95
Bennett	129, 154, 177, 178
van Benthem	11, 77
Bresnan	112, 141
Burmeister	87
Chomsky	102, 103, 131, 138
Cooper	54, 86, 87, 119, 120, 137, 138, 141, 144-158, 166, 167
Curry	122
Delacruz	57
Dik	118
Dowty	43, 57, 69, 95, 122, 124, 177
Ejerhed	192, 193,
Emde Boas	29, 75
Faltz	71, 72, 77, 78
van Fraassen	86
Friedman	7, 22, 24, 37, 118, 131, 132, 167, 221
Gallin	74, 76
Gazdar	21, 103
Goguen	87
Goldblatt	87
Graetzer	87
Groenendijk	70, 71, 76, 124, 128, 143, 150, 158
Hausser	75, 76, 86, 87, 111, 124, 141, 156
Henkin	74
Hopcroft	84
Hopkins	184
Indurkha	7
Janssen	7, 27, 54, 75, 89, 94, 118, 138 144, 155, 157, 178, 206
Kamp	87, 193
Karttunen	166
Keenan	71, 72, 77
Klein	87, 131
von Kutschera	87
Kripke	5
Ladusaw	192
Landman	33, 121, 125
Landsbergen	107, 122
Lewis	10, 48, 76
Link	210
Loebner	210
Lukasiewicz	87
ter Meulen	87
Mikenberg	87
Moerdijk	121, 125
Montague	1, 2, 4, 58, 62, 78, 124, 150, 174, 215

Nemeti	87
Oh	158
Parsons	54, 119
Partee	25, 27, 49, 56, 57, 73, 80, 82, 91-99, 108-112, 117-120, 125, 129, 131, 132, 138, 140, 141, 155, 156, 163, 164, 206
Rodman	60, 141, 143, 147, 157
Rohrer	177,
Saarinen	193
Sag	103
Scha	85
Scott	87
Sintzoff	224
Stokhof	70, 71, 76, 124, 128, 143, 150, 158, 176
Strachey	87
Thomason	63, 82, 84, 94, 143, 155, 159
Tichy	75
Veltman	91
Waldo	85, 87
Warren	7, 37, 167, 221
van Wijngaarden	103, 225

REFERENCES

- Andreka, H., P. Burmeister & I. Nemeti, 1980,
'Quasivarieties of partial algebras. A unifying approach
towards a two-values model theory for partial algebras',
Preprint 557, Fachbereich Mathematics, Technische Hochschule,
Darmstadt.
- Andreka, H., & I. Nemeti, 1982,
'Generalization of variety and quasi variety-concept
to partial algebras through category theory',
Dissertationes Mathematicae 204.
- Bach, E. & R.H. Cooper, 1978,
'The NP-S analysis of relative clauses and compositional
semantics',
Linguistics and Philosophy 2, 145-150.
- Bach, E., 1979a,
'Control in Montague grammar',
Linguistic Inquiry 10, 515-531.
- Bach, E., 1979b,
'Montague grammar and classical transformational grammar',
in Davis & Mithun 1979, pp. 3-49.
- Bach, E., 1980,
'Tenses and aspects as functions on verb-phrases',
in Rohrer 1980, pp. 19-37.
- Bartsch, R., 1978,
'Infinitives and the control problem in categorial grammar',
Theoretical Linguistics 5, 217-250.
- Bartsch, R., 1979,
'The syntax and semantics of subordinate clause
constructions and pronominal reference',
in Heny & Schnelle 1979, pp. 23-59.
- Bennett, M., 1976,
'A variation and extension of a Montague fragment of English',
in Partee 1976, pp. 119-163.
- Bennett, M., 1977,
'A guide to the logic of tense and aspect in English',
Logique et Analyse 80, 491-517.
- Bennett, M., 1978,
'Demonstratives and indexicals in Montague grammar',
Synthese 39, 1-80.
- Bentham, J.F.A.K. van, 1977,
'Tense logic and standard logic',
Logique et Analyse 80, 395-437.
Kennis en methode 5, 94-116.
- Bentham, J.F.A.K. van, 1981,
'Why is semantics what',
in Groenendijk, Janssen & Stokhof 1981, pp. 24-49.
- Bresnan, J.W., 1973,
'Comparative deletion and constraints on transformations',
Linguistic Analysis 1, 25-74.
- Chomsky, N., 1965,
'Aspects of the theory of syntax',
The M.I.T. Press, Cambridge (Mass.)

- Chomsky, N., 1975,
 'Questions of form and interpretation',
 Linguistic Analysis 1,
 Reprinted in N. Chomsky, 'Essays on form and interpretation',
 North Holland, New York, 1977, pp. 25-29.
- Cooper, R.H., 1975,
 'Montague's semantic theory and transformational syntax',
 dissertation, Univ. of Massachusetts, Amherst.
 Published by: Xerox University Microfilms.
- Cooper, R. & T. Parsons, 1976,
 'Montague grammar, generative semantics and interpretative
 semantics',
 in Partee, 1976, pp. 311-362.
- Cooper, R.H., 1978,
 'A fragment of English with questions and relative clauses',
 Unpublished paper, Dept. of Linguistics, Univ. of Wisconsin.
- Cooper, R.H., 1979a
 'The interpretation of pronouns',
 in Heny & Schnelle 1979, pp. 61-92.
- Cooper, R., 1979b,
 'Variable binding and relative clauses',
 in F. Guenther & J.S. Schmidt (eds), 'Formal semantics and
 pragmatics for natural languages', Synthese language library 4,
 Reidel, Dordrecht, 1979, pp. 131-171.
- Curry, H.B., 1961,
 'Some logical aspects of grammatical structure',
 in 'Structure of language and its mathematical aspects',
 Proceedings of Symposia in Applied Mathematics vol. XII,
 American Mathematical Society, Rhode Island, pp. 56-68.
- Davidson, D. & G. Harman (eds), 1972,
 'Semantics of natural language',
 Synthese library 40, Reidel, Dordrecht.
- Davis, S. & M. Mithun (eds), 1979,
 'Linguistics, philosophy and Montague grammar. (Proc. conf.
 Albany 1977)',
 Univ. of Texas Press, Austin, Texas.
- Delacruz, E.B., 1976,
 'Factives and propositional level constructions in Montague
 grammar',
 in Partee 1976, pp.177-199.
- Dik, S.C., 1978,
 'Functional grammar',
 North Holland Linguistics series 37, North Holland,
 Amsterdam.
- Dik, S.C., 1980,
 'Seventeen Sentences: basic principles and application
 of functional grammar',
 in E.A. Moravcsik & J.R. Wirth (eds), 'Current approaches to
 syntax', Syntax and semantics 13, Academic Press, 1980, pp. 45-75.
- Dowty, D.R., 1976,
 'Montague grammar and the lexical decomposition of causative
 verbs',
 in Partee 1976, pp. 201-245.

- Dowty, D., 1978,
 'Governed transformations as lexical rules in a
 Montague grammar',
 Linguistic Inquiry 9, 393-426.
- Dowty, D., 1979a,
 'Dative "movement" and Thomason's extensions of Montague
 grammar',
 in Davis & Mithun 1979, pp. 153-222.
- Dowty, D.R., 1979b,
 'Word meaning and Montague grammar',
 Synthese language library 7, Reidel, Dordrecht.
- Dowty, D., 1982,
 'Grammatical relations and Montague grammar',
 in Jacobson & Pullum 1982, pp. 79-130.
- Ejerhed, E.I., 1981,
 'Tense as a source of intensional ambiguity',
 in Heny 1981, pp. 231-252.
- Fraassen, B.C. van, 1969,
 'Presuppositions, supervaluations and free logic',
 in K. Lambert (ed.), 'The logical way of doing things',
 Yale Univ. Press, New Haven, 1979, pp. 67-91.
- Friedman, J., 1979a,
 'An unlabelled bracketing solution to the problem
 of conjoined phrases in Montague's PTQ',
 Journal of Philosophical Logic 8, 151-169.
- Friedman, J. & D. Warren, 1979b,
 'Erratum',
 Linguistics and Philosophy 3, 39.
- Friedman, J., & D. Warren, 1980a,
 'Notes on an intensional logic for English III. Extensional
 forms',
 Report N-13, Dept. of Computer and Communication Sciences,
 The Univ. of Michigan, Ann Arbor (Mich).
- Friedman, J. & D. Warren, 1980b,
 'Lambda-normal forms in an intensional logic for English',
 Studia Logica 39, 311-324.
- Gallin, D., 1975,
 'Intensional and higher-order modal logic',
 Mathematics studies 17, North Holland, Amsterdam.
- Gazdar, G., 1980,
 'A cross categorial semantics for coordination',
 Linguistics and Philosophy 3, 407-409.
- Gazdar, G. & I.A. Sag, 1981,
 'Passives and reflexives in phrase structure grammar',
 in Groenendijk, Janssen & Stokhof, 1981, pp. 131-152.
- Gazdar, G., 1982,
 'Phrase structure grammar',
 in Jakobson & Pullum 1982, pp. 131-186.
- Goldblatt, R., 1979,
 'Topoi. The categorial analysis of logic',
 Studies in Logic and the foundations of mathematics 98,
 North Holland, Amsterdam.
- Goguen, J., 1978,
 'Abstract errors for abstract data types',
 in Neuhold 1978, pp. 491-525

- Graetzer G., 1968,
 'Universal algebra',
 The univ. series in higher mathematics, van Nostrand, Princeton.
 Second edition published by: Springer, New York, 1979.
- Groenendijk, J. & M. Stokhof, 1976,
 'Some notes on personal pronouns, reflexives and sloppy
 identity',
 in K. Braunmueller & W. Kuerschner (eds), 'Grammatik. Akten
 des 10. linguistischen Kolloquiums. Band 2', Max Niemeyer,
 Tuebingen, 1976, pp. 301-319.
- Groenendijk, J. & M. Stokhof, 1979,
 'Infinitives and context in Montague grammar',
 in Davis & Mithun 1979, pp. 287-310.
- Groenendijk J.A.G., T.M.V. Janssen & M.B.J. Stokhof (eds), 1981,
 'Formal methods in the study of language. Proceedings of the
 third Amsterdam colloquium',
 MC-Tracts 135 & 136, Mathematical Centre, Amsterdam, 1981.
- Groenendijk, J. & M. Stokhof, 1981,
 'Semantics of wh-complements',
 in Groenendijk, Janssen, Stokhof, 1981, pp.153 -181.
- Hausser, R.R., 1976,
 'Presuppositions in Montague grammar',
 Theoretical linguistics 3, 245-280
- Hausser, R.R., 1978,
 'Surface compositionality and the semantics of mood',
 in Groenendijk & Stokhof 1978, pp. 174-193.
- Hausser, R.R., 1979a,
 'A constructive approach to intensional contexts. Remarks
 on the metaphysics of model theory',
 Unpublished paper.
- Hausser, R.R., 1979b,
 'How do pronouns denote?',
 in Heny & Schnelle 1979, pp. 93-139.
- Hausser, R.R., 1980,
 'Surface compositionality and the semantics of mood',
 in J.R. Searle, F. Kiefer & M. Bierwisch, 'Speech act theory and
 pragmatics', Reidel, Dordrecht, 1980, pp. 71-95.
- Hausser, R.R., 1984,
 'Surface compositional grammar',
 'Studies in theoretical linguistics, Fink Verlag, Muenchen.
- Henkin, L., 1950,
 'Completeness in the theory of types',
 Journal of Symbolic Logic 15, 81-91.
- Heny, F. & H.S. Schnelle (eds), 1979,
 'Selections from the third Groningen round table',
 Syntax and Semantics 10, Academic Press, New York.
- Heny, F. (ed.), 1981,
 'Ambiguities in intensional contexts',
 Synthese language library 12, Reidel, Dordrecht.
- Hopcroft, J.E. & J.D. Ullman, 1979,
 'Introduction to automata theory, languages and computation',
 Addison-Wesley, Reading (Mass.).

- Hopkins, M., 1972,
 'A case for the GOTO',
 in 'Proceedings of the ACM (proc. conf. Boston 1972)',
 Association for Computing Machinery, New York, 1972, pp. 787-790.
- Indurkha, B., 1981,
 'Sentence analysis programs based on Montague grammars',
 unpublished paper, Philips international institute
 of technological studies, Eindhoven.
- Jacobson, P., & G.K. Pullum, 1982,
 'The nature of syntactic representation',
 Synthese Language Library 15, Reidel, Dordrecht.
- Janssen, T.M.V. & P. van Emde Boas, 1977a,
 'On the proper treatment of referencing, dereferencing
 and assignment',
 in A. Salomaa & M. Steinby (eds), 'Automata, languages,
 and programming (Proc. 4th. coll. Turku)', Lecture notes in
 computer science 52, Springer, Berlin, 1977, pp. 282-300.
- Janssen, T., 1978a,
 'Compositionality and the form of the rules in Montague
 grammar',
 in Groenendijk & Stokhof 1978, pp. 101-124.
- Janssen, T.M.V., 1978b,
 'Simulation of a Montague grammar',
 Annals of systems research 6, 127-140.
- Janssen, T.M.V., 1980a,
 'Logical investigations on PTQ arising from programming
 requirements',
 Synthese 44, 361-390.
- Janssen, T.M.V., 1980b,
 'On problems concerning the quantification rules in
 Montague grammar',
 in Rohrer 1980, pp. 113-134.
- Janssen, T.M.V., 1981,
 'Compositional semantics and relative clause formation
 in Montague grammar',
 in Groenendijk, Janssen, Stokhof 1981, pp. 237-276.
- Janssen, T.M.V., 1981b,
 'Montague grammar and functional grammar',
 in T. Hoekstra & H v.d. Hulst and M. Moortgat (eds),
 'Perspectives on functional grammar', Foris publications,
 Dordrecht, 273-297.
 also: GLOT 3, 1980, 273-297.
- Kamp, H., 1971,
 'Formal properties of "Now"',
 Theoria 37, 227-273.
- Kamp, H., 1975,
 'Two theories about adjectives',
 in Keenan 1975, pp. 123-155.
- Keenan, E. (ed.), 1975,
 'Formal semantics of natural language. (Coll. Cambridge 1973)',
 Cambridge Univ. Press, 1975.
- Keenan, E.L. & L.M. Faltz, 1978,
 'Logical types for natural language',
 UCLA occasional papers in linguistics 3.

- Klein, E., 1979,
 On sentences which report beliefs, desires and other
 mental attitudes',
 unpublished dissertation.
- Klein, E., 1981,
 'The interpretation of adjectival, nominal and
 adverbial constructions',
 in Groenendijk, Janssen, Stokhof 1981, pp. 381-398.
- Kripke, S., 1972,
 'Naming and necessity',
 in Davidson & Harman 1972, pp. 253-355.
- Kutschera, F. von, 1975,
 'Partial interpretations',
 in Keenan 1975, pp. 156-174.
- Ladusaw, W., 1974,
 'Some problems with tense in PTQ',
 Texas Linguistics Forum 1, p. 89-102.
- Landman, F. & I. Moerdijk, 1981,
 'Morphological features and conditions on rules in
 Montague grammar',
 Amsterdam papers in formal grammar 3, Centrale Interfaculteit
 University of Amsterdam.
- Landman, F., & I. Moerdijk, 1983a,
 'Compositionality and the analysis
 Linguistics and Philosophy 6, pp. 89-114.
- Landsbergen, J., 1981,
 'Adaption of Montague grammar to the requirements of parsing',
 in Groenendijk, Janssen, Stokhof 1981, pp. 399-420.
- Lewis, D., 1970,
 'General semantics',
 Synthese 22, 18-67.
 Reprinted in Davidson & Harman 1972, pp. 169-248.
 Reprinted in Partee 1976, pp. 1-50.
- Lewis, D., 1974,
 'Tensions',
 in M.K. Munitz & P.K. Unger (eds), 'Semantics and
 philosophy', New York Univ. Press, New York, 1974.
- Link, G., 1979,
 'Montague Grammatik. I: Die logische Grundlagen',
 Kritische Information 71, Wilhelm Fink, Muenchen.
- Loebner, S., 1976,
 'Einfuehrung in die Montague Grammatik',
 Monographien Linguistik und Kommunikationswissenschaft 27,
 Scriptor, Kronberg.
- Lukasiewicz, J., 1920,
 'Philosophische Bemerkungen zu mehrwertigen Systemen
 des Aussagenkalkuels',
 Comptes Rendus des Sceances de Varsovie, Cl. iii, 23, pp. 51-77.
 Translated by H. Weber as 'Philosophical remarks on many-valued
 systems of propositional logic', in S. McCall (ed.), 'Polish
 logic, 1920-1939', Clarendon Press, Oxford, 1967.
- Meulen, A. ter, 1980,
 'Substances, quantities and individuals. A study in
 the formal semantics of mass terms',
 distributed by: Indiana Univ. Linguistics Club,
 Bloomington (Ind.).

- Mikenberg, I., 1977,
 'From total to partial algebras',
 in A.I. Arruda, N.C.A. da Costa & R. Chuaqui (eds),
 Mathematical Logic. Proceedings of the first Brazilian
 Conference', M. Dekker, New York, 1977, pp. 203-223.
- Montague, R., 1968,
 'Pragmatics',
 in R. Klibansky (ed.), 'Contemporary philosophy. A survey',
 La Nuovo Italia Editrice, Florence, 1968, pp. 102-122.
 Reprinted in Thomason 1974, pp. 95-118.
- Montague, R., 1970b,
 'Universal grammar',
 Theoria 36, 373-398.
 Reprinted in : Thomason 1974, pp. 222-246.
- Montague, R., 1973,
 'The proper treatment of quantification in ordinary
 English',
 in K.J.J. Hintikka, J.M.E. Moravcsik & P. Suppes (eds),
 'Approaches to natural language', Synthese Library 49,
 Reidel, Dordrecht, 1973, pp. 221-242.
 Reprinted in Thomason 1974, pp. 247-270.
- Oh, Choon-Kyu, 1977,
 'The so-called Bach-Peters paradox explained away',
 Theoretical Linguistics 4, 197-207.
- Partee, B.H., 1971,
 'On the requirement that transformations preserve meaning',
 in C. Fillmore & D.T. Langendoen (eds), 'Studies in
 linguistic semantics', Holt, Rinehart & Wintson, 1971.
- Partee, B.H., 1973,
 'Some transformational extensions of Montague grammar',
 Journal of Philosophical Logic 2, 509-534.
 Reprinted in Partee 1976, pp. 51-76.
- Partee, B., 1975,
 'Montague grammar and transformational grammar',
 Linguistic Inquiry 6, 203-300.
- Partee, B.H. (ed.), 1976,
 'Montague grammar',
 Academic Press, New York.
- Partee, B., 1977a,
 'John is easy to please',
 in A. Zampolli (ed.), 'Linguistic structures processing',
 Fundamental studies in computer science 5, North Holland,
 Amsterdam, 1977, pp. 281-312.
- Partee, B.H., 1977b,
 'Possible world semantics and linguistic theory',
 The Monist 60, 303-326.
- Partee, B.H., 1979a,
 'Constraining transformational Montague grammar: A framework
 and a fragment',
 in Davis & Mithun 1979, pp. 51-102.
- Partee, B.H., 1979b,
 'Montague grammar and the well-formedness constraint',
 in Heny & Schnelle 1979, pp. 275-314.

- Partee, B. & E. Bach, 1981,
 'Quantification, pronouns and VP-anaphora',
 in J.A.G. Groenendijk, T.M.V. Janssen, & M.B.J. Stokhof, 1981,
 'Formal methods in the study of language. Proceeding of the third
 Amsterdam colloquium. Part 2', MC Tract 136, Mathematical Centre,
 Amsterdam, pp. 445-481.
 reprinted in J. Groenendijk, T.M.V. Janssen & M. Stokhof (eds),
 'Truth, interpretation and information', Grass 3, Foris,
 Dordrecht, 1984, 99-130.
- Rodman, R., 1976,
 'Scope phenomena, "movement transformations", and relative
 clauses',
 in Partee 1976, pp. 165-176.
 Gunter Narr, Tuebingen, 1977.
- Rohrer, C., 1980,
 'Time, tense and quantifiers. (Proc. conf Stuttgart 1979)',
 Linguistische Arbeiten 83, Max Niemeyer, Tuebingen.
- Saarinen, E., 1978,
 'Backwards-looking operators in tense logic and in natural
 language',
 in J. Hintikka, I. Niiniluoto, & E. Saarinen (eds), 'Essays
 on mathematical and philosophical logic. Proceedings of the
 4th. Scandinavian logic symposium', Synthese Library 122,
 Reidel, Dordrecht, 1978, pp.341-367.
- Scott, D. & C. Strachey 1971,
 'Towards a mathematical semantics for computer languages',
 in J. Fox (ed.), 'Computers and automata (proc. symp.
 Brooklyn)', Polytechnic Press, Brooklyn (N.Y.), 1971,
 pp. 16-46.
- Sintzoff, M., 1967,
 'Existence of a van Wijgaarden syntax for every
 enumerable set',
 Annales de la Societe Scientifique de Bruxelles 81, 115-118.
- Thomason, R.H., 1972,
 'A semantic theory of sortal incorrectness',
 Journal of Philosophical Logic 1, 209-258.
- Thomason, R.H. (ed.), 1974,
 'Formal philosophy. Selected papers of Richard Montague',
 Yale University Press, New Haven.
- Thomason, R.H., 1976,
 'Some extensions of Montague grammar',
 in Partee 1976, pp. 77-117.
- Tichy, P., 1971,
 'An approach to intensional analysis',
 Nous 5, 273-297.
- Veltman F., 1981,
 'Data semantics',
 in Groenendijk, Janssen & Stokhof 1981, pp. 541-565.
 Revised reprint in Groenendijk, Janssen & Stokhof, 1984,
 pp. 43-63.
- Visentini et al., 1970,
 'Linguaggi nella societa e nella tecnica',
 Edizioni di communita, Milan (distributed by the Olivetti
 Corporation).

- Waldo, J., 1979,
 'A PTQ semantics for sortal incorrectness',
 in Davis & Mithun 1979, pp. 311-331.
- Wijngaarden, A. van, 1970,
 'On the boundary between natural and artificial languages',
 in Visentini et al., 1970, pp. 165-176.
- Wijngaarden, A. van, et al., 1975,
 'Revised report on the algorithmic language ALGOL 68',
 Acta Informatica 5, 1-236.

CWI TRACTS

- 1 D.H.J. Epema. *Surfaces with canonical hyperplane sections*. 1984.
- 2 J.J. Dijkstra. *Fake topological Hilbert spaces and characterizations of dimension in terms of negligibility*. 1984.
- 3 A.J. van der Schaft. *System theoretic descriptions of physical systems*. 1984.
- 4 J. Koene. *Minimal cost flow in processing networks, a primal approach*. 1984.
- 5 B. Hoogenboom. *Intertwining functions on compact Lie groups*. 1984.
- 6 A.P.W. Böhm. *Dataflow computation*. 1984.
- 7 A. Blokhuis. *Few-distance sets*. 1984.
- 8 M.H. van Hoorn. *Algorithms and approximations for queueing systems*. 1984.
- 9 C.P.J. Koymans. *Models of the lambda calculus*. 1984.
- 10 C.G. van der Laan, N.M. Temme. *Calculation of special functions: the gamma function, the exponential integrals and error-like functions*. 1984.
- 11 N.M. van Dijk. *Controlled Markov processes; time-discretization*. 1984.
- 12 W.H. Hundsdorfer. *The numerical solution of nonlinear stiff initial value problems: an analysis of one step methods*. 1985.
- 13 D. Grune. *On the design of ALEPH*. 1985.
- 14 J.G.F. Thiemann. *Analytic spaces and dynamic programming: a measure theoretic approach*. 1985.
- 15 F.J. van der Linden. *Euclidean rings with two infinite primes*. 1985.
- 16 R.J.P. Groothuizen. *Mixed elliptic-hyperbolic partial differential operators: a case-study in Fourier integral operators*. 1985.
- 17 H.M.M. ten Eikelder. *Symmetries for dynamical and Hamiltonian systems*. 1985.
- 18 A.D.M. Kester. *Some large deviation results in statistics*. 1985.
- 19 T.M.V. Janssen. *Foundations and applications of Montague grammar, part 1: Philosophy, framework, computer science*. 1986.
- 20 B.F. Schriever. *Order dependence*. 1986.
- 21 D.P. van der Vecht. *Inequalities for stopped Brownian motion*. 1986.
- 22 J.C.S.P. van der Woude. *Topological dynamix*. 1986.
- 23 A.F. Monna. *Methods, concepts and ideas in mathematics: aspects of an evolution*. 1986.
- 24 J.C.M. Baeten. *Filters and ultrafilters over definable subsets of admissible ordinals*. 1986.
- 25 A.W.J. Kolen. *Tree network and planar rectilinear location theory*. 1986.
- 26 A.H. Veen. *The misconstrued semicolon: Reconciling imperative languages and dataflow machines*. 1986.
- 27 A.J.M. van Engelen. *Homogeneous zero-dimensional absolute Borel sets*. 1986.
- 28 T.M.V. Janssen. *Foundations and applications of Montague grammar, part 2: Applications to natural language*. 1986.

MATHEMATICAL CENTRE TRACTS

- 1 T. van der Walt. *Fixed and almost fixed points*. 1963.
- 2 A.R. Bloemena. *Sampling from a graph*. 1964.
- 3 G. de Leve. *Generalized Markovian decision processes, part I: model and method*. 1964.
- 4 G. de Leve. *Generalized Markovian decision processes, part II: probabilistic background*. 1964.
- 5 G. de Leve, H.C. Tijms, P.J. Weeda. *Generalized Markovian decision processes, applications*. 1970.
- 6 M.A. Maurice. *Compact ordered spaces*. 1964.
- 7 W.R. van Zwet. *Convex transformations of random variables*. 1964.
- 8 J.A. Zonneveld. *Automatic numerical integration*. 1964.
- 9 P.C. Baayen. *Universal morphisms*. 1964.
- 10 E.M. de Jager. *Applications of distributions in mathematical physics*. 1964.
- 11 A.B. Paalman-de Miranda. *Topological semigroups*. 1964.
- 12 J.A.Th.M. van Berckel, H. Brandt Corstius, R.J. Mokken, A. van Wijngaarden. *Formal properties of newspaper Dutch*. 1965.
- 13 H.A. Lauwerier. *Asymptotic expansions*. 1966, out of print; replaced by MCT 54.
- 14 H.A. Lauwerier. *Calculus of variations in mathematical physics*. 1966.
- 15 R. Doornbos. *Slippage tests*. 1966.
- 16 J.W. de Bakker. *Formal definition of programming languages with an application to the definition of ALGOL 60*. 1967.
- 17 R.P. van de Riet. *Formula manipulation in ALGOL 60, part 1*. 1968.
- 18 R.P. van de Riet. *Formula manipulation in ALGOL 60, part 2*. 1968.
- 19 J. van der Slot. *Some properties related to compactness*. 1968.
- 20 P.J. van der Houwen. *Finite difference methods for solving partial differential equations*. 1968.
- 21 E. Wattel. *The compactness operator in set theory and topology*. 1968.
- 22 T.J. Dekker. *ALGOL 60 procedures in numerical algebra, part 1*. 1968.
- 23 T.J. Dekker, W. Hoffmann. *ALGOL 60 procedures in numerical algebra, part 2*. 1968.
- 24 J.W. de Bakker. *Recursive procedures*. 1971.
- 25 E.R. Paërl. *Representations of the Lorentz group and projective geometry*. 1969.
- 26 European Meeting 1968. *Selected statistical papers, part I*. 1968.
- 27 European Meeting 1968. *Selected statistical papers, part II*. 1968.
- 28 J. Oosterhoff. *Combination of one-sided statistical tests*. 1969.
- 29 J. Verhoeff. *Error detecting decimal codes*. 1969.
- 30 H. Brandt Corstius. *Exercises in computational linguistics*. 1970.
- 31 W. Molenaar. *Approximations to the Poisson, binomial and hypergeometric distribution functions*. 1970.
- 32 L. de Haan. *On regular variation and its application to the weak convergence of sample extremes*. 1970.
- 33 F.W. Steutel. *Preservation of infinite divisibility under mixing and related topics*. 1970.
- 34 I. Juhász, A. Verbeek, N.S. Kroonenberg. *Cardinal functions in topology*. 1971.
- 35 M.H. van Emden. *An analysis of complexity*. 1971.
- 36 J. Grasman. *On the birth of boundary layers*. 1971.
- 37 J.W. de Bakker, G.A. Blaauw, A.J.W. Duijvestijn, E.W. Dijkstra, P.J. van der Houwen, G.A.M. Kamsteeg-Kemper, F.E.J. Kruseman Aretz, W.L. van der Poel, J.P. Schaap-Kruseman, M.V. Wilkes, G. Zoutendijk. *MC-25 Informatica Symposium*. 1971.
- 38 W.A. Verloren van Themaat. *Automatic analysis of Dutch compound words*. 1972.
- 39 H. Bavinck. *Jacobi series and approximation*. 1972.
- 40 H.C. Tijms. *Analysis of (s,S) inventory models*. 1972.
- 41 A. Verbeek. *Superextensions of topological spaces*. 1972.
- 42 W. Vervaat. *Success epochs in Bernoulli trials (with applications in number theory)*. 1972.
- 43 F.H. Ruymgaart. *Asymptotic theory of rank tests for independence*. 1973.
- 44 H. Bart. *Meromorphic operator valued functions*. 1973.
- 45 A.A. Balkema. *Monotone transformations and limit laws*. 1973.
- 46 R.P. van de Riet. *ABC ALGOL, a portable language for formula manipulation systems, part 1: the language*. 1973.
- 47 R.P. van de Riet. *ABC ALGOL, a portable language for formula manipulation systems, part 2: the compiler*. 1973.
- 48 F.E.J. Kruseman Aretz, P.J.W. ten Hagen, H.L. Oudshoorn. *An ALGOL 60 compiler in ALGOL 60, text of the MC-compiler for the EL-X8*. 1973.
- 49 H. Kok. *Connected orderable spaces*. 1974.
- 50 A. van Wijngaarden, B.J. Mailloux, J.E.L. Peck, C.H.A. Koster, M. Sintzoff, C.H. Lindsey, L.G.L.T. Meertens, R.G. Fisker (eds.). *Revised report on the algorithmic language ALGOL 68*. 1976.
- 51 A. Hordijk. *Dynamic programming and Markov potential theory*. 1974.
- 52 P.C. Baayen (ed.). *Topological structures*. 1974.
- 53 M.J. Faber. *Metricizability in generalized ordered spaces*. 1974.
- 54 H.A. Lauwerier. *Asymptotic analysis, part 1*. 1974.
- 55 M. Hall, Jr., J.H. van Lint (eds.). *Combinatorics, part 1: theory of designs, finite geometry and coding theory*. 1974.
- 56 M. Hall, Jr., J.H. van Lint (eds.). *Combinatorics, part 2: graph theory, foundations, partitions and combinatorial geometry*. 1974.
- 57 M. Hall, Jr., J.H. van Lint (eds.). *Combinatorics, part 3: combinatorial group theory*. 1974.
- 58 W. Albers. *Asymptotic expansions and the deficiency concept in statistics*. 1975.
- 59 J.L. Mijnheer. *Sample path properties of stable processes*. 1975.
- 60 F. Göbel. *Queueing models involving buffers*. 1975.
- 63 J.W. de Bakker (ed.). *Foundations of computer science*. 1975.
- 64 W.J. de Schipper. *Symmetric closed categories*. 1975.
- 65 J. de Vries. *Topological transformation groups. I: a categorical approach*. 1975.
- 66 H.G.J. Pijls. *Logically convex algebras in spectral theory and eigenfunction expansions*. 1976.
- 68 P.P.N. de Groen. *Singularly perturbed differential operators of second order*. 1976.
- 69 J.K. Lenstra. *Sequencing by enumerative methods*. 1977.
- 70 W.P. de Roever, Jr. *Recursive program schemes: semantics and proof theory*. 1976.
- 71 J.A.E.E. van Nunen. *Contracting Markov decision processes*. 1976.
- 72 J.K.M. Jansen. *Simple periodic and non-periodic Lamé functions and their applications in the theory of conical waveguides*. 1977.
- 73 D.M.R. Leivant. *Absoluteness of intuitionistic logic*. 1979.
- 74 H.J.J. te Riele. *A theoretical and computational study of generalized aliquot sequences*. 1976.
- 75 A.E. Brouwer. *Tree-like spaces and related connected topological spaces*. 1977.
- 76 M. Rem. *Associons and the closure statement*. 1976.
- 77 W.C.M. Kallenberg. *Asymptotic optimality of likelihood ratio tests in exponential families*. 1978.
- 78 E. de Jonge, A.C.M. van Rooij. *Introduction to Riesz spaces*. 1977.
- 79 M.C.A. van Zuijlen. *Empirical distributions and rank statistics*. 1977.
- 80 P.W. Hemker. *A numerical study of stiff two-point boundary problems*. 1977.
- 81 K.R. Apt, J.W. de Bakker (eds.). *Foundations of computer science II, part 1*. 1976.
- 82 K.R. Apt, J.W. de Bakker (eds.). *Foundations of computer science II, part 2*. 1976.
- 83 L.S. van Benthem Jutting. *Checking Landau's "Grundlagen" in the AUTOMATH system*. 1979.
- 84 H.L.L. Busard. *The translation of the elements of Euclid from the Arabic into Latin by Hermann of Carinthia (?), books vii-xii*. 1977.
- 85 J. van Mill. *Supercompactness and Wallman spaces*. 1977.
- 86 S.G. van der Meulen, M. Veldhorst. *Torrix I, a programming system for operations on vectors and matrices over arbitrary fields and of variable size*. 1978.
- 88 A. Schrijver. *Matroids and linking systems*. 1977.
- 89 J.W. de Roever. *Complex Fourier transformation and analytic functionals with unbounded carriers*. 1978.

- 90 L.P.J. Groenewegen. *Characterization of optimal strategies in dynamic games*. 1981.
- 91 J.M. Geysel. *Transcendence in fields of positive characteristic*. 1979.
- 92 P.J. Weeda. *Finite generalized Markov programming*. 1979.
- 93 H.C. Tijms, J. Wessels (eds.). *Markov decision theory*. 1977.
- 94 A. Bijlsma. *Simultaneous approximations in transcendental number theory*. 1978.
- 95 K.M. van Hee. *Bayesian control of Markov chains*. 1978.
- 96 P.M.B. Vitányi. *Lindenmayer systems: structure, languages, and growth functions*. 1980.
- 97 A. Federgruen. *Markovian control problems; functional equations and algorithms*. 1984.
- 98 R. Geel. *Singular perturbations of hyperbolic type*. 1978.
- 99 J.K. Lenstra, A.H.G. Rinnooy Kan, P. van Emde Boas (eds.). *Interfaces between computer science and operations research*. 1978.
- 100 P.C. Baayen, D. van Dulst, J. Oosterhoff (eds.). *Proceedings bicentennial congress of the Wiskundig Genootschap, part 1*. 1979.
- 101 P.C. Baayen, D. van Dulst, J. Oosterhoff (eds.). *Proceedings bicentennial congress of the Wiskundig Genootschap, part 2*. 1979.
- 102 D. van Dulst. *Reflexive and superreflexive Banach spaces*. 1978.
- 103 K. van Harn. *Classifying infinitely divisible distributions by functional equations*. 1978.
- 104 J.M. van Wouwe. *Go-spaces and generalizations of metrizability*. 1979.
- 105 R. Helmers. *Edgeworth expansions for linear combinations of order statistics*. 1982.
- 106 A. Schrijver (ed.). *Packing and covering in combinatorics*. 1979.
- 107 C. den Heijer. *The numerical solution of nonlinear operator equations by imbedding methods*. 1979.
- 108 J.W. de Bakker, J. van Leeuwen (eds.). *Foundations of computer science III, part 1*. 1979.
- 109 J.W. de Bakker, J. van Leeuwen (eds.). *Foundations of computer science III, part 2*. 1979.
- 110 J.C. van Vliet. *ALGOL 68 transput, part I: historical review and discussion of the implementation model*. 1979.
- 111 J.C. van Vliet. *ALGOL 68 transput, part II: an implementation model*. 1979.
- 112 H.C.P. Berbee. *Random walks with stationary increments and renewal theory*. 1979.
- 113 T.A.B. Snijders. *Asymptotic optimality theory for testing problems with restricted alternatives*. 1979.
- 114 A.J.E.M. Janssen. *Application of the Wigner distribution to harmonic analysis of generalized stochastic processes*. 1979.
- 115 P.C. Baayen, J. van Mill (eds.). *Topological structures II, part 1*. 1979.
- 116 P.C. Baayen, J. van Mill (eds.). *Topological structures II, part 2*. 1979.
- 117 P.J.M. Kallenberg. *Branching processes with continuous state space*. 1979.
- 118 P. Groeneboom. *Large deviations and asymptotic efficiencies*. 1980.
- 119 F.J. Peters. *Sparse matrices and substructures, with a novel implementation of finite element algorithms*. 1980.
- 120 W.P.M. de Ruyter. *On the asymptotic analysis of large-scale ocean circulation*. 1980.
- 121 W.H. Haemers. *Eigenvalue techniques in design and graph theory*. 1980.
- 122 J.C.P. Bus. *Numerical solution of systems of nonlinear equations*. 1980.
- 123 I. Yuhász. *Cardinal functions in topology - ten years later*. 1980.
- 124 R.D. Gill. *Censoring and stochastic integrals*. 1980.
- 125 R. Eising. *2-D systems, an algebraic approach*. 1980.
- 126 G. van der Hoek. *Reduction methods in nonlinear programming*. 1980.
- 127 J.W. Klop. *Combinatory reduction systems*. 1980.
- 128 A.J.J. Talman. *Variable dimension fixed point algorithms and triangulations*. 1980.
- 129 G. van der Laan. *Simplicial fixed point algorithms*. 1980.
- 130 P.J.W. ten Hagen, T. Hagen, P. Klint, H. Noot, H.J. Sint, A.H. Veen. *ILP: intermediate language for pictures*. 1980.
- 131 R.J.R. Back. *Correctness preserving program refinements: proof theory and applications*. 1980.
- 132 H.M. Mulder. *The interval function of a graph*. 1980.
- 133 C.A.J. Klaassen. *Statistical performance of location estimators*. 1981.
- 134 J.C. van Vliet, H. Wupper (eds.). *Proceedings international conference on ALGOL 68*. 1981.
- 135 J.A.G. Groenendijk, T.M.V. Janssen, M.J.B. Stokhof (eds.). *Formal methods in the study of language, part I*. 1981.
- 136 J.A.G. Groenendijk, T.M.V. Janssen, M.J.B. Stokhof (eds.). *Formal methods in the study of language, part II*. 1981.
- 137 J. Telgen. *Redundancy and linear programs*. 1981.
- 138 H.A. Lauwerier. *Mathematical models of epidemics*. 1981.
- 139 J. van der Wal. *Stochastic dynamic programming, successive approximations and nearly optimal strategies for Markov decision processes and Markov games*. 1981.
- 140 J.H. van Geldrop. *A mathematical theory of pure exchange economies without the no-critical-point hypothesis*. 1981.
- 141 G.E. Welters. *Abel-Jacobi isogenies for certain types of Fano threefolds*. 1981.
- 142 H.R. Bennett, D.J. Lutzer (eds.). *Topology and order structures, part 1*. 1981.
- 143 J.M. Schumacher. *Dynamic feedback in finite- and infinite-dimensional linear systems*. 1981.
- 144 P. Eijgenraam. *The solution of initial value problems using interval arithmetic; formulation and analysis of an algorithm*. 1981.
- 145 A.J. Brentjes. *Multi-dimensional continued fraction algorithms*. 1981.
- 146 C.V.M. van der Mee. *Semigroup and factorization methods in transport theory*. 1981.
- 147 H.H. Tigelaar. *Identification and informative sample size*. 1982.
- 148 L.C.M. Kallenberg. *Linear programming and finite Markovian control problems*. 1983.
- 149 C.B. Huijsmans, M.A. Kaashoek, W.A.J. Luxemburg, W.K. Vietsch (eds.). *From A to Z, proceedings of a symposium in honour of A.C. Zaanen*. 1982.
- 150 M. Veldhorst. *An analysis of sparse matrix storage schemes*. 1982.
- 151 R.J.M.M. Does. *Higher order asymptotics for simple linear rank statistics*. 1982.
- 152 G.F. van der Hoeven. *Projections of lawless sequences*. 1982.
- 153 J.P.C. Blanc. *Application of the theory of boundary value problems in the analysis of a queueing model with paired services*. 1982.
- 154 H.W. Lenstra, Jr., R. Tijdeman (eds.). *Computational methods in number theory, part I*. 1982.
- 155 H.W. Lenstra, Jr., R. Tijdeman (eds.). *Computational methods in number theory, part II*. 1982.
- 156 P.M.G. Apers. *Query processing and data allocation in distributed database systems*. 1983.
- 157 H.A.W.M. Kneppers. *The covariant classification of two-dimensional smooth commutative formal groups over an algebraically closed field of positive characteristic*. 1983.
- 158 J.W. de Bakker, J. van Leeuwen (eds.). *Foundations of computer science IV, distributed systems, part 1*. 1983.
- 159 J.W. de Bakker, J. van Leeuwen (eds.). *Foundations of computer science IV, distributed systems, part 2*. 1983.
- 160 A. Rezus. *Abstract AUTOMATH*. 1983.
- 161 G.F. Helminck. *Eisenstein series on the metaplectic group, an algebraic approach*. 1983.
- 162 J.J. Dik. *Tests for preference*. 1983.
- 163 H. Schippers. *Multiple grid methods for equations of the second kind with applications in fluid mechanics*. 1983.
- 164 F.A. van der Duyn Schouten. *Markov decision processes with continuous time parameter*. 1983.
- 165 P.C.T. van der Hoeven. *On point processes*. 1983.
- 166 H.B.M. Jonkers. *Abstraction, specification and implementation techniques, with an application to garbage collection*. 1983.
- 167 W.H.M. Zijm. *Nonnegative matrices in dynamic programming*. 1983.
- 168 J.H. Evertse. *Upper bounds for the numbers of solutions of diophantine equations*. 1983.
- 169 H.R. Bennett, D.J. Lutzer (eds.). *Topology and order structures, part 2*. 1983.