

# On the Bahadur-Efficient Testing of Uniformity by means of the Entropy

Peter Harremoës, *Member, IEEE*, and Igor Vajda, *Fellow, IEEE*,

**Abstract**— This paper compares the power divergence statistics of orders  $\alpha > 1$  with the information divergence statistic in the problem of testing the uniformity of a distribution. In this problem the information divergence statistic is equivalent to the entropy statistic. Extending some previously established results about information diagrams, it is proved that the information divergence statistic in this problem is more efficient in the Bahadur sense than any power divergence statistic of order  $\alpha > 1$ . This means that the entropy provides in this sense the most efficient way of characterizing the uniformity of a distribution.

**Index Terms**— Bahadur efficiency, entropy, goodness-of-fit, index of coincidence, information diagram, power divergences.

## I. POWER DIVERGENCE STATISTICS

LET  $M(k)$  denote the set of all discrete probability distributions of the form  $P = (p_1, \dots, p_k)$  and

$$M(k|n) = \{P \in M(k) : nP \in \{0, 1, \dots\}^k\} \quad (1)$$

the subset of types. One of the fundamental problems of mathematical statistics can be described by  $n$  balls distributed into boxes  $1, \dots, k$  independently according to an unknown probability law

$$P_n = (p_{n1}, \dots, p_{nk}) \in M(k) \quad (2)$$

possibly depending on the number of balls  $n$ . This results in frequency counts  $X_{n1}, \dots, X_{nk}$  the vector of which  $\mathbf{X}_n = (X_{n1}, \dots, X_{nk}) \in \{0, 1, \dots\}^k$  has a multinomial distribution with parameters  $k, n, P_n$ ,

$$\mathbf{X}_n \sim \text{Multinomial}_k(n, P_n). \quad (3)$$

The problem is to decide on the basis of observations  $\mathbf{X}_n$  whether the unknown law (2) is equal to a given  $Q = (q_1, \dots, q_k) \in M(k)$  or not.

The observations  $\mathbf{X}_n$  are represented by the (random) empirical probability distribution

$$\hat{P}_n = (\hat{p}_{n1} \triangleq X_{n1}/n, \dots, \hat{p}_{nk} \triangleq X_{nk}/n) \in M(k|n) \quad (4)$$

and the hypothesis  $Q$  about  $P_n$  is usually decided by means of a procedure  $\mathcal{T}$  called a test. This procedure uses a statistic  $T_n(\hat{P}_n, Q)$  which characterizes the goodness-of-fit between

Manuscript was submitted Feb. 2007. P. Harremoës was supported by grants from the Willum Kann Rasmussen Foundation and Danish Natural Science Research Council. I. Vajda was supported by the GAČR grant 102/07/1131 and the MŠMT grant 1M0572.

P. Harremoës is with Centrum voor Wiskunde en Informatica, Amsterdam, NL-1090 GB, The Netherlands (e-mail: P.Harremoës@cwi.nl).

I. Vajda is with Inst. Inform. Theory and Automation, Acad. Sciences Czech Republic, CZ-182 Prague, Czech Republic (e-mail: vajda@utia.cas.cz)

the distributions  $\hat{P}_n$  and  $Q$ . The test  $\mathcal{T}$  rejects the hypothesis  $P_n = Q$  if  $T = T_n(\hat{P}_n, Q)$  exceeds certain rejection level  $r_n \in \mathbb{R}$ .

The goodness-of-fit statistic is usually one of the *power divergence statistic*

$$T_\alpha = T_{\alpha,n} = 2n D_\alpha(\hat{P}_n, Q), \quad \alpha \in \mathbb{R} \quad (5)$$

where  $D_\alpha(P, Q)$  denotes the so-called  $\alpha$ -divergence (*power divergence* of order  $\alpha$ ) of distributions  $P, Q \in M(k)$  defined by

$$D_\alpha(P, Q) = \sum_{j=1}^k q_j \phi_\alpha\left(\frac{p_j}{q_j}\right), \quad \alpha \in \mathbb{R}, \quad (6)$$

for the power function  $\phi_\alpha$  of order  $\alpha \in \mathbb{R}$  given in the domain  $t > 0$  by the formula

$$\phi_\alpha(t) = \frac{t^\alpha - \alpha(t-1) - 1}{\alpha(\alpha-1)} \quad \text{when } \alpha(\alpha-1) \neq 0 \quad (7)$$

and by the corresponding limits

$$\phi_0(t) = -\ln t + t - 1, \quad (8)$$

$$\phi_1(t) = t \ln t - t + 1. \quad (9)$$

For details about the definition (6) and properties of power divergences, see [1] or [2]. Next we cite the best known members of the family of statistics (5) with a reference to the skew symmetry  $D_\alpha(P, Q) = D_{1-\alpha}(Q, P)$  of the power divergences (6).

*Example 1: The quadratic divergences*

$$D_2(P, Q) = D_{-1}(Q, P) = \frac{1}{2} \sum_{j=1}^k \frac{(p_j - q_j)^2}{q_j}$$

lead to the well known *Pearson statistic*

$$T_2 = T_{2,n} = \sum_{j=1}^k \frac{(X_{nj} - nq_j)^2}{nq_j} \quad (10)$$

and *Neyman statistics*

$$T_{-1} = T_{-1,n} = \sum_{j=1}^k \frac{(X_{nj} - nq_j)^2}{X_{nj}}.$$

*The logarithmic divergences*

$$D_1(P, Q) = D_0(Q, P) = \sum_{j=1}^k p_j \ln \frac{p_j}{q_j} \quad (11)$$

lead to the *log-likelihood ratio statistic*

$$T_1 = T_{1,n} = 2 \sum_{j=1}^k X_{nj} \ln \frac{X_{nj}}{nq_j} \quad (12)$$

and *reversed log-likelihood ratio statistic*

$$T_0 = T_{0,n} = 2nq_j \sum_{j=1}^k \ln \frac{nq_j}{X_{nj}}.$$

The symmetric *Hellinger divergence*

$$D_{1/2}(P, Q) = D_{1/2}(Q, P) = 4 \sum_{j=1}^k (\sqrt{p_j} - \sqrt{q_j})^2$$

leads to the *Freeman–Tukey statistic*

$$T_{1/2} = T_{1/2,n} = 2 \sum_{j=1}^k \left( \sqrt{X_{nj}} - \sqrt{nq_j} \right)^2. \quad (13)$$

In this paper we would like to find the power divergence statistic  $T_\alpha$ ,  $\alpha \in \mathbb{R}$  that is most suitable for testing the hypothesis that the true distribution  $P_n$  is uniform, i.e. the hypothesis  $\mathcal{H} : P_n = U \triangleq (1/k, \dots, 1/k)$ . Hence in our model

$$\mathbf{X}_n \sim \text{Multinomial}_k(n, U) \quad \text{under } \mathcal{H}. \quad (14)$$

The alternative to the hypothesis  $\mathcal{H}$  is denoted by  $\mathcal{A}_n$ . Thus by (3),

$$\mathbf{X}_n \sim \text{Multinomial}_k(n, P_n) \quad \text{under } \mathcal{A}_n \quad (15)$$

for  $P_n$  assumed in (2).

Next follows a typical example of the hypotheses testing model introduced in (14) - (15).

*Example 2:* Let  $\mu, \nu$  be two different probability measures on the Borel line  $(\mathbb{R}, \mathcal{B})$  with absolutely continuous distribution functions  $F, G$  and  $Y_1, \dots, Y_n$  an i.i.d. sample from the probability space  $(\mathbb{R}, \mathcal{B}, \mu)$ . Consider a statistician who knows neither the probability measure  $\mu$  governing the random sample  $(Y_1, \dots, Y_n)$  nor this sample itself. Nevertheless he observes the frequencies  $\mathbf{X}_n = (X_{n1}, \dots, X_{nk})$  of the samples  $Y_1, \dots, Y_n$  in an interval partition  $\mathcal{P}_n = \{A_{n1}, \dots, A_{nk}\}$  of  $\mathbb{R}$  chosen by him. Using  $\mathbf{X}_n$  he has to decide about the hypothesis  $\mathcal{H}$  that the unknown probability measure on  $(\mathbb{R}, \mathcal{B})$  is the given  $\nu$ . Thus for a partition  $\mathcal{P}_n = \{A_{n1}, \dots, A_{nk}\}$  under his control he obtains the observations

$$\mathbf{X}_n \sim \text{Multinomial}_k(n, P_n) \quad (16)$$

where

$$P_n = (\mu(A_{n1}), \dots, \mu(A_{nk}))$$

and his task is to test the hypothesis  $\mathcal{H} : \mu = \nu$ . Knowing  $\nu$ , he can use the quantile function  $G^{-1}$  of  $\nu$  or, more precisely, the quantiles  $G^{-1}(j/k)$  of the orders  $j/k$  for  $1 \leq j \leq k$  cutting  $\mathbb{R}$  into a special system of intervals  $\mathcal{P}_n = \{A_{n1}, \dots, A_{nk}\}$  with the property  $\nu(A_{nj}) = 1/k$  for  $1 \leq j \leq k$ . Hence for this special partition we get from (16)

$$P_n = U = (1/k, \dots, 1/k) \in M(k|n) \quad \text{under } \mathcal{H} \quad (17)$$




Fig. 1. The domain of a Cauchy distribution divided in five bins with equal probabilities.

and

$$P_n = (\mu(A_{n1}), \dots, \mu(A_{nk})) \in M(k) \quad \text{under } \mathcal{A}_n. \quad (18)$$

We see from (16) - (18) that the quantile-generated partitions  $\mathcal{P}_n$  lead exactly to the situation assumed in (14) - (15). This idea is illustrated in Figure 1.

The formulas for divergences  $D_\alpha(P, Q)$  simplify when  $Q = U$ , e.g.,

$$D_1(P, U) = \ln k - H(P) \quad \text{for } P \in M(k) \quad (19)$$

where  $H(P)$  denotes the Shannon entropy

$$H(P) = - \sum_{j=1}^k p_j \ln p_j.$$

Similarly, (6) and (7) imply for all  $\alpha > 1$  and  $P \in M(k)$

$$\begin{aligned} D_\alpha(P, U) &= \frac{\sum_{j=1}^k p_j^\alpha (1/k)^{1-\alpha} - 1}{\alpha(\alpha - 1)} \\ &= \frac{k^{\alpha-1} \sum_{j=1}^k p_j^\alpha - 1}{\alpha(\alpha - 1)} \\ &= \frac{k^{\alpha-1} IC_\alpha(P) - 1}{\alpha(\alpha - 1)}. \end{aligned} \quad (20)$$

Here

$$IC_\alpha(P) = \sum_{j=1}^k p_j^\alpha \quad \text{for } P \in M(k) \quad (21)$$

is the *index of coincidence* of  $P$  of order  $\alpha > 1$  introduced in [3], taking on values between  $k^{1-\alpha}$  (when  $P$  is the uniform distribution  $U$ ) and 1 (when  $P$  is the Dirac distribution  $P_j$  with  $p_j = 1$  for some  $1 \leq j \leq k$ ).

From (19) we see that the log-likelihood ratio statistics statistic  $T_{1,n} = 2nD_1(\hat{P}_n, Q)$  is one-one related to the entropy statistic  $2nH(\hat{P}_n)$ , and from (20) we see that for each  $\alpha > 1$

the power divergence statistic  $T_{\alpha,n} = 2nD_\alpha(\hat{P}_n, Q)$  is one-one related to the corresponding  $IC$ -statistic  $2nIC_\alpha(\hat{P}_n)$ . The entropy  $H(\hat{P}_n)$  as well as the indices of coincidence  $IC_\alpha(\hat{P}_n)$ ,  $\alpha > 1$  characterize the uniformity of the distribution  $\hat{P}_n$ . We are interested in the characterization which is most efficient from the statistical point of view.

The rest of the paper is organized as follows. In Section II the basic idea behind Bahadur efficiency is explained and previous result related to efficiency of certain tests are mentioned. These previous results have been an important inspiration for developing the results of this paper, but they are not used directly. In Section III conditions for the plug-in estimator of power divergence to be consistent are given. These conditions play an important role in the formulation and the proof of the main results, but the results should also be of independent interest. Section IV is the most technical where Bahadur functions are introduced and the link to result from [3] is established.

Section V states the main result of the paper and proves it using results of the previous sections. This result is then discussed in Section VI.

## II. BAHADUR EFFICIENCY

In the previous section we introduced the family of power divergence statistics  $T_\alpha$ , or the one-one related family of statistics  $D_\alpha(\hat{P}_n, Q)$ ,  $\alpha \in \mathbb{R}$ . In the rest of this paper we are interested in the relative asymptotic efficiencies of these statistics for  $1 \leq \alpha_1 < \alpha_2 < \infty$  when applied in testing the uniformity hypothesis (14). To this end we use the concept of *Bahadur asymptotic relative efficiency* of  $T_{\alpha_1}$  with respect to  $T_{\alpha_2}$  (briefly *Bahadur efficiency*, in symbols  $BE(T_{\alpha_1} | T_{\alpha_2})$ ). Roughly speaking, this efficiency compares the sample sizes  $n_i$  needed by the  $T_{\alpha_i}$ -tests of the same powers to achieve the same asymptotic sizes. It differs from the *Pitman asymptotic relative efficiency* of  $T_{\alpha_1}$  with respect to  $T_{\alpha_2}$  which compares the sample sizes  $n_i$  needed by the  $T_{\alpha_i}$ -tests of the same sizes to achieve the same asymptotic powers (cf. [4, pp. 332–341] or [5]). We use the general concept of Bahadur efficiency introduced in [6] where it was extended the original concept of [7]. Before its formal definition, we briefly review some useful preliminary testing results.

Let us first suppose that  $k$  remains fixed while  $n$  tends to infinity. In this case the goodness-of-fit statistic (5)–(13) have been studied systematically in [2]. They proved under  $\mathcal{H}$  the limit law

$$T_{\alpha,n} \xrightarrow{\mathcal{L}} \chi_{k-1}^2 \quad \text{as } n \rightarrow \infty, \quad \alpha \in \mathbb{R} \quad (22)$$

where  $\chi_{k-1}^2$  stands for the  $\chi^2$ -distributed random variable with  $k-1$  degrees of freedom and where  $\rightarrow$  here denotes convergence in distribution. In [2] the authors also proved a modification of (22) under the local alternatives

$$\mathcal{A}_n : P_n = (1 - 1/\sqrt{n})U + P/\sqrt{n} \quad \text{for } P \in M(k) \text{ fixed.} \quad (23)$$

An extension of (22) to the case where (14) remains valid but

$k = k_n$  increases slowly to  $\infty$  as  $n \rightarrow \infty$  in the sense

$$\lim_{n \rightarrow \infty} \gamma_n = 0 \quad \text{for } \gamma_n = \frac{k}{n} \quad (24)$$

has been studied for  $\alpha = 2$  in [8] and for arbitrary positive integers  $\alpha$  in [9].

The asymptotic normality

$$\frac{T_{\alpha,n} - k}{\sqrt{2k}} \xrightarrow{\mathcal{L}} N(0, 1) \quad \text{as } n \rightarrow \infty, \quad \alpha \in \mathbb{R} \quad (25)$$

has been proved under  $\mathcal{H}$  subsequently in [10], [11] and [12] under stronger alternatives to the slow convergence condition (24), namely

$$\lim_{n \rightarrow \infty} \gamma_n = 0 \quad \text{for } \gamma_n = \frac{k^2 \ln^2 n}{n}, \quad \frac{k^2 \ln k}{n} \quad \text{and} \quad \frac{k^2}{n} \quad (26)$$

respectively. Extension of (25) to a local alternative of the type (23) can be found for  $\alpha = 1$  and  $\alpha = 2$  in [9], and for arbitrary  $\alpha \in \mathbb{R}$  in [13].

If contrary to (24) or (26),  $k = k_n$  increases fast to  $\infty$  in the sense

$$\lim_{n \rightarrow \infty} \gamma_n = \gamma > 0 \quad \text{for } \gamma_n = \frac{k}{n} \quad (27)$$

then (25) has to be replaced by more complicated limit laws established in [14], [15], [2] and [16]. However, the practical situations where the model satisfies (27) are rare. In our introductory example with distribution of balls, this assumption means that the number of boxes is comparable with the number of balls. Hence either the frequencies  $X_{n1}, \dots, X_{nk}$  of balls in all boxes remain bounded as  $n \rightarrow \infty$ , or majority of the boxes remains empty.

In what follows we restrict ourselves to the usual situations where  $k = k_n$  satisfies the conditions of the type (24) or (26). The limit laws mentioned above enable us to specify for any  $\alpha \in \mathbb{R}$  the  $T_\alpha$ -based test of the hypothesis  $\mathcal{H}$  of an arbitrary asymptotic size  $s \in ]0, 1[$ . Under the normal law (25) such a test is defined by the rejection rule

$$T_\alpha > r_n(s) \quad \text{for } r_n(s) = k_n + \sqrt{2k_n} \Phi^{-1}(1-s) \quad (28)$$

for the quantile of the order  $1-s$  of the standard normal distribution function  $\Phi$ . We would like to choose the optimal statistic  $T_{\alpha_{\text{opt}}}$  from the family  $T_\alpha$ ,  $\alpha \in \mathbb{R}$ . This leads to the comparison of the asymptotic relative efficiencies in this family.

If  $k = k_n$  increases slowly as assumed in (24) or (26), then the *Pitman asymptotic relative efficiencies* of all statistics  $T_\alpha$ ,  $\alpha \in \mathbb{R}$  coincide (cf. e.g. [2]). In this situation preferences between these statistics must be based on the Bahadur efficiencies  $BE(T_{\alpha_1} | T_{\alpha_2})$ . The key result in this direction was established in [6] where it was demonstrated that  $BE(T_1 | T_2) = \infty$  so that the log-likelihood ratio statistic  $T_1$  is more Bahadur efficient than the Pearson statistic  $T_2$ . Using the results from [17], this first achievement was extended in [18] where it was proved that the Bahadur efficiencies of the reversed log-likelihood ratio statistic  $T_0$  and the Neyman statistic  $T_{-1}$  coincide and both are less Bahadur efficient than the Pearson's  $T_2$ . A problem left open in the previous literature is

to evaluate the Bahadur efficiencies of the remaining statistics  $T_\alpha$ ,  $\alpha \in \mathbb{R}$ , in particular to confirm or reject the conjecture that the log-likelihood ratio statistic  $T_1$  is most Bahadur efficient in the class of all power divergence statistics  $T_\alpha$ ,  $\alpha \in \mathbb{R}$ . In this paper we present a partial solution to this problem for  $\alpha \geq 1$ . Our solution is based on the results for indices of coincidence established in [3].

The above mentioned Bahadur efficiency  $BE(T_{\alpha_1} | T_{\alpha_2})$  is defined under the condition that for  $\alpha = \alpha_1$  and  $\alpha = \alpha_2$  the statistic  $D_\alpha(\hat{P}_n, U)$  is consistent and admits the so-called Bahadur function. These two concepts are given in Definition 1 and 2 below. In what follows we often use the statistics  $D_\alpha(\hat{P}_n, U)$  instead of the one-one related  $T_\alpha = T_{\alpha, n}$ . Further by  $P(B_n)$  we shall denote the probabilities of events  $B_n$  depending on the random observations  $\mathbf{X}_n$  (cf. (14) and (15)) and by  $E$  the corresponding expectations.

*Definition 1:* For  $\alpha \in \mathbb{R}$  we say that

- 1) the model satisfies the Bahadur condition if there exists  $0 < \Delta_\alpha < \infty$  such that under the alternatives  $\mathcal{A}_n$

$$\lim_{n \rightarrow \infty} D_\alpha(P_n, U) = \Delta_\alpha, \quad (29)$$

- 2) the statistic  $D_\alpha(\hat{P}_n, U)$  is consistent if the Bahadur condition holds and for  $n \rightarrow \infty$

$$ED_\alpha(\hat{P}_n, U) \longrightarrow 0 \quad \text{under } \mathcal{H} \quad (30)$$

and

$$D_\alpha(\hat{P}_n, U) \xrightarrow{p} \Delta_\alpha \quad \text{under } \mathcal{A}_n. \quad (31)$$

The inequality  $0 < \Delta_\alpha < \infty$  in the Bahadur condition means that in term of the statistic  $D_\alpha(\hat{P}_n, U)$ , the alternatives  $\mathcal{A}_n$  are neither too near to nor too far from the hypothesis  $\mathcal{H}$ . The next example demonstrates that in the model of Example 2 this important condition holds.

*Example 3:* Let us consider the typical situation of Example 2 leading to the present statistical testing model. If the probability measure  $\mu$  considered there is dominated by  $\nu$  then, by [19, Theorem 2],

$$\lim_{n \rightarrow \infty} D_\alpha(P_n, U) = \int_{-\infty}^{\infty} \phi_\alpha \left( \frac{d\mu}{d\nu} \right) d\nu \quad \text{for all } \alpha \in \mathbb{R}.$$

The integrals are  $\alpha$ -divergences  $D_\alpha(\mu, \nu)$  of probability measures  $\mu$  and  $\nu$ , see [1]. Thus (29) holds for  $\Delta_\alpha = D_\alpha(\mu, \nu)$  when  $\mu$  is dominated by  $\nu$  and  $\Delta_\alpha > 0$  unless  $\mu = \nu$  (i.e.  $\mathcal{H} = \mathcal{A}_n$  for all  $n = 1, 2, \dots$ ). This means that if the model of Example 2 is nontrivial then then the Bahadur condition holds for all  $\alpha \in \mathbb{R}$  such that  $D_\alpha(\mu, \nu) < \infty$ .

The consistency of  $D_\alpha(\hat{P}_n, U)$  introduced in Definition 1 means that  $D_\alpha(\hat{P}_n, U)$ -based test of the hypothesis  $\mathcal{H} : U$  against the alternative  $\mathcal{A}_n : P_n$  of any fixed size has power tending to 1. Indeed, under  $\mathcal{H}$  we have  $D_\alpha(\hat{P}_n, U) \xrightarrow{p} 0$  so that the rejection level  $r_n(s)$  of the  $D_\alpha(\hat{P}_n, U)$ -based test of size  $s \in ]0; 1[$  tends to 0 for  $n \rightarrow \infty$  while under  $\mathcal{A}_n$  we have  $D_\alpha(\hat{P}_n, U) \xrightarrow{p} \Delta_\alpha > 0$ .

*Definition 2:* For  $\alpha \in \mathbb{R}$  we say that  $g_\alpha$  is the Bahadur function for the statistic  $T_\alpha = 2nD_\alpha(\hat{P}_n, U)$  if  $g_\alpha : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  is continuous and there exists a sequence  $c_{\alpha, n} > 0$  such that under  $\mathcal{H}$

$$\lim_{n \rightarrow \infty} -\frac{c_{\alpha, n}}{n} \ln \mathbb{P}(D_\alpha(\hat{P}_n, U) \geq \Delta) = g_\alpha(\Delta), \quad \Delta > 0. \quad (32)$$

*Remark 1:* One should note that the Bahadur function depends on the sequence  $c_{\alpha, n}$ . For the kind of results that we are interested in, the crucial thing is to determine the asymptotic behavior of sequences  $c_{\alpha, n}$  admitting the Bahadur function rather than the exact value of this. Nevertheless we shall calculate the exact value of the Bahadur function for certain sequences because this will allow us to use standard terminology and because the determination of the Bahadur function may be of independent interest.

Next follows the basic definition of the present paper where  $\Delta_{\alpha_1}, \Delta_{\alpha_2}$  are the limits from the Bahadur condition and  $g_{\alpha_1}, g_{\alpha_2}$  and  $c_{\alpha_1, n}, c_{\alpha_2, n}$  are the functions and sequences from the definition of Bahadur function. In this definition we apply to the power divergence statistics  $T_{\alpha_1}$  and  $T_{\alpha_2}$  the concept of the Bahadur efficiency  $BE(T_1 | T_2)$  introduced for more general statistics  $T_1$  and  $T_2$  in [6, p. 732].

*Definition 3:* Let the statistic  $D_{\alpha_1}(\hat{P}_n, U)$  and  $D_{\alpha_2}(\hat{P}_n, U)$  be consistent and let the Bahadur functions  $g_{\alpha_1}$  and  $g_{\alpha_2}$  of the power divergence statistics  $T_{\alpha_1}$  and  $T_{\alpha_2}$  exist. Then the Bahadur efficiency  $BE(T_{\alpha_1} | T_{\alpha_2})$  of  $T_{\alpha_1}$  with respect to  $T_{\alpha_2}$  is defined by

$$BE(T_{\alpha_1} | T_{\alpha_2}) = \frac{g_{\alpha_1}(\Delta_{\alpha_1})}{g_{\alpha_2}(\Delta_{\alpha_2})} \lim_{n \rightarrow \infty} \frac{c_{\alpha_1, n}}{c_{\alpha_2, n}} \quad (33)$$

provided the limit exists in  $[0; \infty]$ . Therefore this efficiency takes on values in the domain  $[0; \infty]$ .

Assume that the statistics  $D_{\alpha_i}(\hat{P}_n, U)$  are consistent for  $i \in \{1, 2\}$  and that there exist Bahadur functions  $g_{\alpha_i}$  satisfying (32) for some sequences  $c_{\alpha_i, n} > 0$ . Then the definition of consistency implies that both the  $T_{\alpha_i}$ -tests of the uniformity hypothesis  $\mathcal{H} : U$  will achieve identical powers

$$\pi = \mathbb{P}(D_{\alpha_i}(\hat{P}_n, U) \geq r_{n,i})$$

for  $\pi \in ]0, 1[$  and  $i \in \{1, 2\}$  under  $\mathcal{A}_n$  if and only if  $r_{n,i} \downarrow \Delta_{\alpha_i}$  for  $i \in \{1, 2\}$  as  $n \rightarrow \infty$ . The convergence  $r_{n,i} \downarrow \Delta_{\alpha_i}$  leads to the approximate  $T_{\alpha_i}$ -test sizes

$$\begin{aligned} s_{n,i} &\triangleq \mathbb{P}(D_{\alpha_i}(\hat{P}_n, U) \geq \Delta_{\alpha_i}) \\ &\approx \mathbb{P}(D_{\alpha_i}(\hat{P}_n, U) \geq r_{n,i}) \end{aligned}$$

for  $i \in \{1, 2\}$  under  $\mathcal{H}$  where  $s_{n,i} \rightarrow 0$  as  $n \rightarrow \infty$  for  $i \in \{1, 2\}$  under  $\mathcal{H}$ . By (32), the  $T_{\alpha_i}$ -tests need different sample sizes

$$n_i = \frac{c_{\alpha_i, n}}{g_{\alpha_i}(\Delta_{\alpha_i})} \ln \frac{1}{s_n}, \quad i \in \{1, 2\} \quad (34)$$

to achieve the same approximate test sizes  $s_n = s_{n,1} = s_{n,2}$  when  $n$  is here playing the role of a formal parameter that




Fig. 2. The function  $y \rightarrow y^\alpha$  illustrated for  $\alpha = 3/2$  by the full line. The lower bound given in (35) is indicated by the dashed straight line for  $x = 0.5$ . The upper bound in (36) is indicated by the dotted line.

increases to  $\infty$ . Thus Definition 3 formalizes the concept of asymptotic relative efficiency announced at the beginning of this section.

Before presenting the main results based on Definition 3 in Section V, we investigate sufficient conditions for consistency of the statistic  $D_\alpha(\hat{P}_n, U)$  in the domain  $\alpha \geq 1$  in Section III. Section IV presents conditions for the existence of the corresponding Bahadur functions  $g_\alpha$ ,  $\alpha \geq 1$  and explicitly evaluates these functions.

### III. CONSISTENCY

When a statistician uses  $D_\alpha(\hat{P}_n, U)$  as a statistic to distinguish between  $P_n$  and  $U$  then he does so because he considers the plug-in estimator  $D_\alpha(\hat{P}_n, U)$  as a good estimate of  $D_\alpha(P_n, U)$ . This idea was made precise in Definition 1 dealing with the important concept of consistency of  $D_\alpha(\hat{P}_n, U)$ . Our next theorem presents consistency conditions for all statistic  $D_\alpha(\hat{P}_n, U)$ ,  $\alpha \geq 1$ . It is based on the following auxiliary result.

*Lemma 1:* For  $x \in [0; 1[$  and  $y \in [0; 1]$  and  $\alpha \in ]1; 2[$  we have

$$|y^\alpha - x^\alpha| \leq \alpha x^{\alpha-1} |y - x| + (\alpha - 1) x^{\alpha-2} (y - x)^2.$$

*Proof:* First we observe that

$$y^\alpha \geq x^\alpha + \alpha x^{\alpha-1} (y - x) \quad (35)$$

because the function  $y \rightarrow y^\alpha$  is convex. Next we prove the inequality

$$y^\alpha \leq x^\alpha + \alpha x^{\alpha-1} (y - x) + (\alpha - 1) x^{\alpha-2} (y - x)^2. \quad (36)$$

The upper and lower bounds in (35) and (36) are illustrated in Figure 2.

We have to prove that

$$y^\alpha - \left( x^\alpha + \alpha x^{\alpha-1} (y - x) + (\alpha - 1) x^{\alpha-2} (y - x)^2 \right)$$

is negative. This is obvious for  $y = x$  and for  $y = 0$ . The derivative is

$$\begin{aligned} \alpha y^{\alpha-1} - (\alpha x^{\alpha-1} + (\alpha - 1) x^{\alpha-2} 2(y - x)) \\ = \alpha y^{\alpha-1} + (\alpha - 2) x^{\alpha-1} + (2 - 2\alpha) x^{\alpha-2} y. \end{aligned}$$

The derivative is 0 for  $y = x$ . Differentiate once more and get

$$\begin{aligned} \alpha(\alpha - 1) y^{\alpha-2} + (2 - 2\alpha) x^{\alpha-2} \\ = (\alpha - 1) (\alpha y^{\alpha-2} - 2x^{\alpha-2}), \end{aligned}$$

which is positive for  $y \leq \left(\frac{\alpha}{2}\right)^{\frac{1}{2-\alpha}} x \leq x$ . Combining (36) and (35) leads to

$$0 \leq y^\alpha - x^\alpha - \alpha x^{\alpha-1} (y - x) \leq (\alpha - 1) x^{\alpha-2} (y - x)^2. \quad (37)$$

Now  $1 - x^\alpha - \alpha x^{\alpha-1} (1 - x)$  is increasing in  $x$  and equals 0 for  $x = 1$  so the lower bound in (37) side is negative and we have

$$|y^\alpha - x^\alpha - \alpha x^{\alpha-1} (y - x)| \leq (\alpha - 1) x^{\alpha-2} (y - x)^2.$$

The inequality follows because

$$\begin{aligned} |y^\alpha - x^\alpha| &\leq |\alpha x^{\alpha-1} (y - x)| \\ &\quad + |y^\alpha - x^\alpha - \alpha x^{\alpha-1} (y - x)|. \end{aligned}$$

We shall also use the following upper bound a number of times

$$\begin{aligned} E(\hat{p}_j - p_j)^2 &= \frac{p_j(1 - p_j)}{n} \\ &\leq \frac{p_j}{n} \end{aligned}$$

For divergence of order 2 it gives

$$\begin{aligned} ED_2(\hat{P}_n \| P) &= \sum_{j=1}^k \frac{E(\hat{p}_j - p_j)^2}{p_j} \\ &\leq \sum_{j=1}^k \frac{1}{n} \\ &= \frac{k}{n}. \end{aligned} \quad (38)$$

*Theorem 1:* For all  $\alpha \geq 1$  let the Bahadur condition (29) hold. Then  $D_\alpha(\hat{P}_n, U)$  is consistent if

$$\alpha \in [1; 2] \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{k}{n} = 0, \quad (39)$$

or

$$\alpha > 2 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{k^{\alpha-1}}{n} = 0. \quad (40)$$

*Proof:* Under  $\mathcal{H}$  we have  $D_\alpha(P_n, U) = D_\alpha(U, U) = 0$ . Hence it suffices to prove that under both  $\mathcal{H}$  and  $\mathcal{A}_n$

$$\lim_{n \rightarrow \infty} E \left| D_\alpha(\hat{P}_n, U) - D_\alpha(P_n, U) \right| = 0 \quad \text{for all } \alpha \geq 1.$$

Put for brevity  $\Lambda_{\alpha,n} = D_{\alpha}(\hat{P}_n, U) - D_{\alpha}(P_n, U)$  and denote the variance by  $\text{Var}$  and the covariance by  $\text{Cov}$ . The cases **i)**  $\alpha = 1$ , **ii)**  $\alpha \in ]1; 2]$  and **iii)**  $\alpha > 2$  have to be treated separately.

**(i)** For  $\alpha = 1$  we have

$$\Lambda_{1,n} = \sum_{j=1}^k (\hat{p}_j \ln \hat{p}_j - p_j \ln p_j), \quad (41)$$

where we dropped the subscript  $n$  everywhere in the sum. From (41) we obtain

$$\Lambda_{1,n} = \sum_{j=1}^k \hat{p}_j \ln \frac{\hat{p}_j}{p_j} - \sum_{j=1}^k (\hat{p}_j - p_j) \ln \frac{1}{p_j}$$

and hence

$$|\Lambda_{1,n}| \leq D_1(\hat{P}_n, P_n) + \left| \sum_{j=1}^k (\hat{p}_j - p_j) \ln \frac{1}{p_j} \right|.$$

If  $\hat{n}_i = \hat{p}_i n$  is the number of observations of type  $i$  then

$$D_1(\hat{P}_n, P_n) \leq 2D_2(\hat{P}_n, P_n).$$

Therefore

$$\mathbb{E} |\Lambda_{1,n}| \leq 2\mathbb{E} D_2(\hat{P}_n, P_n) + \mathbb{E} \left| \sum_{j=1}^k (\hat{p}_j - p_j) \ln \frac{1}{p_j} \right|. \quad (42)$$

The last term on the right hand side can be bounded using Jensen's Inequality.

$$\begin{aligned} \mathbb{E} \left| \sum_{j=1}^k (\hat{p}_j - p_j) \ln \frac{1}{p_j} \right| &\leq \left( \mathbb{E} \left( \sum_{j=1}^k (\hat{p}_j - p_j) \ln \frac{1}{p_j} \right)^2 \right)^{1/2} \\ &= \left( \sum_{i,j=1}^k \ln p_j \ln p_i \text{COV}(\hat{p}_i, \hat{p}_j) \right)^{1/2} \\ &= \left( \sum_{i,j=1}^k \ln p_j \ln p_i \frac{\text{COV}(\hat{n}_i, \hat{n}_j)}{n^2} \right)^{1/2}, \quad (43) \end{aligned}$$

Further,

$$\begin{aligned} &\sum_{i,j=1}^k \ln p_j \ln p_i \frac{\text{Cov}(\hat{n}_i, \hat{n}_j)}{n^2} \\ &= \sum_{i=1}^k (\ln p_i)^2 \frac{\text{Var}(\hat{n}_i)}{n^2} + \sum_{i \neq j} \ln p_j \ln p_i \frac{\text{Cov}(\hat{n}_i, \hat{n}_j)}{n^2} \\ &\leq \sum_{i=1}^k (\ln p_i)^2 \frac{p_i}{n} + \sum_{i \neq j} \ln p_j \ln p_i \frac{np_i p_j}{n^2} \\ &= \frac{1}{n} \sum_{i=1}^k p_i (\ln p_i)^2 + \frac{1}{n} \left( \sum_{i=1}^k p_i \ln p_j \right)^2. \quad (44) \end{aligned}$$

The function  $x \rightarrow x \ln^2 x$  is concave in the interval  $[0; e^{-1}]$  and convex in the interval  $[e^{-1}; 1]$ . Therefore we are able to

use [3, Theorem 3.1] to see that  $\sum_{i=1}^k p_i (\ln p_i)^2$  attains its maximum for a mixture of uniform distributions on  $k$  and  $k-1$  points. Thus

$$\begin{aligned} \sum_{i=1}^k p_i (\ln p_i)^2 &\leq \sum_{i=1}^k \frac{1}{k-1} \left( \ln \frac{1}{k} \right)^2 \\ &= \frac{k (\ln k)^2}{k-1} \\ &\leq 2 (\ln k)^2. \end{aligned} \quad (45)$$

The sum  $\sum_{i=1}^k p_i \ln p_j$  equals minus the entropy, which has maximum  $\ln k$ . By combining (38), (42), (43), (44), and (45) we get

$$\mathbb{E} |\Lambda_{1,n}| \leq \frac{2k}{n} + \left( \frac{3 (\ln k)^2}{n} \right)^{1/2}$$

and the right hand side tends to zero under the condition (39) for  $n$  tending to infinity. This proves (39) for  $\alpha = 1$ .

**(ii)** For every  $\alpha \in ]1; 2[$  we have

$$\Lambda_{\alpha,n} = \frac{k^{\alpha-1}}{\alpha(\alpha-1)} \sum_{j=1}^k (\hat{p}_j^{\alpha} - p_j^{\alpha}).$$

Using the abbreviation

$$D_{\alpha,n} = D_{\alpha}(P_n, U), \quad (46)$$

we obtain

$$\begin{aligned} |\Lambda_{\alpha,n}| &\leq \frac{k^{\alpha-1}}{\alpha(\alpha-1)} \sum_{j=1}^k \left( \alpha p_j^{\alpha-1} |\hat{p}_j - p_j| + (\alpha-1) p_j^{\alpha-2} (\hat{p}_j - p_j)^2 \right) \leq \\ &\frac{k^{\alpha-1}}{\alpha-1} \left( \sum_{j=1}^k (p_j^{\alpha/2})^2 \right)^{1/2} \left( \sum_{j=1}^k p_j^{\alpha-2} (\hat{p}_j - p_j)^2 \right)^{1/2} \\ &\quad + \frac{k^{\alpha-1}}{\alpha} \sum_{j=1}^k p_j^{\alpha-2} (\hat{p}_j - p_j)^2 = \\ &\frac{k^{\frac{\alpha-1}{2}} \left( \alpha(\alpha-1) D_{\alpha,n} + 1 \right)^{1/2}}{\alpha-1} \left( \sum_{j=1}^k p_j^{\alpha-2} (\hat{p}_j - p_j)^2 \right)^{1/2} \\ &\quad + \frac{k^{\alpha-1}}{\alpha} \sum_{j=1}^k p_j^{\alpha-2} (\hat{p}_j - p_j)^2 \end{aligned}$$

because  $\sum_{j=1}^k p_j^\alpha = [\alpha(\alpha-1)D_{\alpha,n} + 1]/k^{\alpha-1}$ . Thus

$$\begin{aligned} \mathbb{E}|\Lambda_{\alpha,n}| &\leq \frac{k^{\frac{\alpha-1}{2}}}{\alpha-1} (\alpha(\alpha-1)D_{\alpha,n} + 1)^{1/2} \left( \sum_{j=1}^k p_j^{\alpha-2} \frac{p_j}{n} \right)^{1/2} \\ &\quad + \frac{k^{\alpha-1}}{\alpha} \sum_{j=1}^k p_j^{\alpha-2} \frac{p_j}{n} \\ &= \frac{(\alpha(\alpha-1)D_{\alpha,n} + 1)^{1/2}}{\alpha-1} \left( \frac{k^{\alpha-1} \sum_{j=1}^k p_j^{\alpha-1}}{n} \right)^{1/2} \\ &\quad + \frac{k^{\alpha-1}}{\alpha n} \sum_{j=1}^k p_j^{\alpha-1}. \end{aligned}$$

The sequence  $D_{\alpha,n}$  is upper bounded and that implies that there exists a constant  $c < 1$  such that  $p_j \leq c$  for all  $n, j$ . Thus

$$\begin{aligned} \mathbb{E}|\Lambda_{\alpha,n}| &\leq \frac{\left( \frac{\alpha(\alpha-1)D_{\alpha,n} + 1}{\alpha-1} \right)^{1/2} \left( \frac{k^{\alpha-1} k \left(\frac{1}{k}\right)^{\alpha-1}}{n} \right)^{1/2}}{\alpha-1} \\ &\quad + \frac{2 k^{\alpha-1} k \left(\frac{1}{k}\right)^{\alpha-1}}{\alpha n} \\ &= \frac{(\alpha(\alpha-1)D_{\alpha,n} + 1)^{1/2} \left(\frac{k}{n}\right)^{1/2}}{\alpha-1} + \frac{k}{\alpha n}. \end{aligned}$$

This proves (39) for  $\alpha \in [1; 2]$ .

(iii) For  $\alpha \geq 2$  we shall use the second order Taylor expansion

$$\hat{p}_j^\alpha = p_j^\alpha + \alpha p_j^{\alpha-1} (\hat{p}_j - p_j) + \frac{\alpha(\alpha-1)}{2} \xi_j^{\alpha-2} (\hat{p}_j - p_j)^2,$$

leading to

$$|\Lambda_{\alpha,n}| \leq \frac{k^{\alpha-1}}{\alpha-1} \sum_{j=1}^k p_j^{\alpha-1} |\hat{p}_j - p_j| + \frac{k^{\alpha-1}}{2} \sum_{j=1}^k (\hat{p}_j - p_j)^2. \quad (47)$$

We use Cauchy-Schwarz' inequality on the first term on the right hand side of (47) to get

$$\begin{aligned} \frac{k^{\alpha-1}}{\alpha-1} \sum_{j=1}^k p_j^{\alpha-1} |\hat{p}_j - p_j| &\leq \frac{k^{\alpha-1}}{\alpha-1} \left( \sum_{j=1}^k p_j^{2(\alpha-1)} \right)^{1/2} \left( \sum_{j=1}^k (\hat{p}_j - p_j)^2 \right)^{1/2}. \quad (48) \end{aligned}$$

Using the sequence  $D_{\alpha,n}$  introduced in (46) we get

$$\begin{aligned} \sum_{j=1}^k p_j^{2(\alpha-1)} &\leq \sum_{j=1}^k p_j^\alpha \\ &= \frac{\alpha(\alpha-1)D_{\alpha,n} + 1}{k^{\alpha-1}}. \end{aligned} \quad (49)$$

By combining (47), (48) and (49) we get

$$\begin{aligned} \mathbb{E}|\Lambda_{\alpha,n}| &\leq \frac{k^{\alpha-1}}{\alpha-1} \left( \frac{\alpha(\alpha-1)D_{\alpha,n} + 1}{k^{\alpha-1}} \right)^{1/2} \left( \sum_{j=1}^k \mathbb{E}(\hat{p}_j - p_j)^2 \right)^{1/2} \\ &\quad + \frac{k^{\alpha-1}}{2} \sum_{j=1}^k \mathbb{E}(\hat{p}_j - p_j)^2 \\ &\leq \frac{1}{\alpha-1} \left( \frac{k^{\alpha-1}}{n} (\alpha(\alpha-1)D_{\alpha,n} + 1) \right)^{1/2} + \frac{k^{\alpha-1}}{2n}. \end{aligned}$$

But  $D_{\alpha,n}$  is zero under the hypothesis of a uniform distribution and, by (29), has a finite limit under the alternative. This completes the proof of (40). ■

*Example 4:* Assume that for  $\alpha = 3$  the model satisfies the Bahadur condition, in particular that (29) holds with  $\alpha = 3$ . Then

$$\mathbb{E}D_3(\hat{P}_n, U) = \frac{k^2 \mathbb{E} \left( \sum_{j=1}^k \hat{p}_j^3 \right) - 1}{6}$$

where

$$\hat{p}_j^3 = p_j^3 + 3p_j^2(\hat{p}_j - p_j) + 3p_j(\hat{p}_j - p_j)^2 + (\hat{p}_j - p_j)^3.$$

Therefore

$$\begin{aligned} \mathbb{E}D_3(\hat{P}_n, U) &= \frac{k^2 \sum_{j=1}^k \mathbb{E} \left( \frac{p_j^3 + 3p_j^2(\hat{p}_j - p_j)}{+3p_j(\hat{p}_j - p_j)^2 + (\hat{p}_j - p_j)^3} \right) - 1}{6} \\ &= \frac{k^2 p_j^3 - 1}{6} \\ &\quad + \frac{k^2}{6} \sum_{j=1}^k \left( 3p_j \mathbb{E}(\hat{p}_j - p_j)^2 + \mathbb{E}(\hat{p}_j - p_j)^3 \right). \end{aligned}$$

By taking mean values we get

$$\begin{aligned} \mathbb{E}D_3(\hat{P}_n, U) &= D_3(P_n, U) \\ &\quad + \frac{k^2}{6} \sum_{j=1}^k \left( 3p_j \frac{p_j(1-p_j)}{n} + \frac{p_j(1-p_j)(1-2p_j)}{n} \right) \\ &= D_3(P_n, U) + \frac{k^2}{6n} \sum_{j=1}^k (p_j - p_j^3) \\ &= D_3(P_n, U) + \frac{k^2 \left( 1 - \sum_{j=1}^k p_j^3 \right)}{6n} \\ &= D_3(P_n, U) + \frac{k^2 - 6D_3(P_n, U) - 1}{6n}. \end{aligned}$$

By (29),  $D_3(P_n, U)$  is bounded away from 0 under  $\mathcal{A}_n$  uniformly for all sufficiently large  $n$ . Therefore (40) is not only sufficient but also necessary for the consistency of statistic  $D_\alpha(\hat{P}_n, U)$  in the spacial case  $\alpha = 3$ .

## IV. BAHADUR FUNCTIONS

Throughout this section we consider the statistical testing model (14) - (15) under the hypothesis  $\mathcal{H}$ . This means that  $P(B_n)$  denotes the probability of the random events  $B_n$  depending on  $\mathbf{X}_n$  with a multinomial distribution in the sense of (14). As before, we consider  $k = k_n$  depending on the sample size  $n$  and we study the Bahadur functions (32) corresponding to the statistic  $D_\alpha(\hat{P}_n, U)$  for  $\alpha \geq 1$ .

*Example 5:* Let  $k = k_n$  increase so slowly that

$$\lim_{n \rightarrow \infty} \frac{k \ln n}{n} = 0. \quad (50)$$

Then (32) holds for the sequence  $c_{1,n} \equiv 1$  and function

$$g_1(\Delta) = \Delta \quad \text{for all } \Delta > 0, \quad (51)$$

i.e., (51) is the Bahadur function for the log-likelihood ratio statistic

$$T_1 = 2nD_1(\hat{P}_n, U).$$

This result was first obtained independently in [20, Corollary 2.4] and [6, Theorem 2]. Using the simple method based on the inequality (52) below, it was obtained in [17, Theorem 2].

According to the result of [21] made precise in [22, Problem 1.2.11] and in [23, p. 16], for every subset  $A \subset M(k)$  the divergence  $D_1(P, U)$  defined by (11) satisfies the inequality

$$\left| \inf_{P \in A \cap M(k|n)} D_1(P, U) + \frac{1}{n} \ln P(\hat{P}_n \in A) \right| \leq \frac{k \ln(n+1)}{n}. \quad (52)$$

Hence the approximation of  $-\frac{1}{n} \ln P(\hat{P}_n \in A)$  in (32) by means of the infimum appearing in (52) is possible under the restriction

$$\lim_{n \rightarrow \infty} c_{\alpha,n} \frac{k \ln n}{n} = 0 \quad (54)$$

on the sequence  $k = k_n$ , in addition to (24).

In the rest of this section we present an alternative to the formula (32) for the Bahadur functions  $g_\alpha$ ,  $\alpha \in \mathbb{R}$  which is based on the inequality (52). These formulas are given in terms of the Shannon entropy  $H(P)$  maximized on the sets

$$A_{\alpha,\Delta}(k) = \{P \in M(k) : D_\alpha(P, U) \geq \Delta\} \quad (55)$$

and

$$A_{\alpha,\Delta}(k|n) = A_{\alpha,\Delta}(k) \cap M(k|n) \quad (56)$$

or, equivalently, in terms of the information divergence  $D_1(\hat{P}_n, U)$  minimized on these sets.

*Lemma 2:* Assume that for some  $\alpha \in \mathbb{R}$  and  $k = k_n$  there exist  $c_{\alpha,n} > 0$  satisfying (54) such that the sequence of functions

$$G_{\alpha,\Delta}(k|n) = c_{\alpha,n} \left( \inf_{P \in A_{\alpha,\Delta}(k|n)} D_\alpha(P, U) \right), \quad \Delta > 0 \quad (57)$$

converges to a positive limit limit

$$g_\alpha(\Delta) = \lim_{n \rightarrow \infty} G_{\alpha,\Delta}(k|n), \quad \Delta > 0. \quad (58)$$




Fig. 3. In the simplex of distributions on a 3-element set, the distributions with index of coincidence less a certain value are indicated by the shaded area. A level curve of the entropy function is indicated by a full curve. Mixtures of the uniform distribution  $U$  and a Dirac distribution at one of the extreme points is indicated by a dashed line. Maximal entropy over the shaded area is obtained at the point where the dashed line leaves the shaded area. The two other points with maximal entropy are indicated as well.

Then the Bahadur function for the power divergence statistic  $T_\alpha$  is equal to  $g_\alpha$ .

*Proof:* Using (19) and (52) we get that the functions (57) satisfy the inequality

$$\left| \frac{c_{\alpha,n}}{n} \ln P(D_\alpha(\hat{P}_n, U) \geq \Delta) + G_{\alpha,\Delta}(k|n) \right| \leq \frac{k c_{\alpha,n} \ln(n+1)}{n}.$$

Since (54) holds, (58) follows from here and from (33). ■

In the following assertion we consider for arbitrary  $\alpha \in \mathbb{R}$ ,  $k = k_n$  and  $c_{\alpha,n} > 0$  the sequence of functions

$$G_{\alpha,\Delta}(k) = c_{\alpha,n} \left( \inf_{P \in A_{\alpha,\Delta}(k)} D_\alpha(P, U) \right), \quad \Delta > 0. \quad (59)$$

Obviously,  $G_{\alpha,\Delta}(k) \leq G_{\alpha,\Delta}(k|n)$ .

*Lemma 3:* Let for some  $\alpha \in \mathbb{R}$  the Bahadur condition hold and  $c_{\alpha,n} > 0$  satisfy (54). If the corresponding sequences of functions (57) and (59) asymptotically coincide in the sense

$$\lim_{n \rightarrow \infty} [G_{\alpha,\Delta}(k|n) - G_{\alpha,\Delta}(k)] = 0 \quad (60)$$

and at the same time  $G_{\alpha,\Delta}(k)$  converges to a positive limit

$$g_\alpha(\Delta) = \lim_{n \rightarrow \infty} G_{\alpha,\Delta}(k), \quad \Delta > 0 \quad (61)$$

then  $g_\alpha$  is the Bahadur function for the power divergence statistic  $T_\alpha$ .

*Proof:* Clear from the assumption (60) and Lemma 2. ■

In [3] it was proved that for every  $x \in [k^{1-\alpha}; 1]$  and for the Dirac distribution  $\mathbf{1} = (1, 0, \dots, 0) \in M(k)$ , the equation

$$IC_\alpha(s\mathbf{1} + (1-s)U) = x \quad (62)$$



has a unique solution  $s \in [0; 1]$  and that this solution satisfies the relation

$$\sup_{IC_\alpha(P) \geq x} H(P) = H(s\mathbf{1} + (1-s)U). \quad (63)$$

This result is illustrated in Figure 3. It leads to the following lemma using the constants

$$s_{\alpha,k} = \frac{1 - (1 - 1/k)^{1/(\alpha-1)}}{1 + (1 - 1/k)^{\alpha/(\alpha-1)}} \in ]0; 1[. \quad (64)$$

*Lemma 4:* For every  $\alpha > 1$  and

$$\frac{1}{\alpha(\alpha-1)k} < \Delta \leq \frac{k^{\alpha-1} - 1}{\alpha(\alpha-1)}, \quad (65)$$

the equation

$$\frac{1}{k}(ks + 1 - s)^\alpha + (1 - s)^\alpha = 1 + \alpha(\alpha-1)\Delta \quad (66)$$

has a unique solution  $s \in [0; 1]$  and this solution satisfies the inequality

$$s_{\alpha,k} < s \leq 1 \quad (67)$$

and the equality

$$\begin{aligned} \inf_{P \in A_{\alpha,\Delta}(k)} D_1(P, U) = \\ \frac{1}{k}(ks + 1 - s) \ln(ks + 1 - s) + (1 - s) \ln(1 - s). \end{aligned} \quad (68)$$

*Proof:* By definition of  $A_{\alpha,\Delta}(k)$  in (55) and (20),  $P \in A_{\alpha,\Delta}(k)$  if and only if  $IC_\alpha(P) \geq x$  for

$$x = k^{1-\alpha}[1 + \alpha(\alpha-1)\Delta].$$

By the definition of  $IC_\alpha(P)$  in (21),

$$\begin{aligned} IC_\alpha(s\mathbf{1} + (1-s)U) \\ = k^{1-\alpha} \left( \frac{1}{k}(ks + 1 - s)^\alpha + (1 - s)^\alpha \right) \end{aligned} \quad (69)$$

so that the equation (62) is equivalent to (66). Further,

$$\begin{aligned} D_1(s\mathbf{1} - (1-s)U, U) = \\ \frac{1}{k}(ks + 1 - s) \ln(ks + 1 - s) + (1 - s) \ln(1 - s). \end{aligned} \quad (70)$$

Therefore, by (63) and (62), the relation (68) will be proved if we prove that the equation (66) has a unique solution  $s \in [0; 1]$ . One can verify by differentiation that the continuous function

$$\psi(s) = \frac{1}{k}(ks + 1 - s)^\alpha + (1 - s)^\alpha, \quad s \in [0, 1] \quad (71)$$

appearing on the left of (66) is decreasing on the interval  $[0; s_{\alpha,k}]$  and increasing on the complement  $]s_{\alpha,k}; 1]$ . Since

$$\psi(0) = \frac{1}{k} + 1 \quad \text{and} \quad \psi(1) = k^{\alpha-1},$$

for each  $\Delta$  satisfying (65) the solution  $s \in [0; 1]$  is unique and strictly greater than  $s_{\alpha,k}$ . Thus not only (68) but also (67) is valid. ■

In the following lemma and everywhere in the sequel, convergence as well as the symbols  $o(\cdot)$  and  $O(\cdot)$ , are considered

for  $n \rightarrow \infty$ . We remind that  $k = k_n$  is assumed to satisfy (24).

*Lemma 5:* For every  $\alpha > 1$  and  $\Delta > 0$ ,

$$\begin{aligned} \inf_{P \in A_{\alpha,\Delta}(k)} D_1(P, U) = \\ \left( [\alpha(\alpha-1)\Delta]^{1/\alpha} + o(1) \right) \frac{k^{1/\alpha} \ln k^{1/\alpha}}{k}. \end{aligned}$$

*Proof:* Consider arbitrary  $\alpha > 1$  and  $\varepsilon > 0$ . Since  $k = k_n$  satisfies (24), Lemma 4 implies for all sufficiently large  $n$  that the equation (66) has in the interval  $]s_{\alpha,k}; 1]$  a unique solution  $s = s_k$  satisfying (68). Therefore it suffices to prove that the sequence

$$x_k = \frac{1}{k}(ks_k + 1 - s_k) \ln(ks_k + 1 - s_k) \quad (72)$$

$$+ (1 - s_k) \ln(1 - s_k) \quad (73)$$

and the positive constant

$$\delta = [\alpha(\alpha-1)\Delta]^{1/\alpha}$$

satisfy the asymptotic relation

$$x_k = (\delta + o(1)) \frac{k^{1/\alpha} \ln k^{1/\alpha}}{k}. \quad (74)$$

By (64) and (67),  $s_k$  is a positive sequence and (66) with  $s$  replaced by  $s_k$  obviously contradicts the assumption

$$\limsup_{k \rightarrow \infty} s_k > 0.$$

Therefore, under (24),  $s_k = o(1)$  and, consequently, (66) with  $s$  replaced by  $s_k$  leads to the asymptotic relation

$$\frac{1}{k}[ks_k + O(1)]^\alpha + 1 + o(1) = 1 + \delta^\alpha.$$

This relation implies that

$$s_k = \frac{\delta k^{1/\alpha}}{k} + o\left(\frac{k^{1/\alpha}}{k}\right) \quad (75)$$

and the desired relation (74) follows from here and from the definition of  $x_k$  in (72). ■

In the rest of the paper we are interested in the sequences

$$c_{\alpha,n} = \frac{k}{k^{1/\alpha} \ln k^{1/\alpha}} \quad (76)$$

for  $\alpha > 1$  and  $k = k_n$  satisfying (24).

*Lemma 6:* If  $c_{\alpha,n}$  is given by (76) and (24) is satisfied then (60) holds for every  $\alpha > 1$  and  $\Delta > 0$ .

*Proof:* Let  $\alpha$ ,  $\Delta$  and  $s_k$  be the same as in the previous proof. Further, denote by  $\ell_k$  the integer part of  $n(1 - s_k)/k$ ,

$$\ell_k = \left\lfloor \frac{n(1 - s_k)}{k} \right\rfloor,$$

and define

$$\tilde{s}_k = \frac{n - k\ell_k}{n},$$

$$\tilde{P}_k = \tilde{s}_k \mathbf{1} + (1 - \tilde{s}_k) U,$$

$$P_k = s_k \mathbf{1} + (1 - s_k) U,$$

where  $\mathbf{1}$  and  $U$  are the same elements of  $M(k)$  as in (69) and (70). Then

$$s_k \leq \tilde{s}_k \leq s_k + \frac{k}{n}, \quad (77)$$

and one obtains from (20), (69) and (71)

$$D_\alpha(\tilde{P}_k, U) = \frac{1}{\alpha(\alpha-1)} (k^{\alpha-1}\psi(\tilde{s}_k) - 1)$$

and

$$D_\alpha(P_k, U) = \frac{1}{\alpha(\alpha-1)} (k^{\alpha-1}\psi(s_k) - 1).$$

The distribution  $P_k$  belongs to  $A_{\alpha,\Delta}(k)$  of (55). Indeed,  $s_k$  satisfies (66) and, consequently,  $D_\alpha(P_k, U) = \Delta$ . The distribution  $\tilde{P}_k$  belongs to  $M(k|n) \subset M(k)$  defined by (1). Further, in the proof of Lemma 4 we argued that the function  $\psi(s)$  of (71) is increasing in the domain  $]s_k; 1] \subset ]s_{k,\alpha}; 1]$ . Therefore the left-hand side of (77) implies  $D_\alpha(\tilde{P}_k, U) \geq \Delta$ , which means that  $\tilde{P}_k$  belongs to  $A_{\alpha,\Delta}(k|n)$  of (56). Consequently,

$$\inf_{P \in A_{\alpha,\Delta}(k|n)} D_1(P, U) \leq D_1(\tilde{P}_k, U)$$

where

$$\begin{aligned} D_1(\tilde{P}_k, U) &= \ln k - H(\tilde{P}_k) \\ &= \ln k - H(\tilde{s}_k \mathbf{1} + (1 - \tilde{s}_k) U) \\ &= \tilde{x}_k \quad \text{cf. (70)} \end{aligned}$$

for  $\tilde{x}_k$  defined by (72) with  $s_k$  replaced by  $\tilde{s}_k$ . Further, in the previous proof we deduced for  $x_k$  of (72) the formula (74) from the asymptotic property (75) of  $s_k$ . However, under (24) the sequence  $\tilde{s}_k$  satisfies this asymptotic property too. Therefore (74) remains to be valid with  $x_k$  replaced by  $\tilde{x}_k$ . This means that under (24) takes place the asymptotic relation

$$\begin{aligned} \inf_{P \in A_{\alpha,\Delta}(k|n)} D_1(P, U) & \\ & \leq \left( [\alpha(\alpha-1)\Delta]^{1/\alpha} + o(1) \right) \frac{k^{1/\alpha} \ln k^{1/\alpha}}{k}. \end{aligned}$$

Combining this with the result of Lemma 5, we obtain the desired relation (60).  $\blacksquare$

In (51) we presented a simple explicit formula for the Bahadur function  $g_1$  of the log-likelihood ratio statistic  $T_1$ . Now we can give explicit formulas for the Bahadur functions of the remaining statistic  $T_\alpha$ ,  $\alpha > 1$ .

*Theorem 2:* Let  $k = k_n$  increase to infinity slowly in the sense that for some  $\alpha > 1$

$$\lim_{n \rightarrow \infty} \frac{k_n^{2-(1/\alpha)} \ln n}{n \ln k_n} = 0. \quad (78)$$

Then (33) holds for the sequence  $c_{\alpha,n}$  given by (76) and for the function

$$g_\alpha(\Delta) = [\alpha(\alpha-1)\Delta]^{1/\alpha}, \quad \Delta > 0 \quad (79)$$

i.e., (79) is the Bahadur function of the statistic  $T_\alpha$ .

*Proof:* Let  $\alpha > 1$  be arbitrary fixed. If  $c_{\alpha,n}$  is given by (76) then (78) implies (24) as well as (54). Hence it follows from Lemmas 3 and 6 that (32) holds for  $c_{\alpha,n}$  under consideration and for  $g_\alpha$  given by (79). Employing Lemma 4

we find that (61) reduces to (79) which completes the proof.  $\blacksquare$

The particular case of Theorem 2 for  $\alpha = 2$  was obtained in [6, Theorem 1] by using more complicated analytic methods involving limit theorems for multinomial and Poisson distributions. This particular case has been obtained also by [17] by using similar simple method as here, based on the inequality (52).

## V. MAIN RESULTS

The functions  $g_\alpha$  as well as the normalizing sequences  $c_{\alpha,n}$  have been explicitly evaluated in Theorem 2 and Example 5 for all  $\alpha \geq 1$ . Therefore (33) provides explicit Bahadur efficiencies  $BE(T_{\alpha_1} | T_{\alpha_2})$  on the whole domain  $\alpha_1, \alpha_2 \geq 1$ . These efficiencies are given in the following main result of this paper.

*Theorem 3:* Let  $1 \leq \alpha_1 < \alpha_2 < \infty$ .

(i) If the statistics  $D_{\alpha_1}(\hat{P}_n, U)$  and  $D_{\alpha_2}(\hat{P}_n, U)$  are consistent and  $k = k_n$  increases so slowly that

$$\lim_{n \rightarrow \infty} \frac{k^{2-1/\alpha_2} \ln n}{n} = 0 \quad (80)$$

then the Bahadur efficiency of the statistic  $T_{\alpha_1}$  with respect to  $T_{\alpha_2}$  satisfies the relation

$$BE(T_{\alpha_1} | T_{\alpha_2}) = \infty. \quad (81)$$

(ii) If  $k = k_n$  increases to infinity slowly in the sense that for some  $\beta \geq 3$

$$\lim_{n \rightarrow \infty} \frac{k^\beta}{n} = 0 \quad (82)$$

then (80) and the consistency required in (i) hold for all  $1 \leq \alpha_1 < \alpha_2 \leq \beta + 1$ . Hence in this case also the Bahadur efficiency relation (81) holds for all  $1 \leq \alpha_1 < \alpha_2 \leq \beta + 1$ .

*Proof:* (i) Let the assumptions of (i) hold for some  $1 < \alpha_1 < \alpha_2 < \infty$ . Then (80) implies (78) for  $\alpha = \alpha_1$  and  $\alpha = \alpha_2$ . By Theorem 2, the sequences  $c_{\alpha_1,n}$  and  $c_{\alpha_2,n}$  given by (76) for lead to the corresponding Bahadur functions  $g_{\alpha_1}$  and  $g_{\alpha_2}$  given by (32) and to the limit

$$\lim_{n \rightarrow \infty} \frac{c_{\alpha_2,n}}{c_{\alpha_1,n}} = \lim_{n \rightarrow \infty} \frac{k^{1/\alpha_1} \ln k^{1/\alpha_1}}{k^{1/\alpha_2} \ln k^{1/\alpha_2}} = \infty. \quad (83)$$

Relation (81) thus follows from (33) in Definition 3. If the assumptions of (i) hold for  $1 = \alpha_1 < \alpha_2 < \infty$  then instead of the above considered Bahadur function  $g_{\alpha_1}$  given by (32) we have  $g_{\alpha_1}(\Delta) = \Delta$  given by (51), and instead of  $c_{\alpha_1,n} = k_n/k_n^{1/\alpha_1} \ln k_n^{1/\alpha_1}$  given by (76) we have  $c_{\alpha_1,n} = 1$  given in Example 5. Therefore the limit

$$\lim_{n \rightarrow \infty} \frac{c_{\alpha_2,n}}{c_{\alpha_1,n}} = \lim_{n \rightarrow \infty} \frac{k_n}{k_n^{1/\alpha_2} \ln k_n^{1/\alpha_2}}$$

remains to be infinite as in (83).

(ii) If (82) holds for  $\beta \geq 3$  then  $(\beta - 2)/\beta > 0$  so that (82) implies

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \frac{k^2}{n^{2/\beta}} \\ &= \lim_{n \rightarrow \infty} \frac{k^2 \ln n}{n^{2/\beta} n^{(\beta-2)/\beta}} \\ &= \lim_{n \rightarrow \infty} \frac{k^2 \ln n}{n}. \end{aligned}$$

Therefore (82) holds for all  $\alpha_2 \geq 1$ . Further, if

$$\lim_{n \rightarrow \infty} \frac{k^3}{n} = 0$$

then it is easy to verify that the consistency conditions of Theorem 1 are satisfied for all  $1 \leq \alpha \leq 3$ , and if (82) holds then these conditions are satisfied for all  $1 \leq \alpha \leq \beta + 1$ . This completes the proof. ■

## VI. DISCUSSION

The special case of (81) with  $\alpha_1 = 1$  and  $\alpha_2 = 2$  with increasing  $k = k_n$  has been obtained in [6]. In the present paper we extended the fact that the log-likelihood ratio statistic  $T_1$  is more Bahadur efficient than the classical Pearson statistic  $T_2$  by proving that  $T_1$  is more Bahadur efficient than any statistics  $T_\alpha$  with  $\alpha > 1$ . Moreover, we found that the Bahadur efficiency of the power divergence statistic  $T_\alpha$  strictly decreases with  $\alpha$  increasing in the domain  $[1; \infty[$ . In particular any statistic  $T_\alpha$ ,  $1 \leq \alpha < 2$ , is more Bahadur efficient than the Pearson's  $T_2$ .

One of the aims of this paper was to verify whether there is a statistic  $T_\alpha$ ,  $\alpha \in \mathbb{R}$  more efficient in the Bahadur sense than  $T_1$ . In this respect, the result of Theorem 3 is negative. All we can say is that, if such a statistic exists, then it is most likely that it is of the form  $T_\alpha$  with  $\alpha \in ]0; 1[$ . Let us comment this conclusion in more detail.

In spite of that we do not have a systematic result for  $\alpha < 1$ , some fragments of such a result are available. Namely, [17] found the Bahadur functions  $g_0(\Delta) = g_{-1}(\Delta) = \Delta$  for all  $\Delta > 0$ , under the identical sequences  $c_{0,n} = c_{-1,n} = k_n$  figuring in (32). There is a small problem with the condition (29), because the event  $\min_j \hat{p}_{nj} = 0$  takes place with a positive probability and implies  $D_\alpha(P_n, U) = \infty$  for all  $\alpha \leq 0$ . Nevertheless the probability of this unpleasant event tends exponentially to zero, and one can modify the statistic  $T_{0,n}$  and  $T_{-1,n}$  in such a way that the above evaluated Bahadur functions and sequences remain unaltered and, at the same time, the consistency condition (30) hold, see [18] and [24]. Therefore in the light of present Theorem 2, the result of [17] means that the reversed log-likelihood ratio statistic  $T_0$ , and the Neyman statistic  $T_{-1}$ , are mutually equally Bahadur efficient, and each of them is less Bahadur efficient than any  $T_\alpha$ ,  $\alpha \geq 1$ . This extends the previous result of [18] who found  $T_0$  and  $T_{-1}$  to be less Bahadur efficient than  $T_2$ . If the low Bahadur efficiency of  $T_0$  and  $T_{-1}$  is shared by all statistics  $T_\alpha$  of the non-positive powers  $\alpha \leq 0$  then the possibility to find  $T_\alpha$  comparable with  $T_1$  or better is restricted to  $\alpha \in (0, 1)$ , as conjectured above.

**Acknowledgement:** We would like to thank Wouter Koolen-Wijkstra for comments that have improved the quality of this paper.

## REFERENCES

- [1] F. Liese and I. Vajda, "On divergence and informations in statistics and information theory," *IEEE Trans. Inform. Theory*, vol. 52, no. 10, pp. 4394 – 4412, Oct. 2006.
- [2] T. R. C. Read and N. Cressie, *Goodness of Fit Statistics for Discrete Multivariate Data*. Berlin: Springer, 1988.
- [3] P. Harremoës and F. Topsøe, "Inequalities between entropy and index of coincidence derived from information diagrams," *IEEE Trans. Inform. Theory*, vol. 47, no. 7, pp. 2944–2960, Nov. 2001. [Online]. Available: <http://www.math.ku.dk/topsoe/InfDiagrams.pdf>
- [4] R. J. Serfling, *Approximation Theorems of Mathematical Statistics*. New York: Wiley, 1980.
- [5] Y. Nikulin, *Asymptotic Efficiency of Nonparametric Tests*. Cambridge: Cambridge University Press, 1995.
- [6] M. P. Quine and J. Robinson, "Efficiencies of chi-square and likelihood ratio goodness-of-fit tests." *Ann. Statist.*, vol. 13, pp. 727–742, 1985.
- [7] R. R. Bahadur, *Some Limit Theorems in Statistics*. Philadelphia: SIAM, 1981.
- [8] S. H. Tumanyan, "On the asymptotic distribution of the chi-square criterion (in russian)." *Dokl. Akad. Nauk SSSR*, vol. 94, pp. 1011–1012, 1954, in Russian.
- [9] C. Morris, "Central limit theorems for multinomial sums." *Ann. Statist.*, vol. 3, pp. 165–188, 1975.
- [10] T. Inglot, T. Jurlewicz, and T. Ledwina, "Asymptotics for multinomial goodness of fit tests for simple hypothesis.35," *Th. of Probab. and Appl.*, vol. 35, pp. 797–803, 1990.
- [11] M. C. Pardo and I. Vajda, "On asymptotic properties of information-theoretic divergences." *IEEE Trans. Inform. Theory*, vol. 49, no. 7, pp. 1860–1867, 2003.
- [12] L. Györfi and I. Vajda, "Asymptotic distributions for goodness-of-fit statistics in a sequence of multinomial models," *Probability and Statistics Letters*, vol. 56, no. 1, pp. 57–67, 2002.
- [13] I. Vajda, "Asymptotic laws for stochastic disparity statistics," *Tatra Mount. Mathem. Publ.*, vol. 26, pp. 269–280, 2003, to be published.
- [14] L. Holst, "Asymptotic normality and efficiency for certain goodness-of-fit tests," *Biometrika*, vol. 59, pp. 138–145, 1972.
- [15] G. I. Ivchenko and Y. I. Medvedev, "Separable statistics and hypotheses testing. the case of small samples." *Th. of Probab. and Appl.*, vol. 23, pp. 764–775, 1978.
- [16] M. Menéndez, D. Morales, L. Pardo, and I. Vajda, "Asymptotic distributions of f-divergences of hypothetical and observed frequencies on refined partitions." *Statistica Neerlandica*, vol. 52, pp. 71–89, 1998.
- [17] J. Beirlant, L. Devroye, L. Györfi, and I. Vajda, "Large deviations of divergence measures on partitions," *J. Statist. Planning and Infer.*, vol. 93, pp. 1–16, 2001.
- [18] L. Györfi, G. Morvai, and I. Vajda, "Asymptotic results for information-theoretic and statistical criteria of goodness of fit," Institute of Information Theory and Automation, Prague., Res. Rep. 1961, 2000.
- [19] I. Vajda, "On convergence of information contained in quantized observations," *IEEE Trans. Inform. Theory*, vol. 48, no. 8, pp. 2163–2172, Aug. 2002.
- [20] W. C. M. Kallenberg, "On moderate and large deviations in multinomial distributions." *Ann. Statist.*, vol. 13, pp. 1554–1580, 1985.
- [21] I. N. Sanov, "On the probability of large deviations of random variables," *Matemat. Sbornik*, vol. 42, pp. 11–44, 1957.
- [22] I. Csiszár and J. Körner, *Information Theory: Coding Theorems for Discrete Memoryless Systems*. New York: Academic, 1981.
- [23] A. Dembo and O. Zeitouni, *Large Deviations Techniques and Applications*. Boston: Jones and Bartlett Publishers International, 1993.
- [24] L. Györfi, G. Morvai, and I. Vajda, "Information-theoretic methods in testing the goodness-of-fit," in *Proc. International Symposium on Information Theory, Sorrento, Italy, June 25-30, 2000*, p. 28.
- [25] L. Györfi and I. Vajda, "A class of modified Pearson and Neyman statistics," *Statistics and Decisions*, vol. 19, pp. 239–251, 2001.
- [26] W. C. M. Kallenberg, J. Oosterhof, and B. F. Shriever, "The number of classes in chi-squared goodness-of-fit tests." *J. Amer. Statist. Assoc.*, vol. 80, pp. 959–968, 1985.

Peter Harremoës (M'00) was born in 1964. He became master of science from University of Copenhagen in 1989 with mathematics as major and archaeology as minor subject. He received his Ph.D. from Roskilde University 1993. From 2001 to 2005 he had a post. doc. position at Mathematical Department, University of Copenhagen. Since 2006 he has been scientific staff member at Centrum voor Wiskunde en Informatica, Amsterdam and is working in the group Quantum Computing and Information Theoretic Learning. He is associate editor of *Entropy*.

Igor Vajda (M'90-SM'91-F'01) was born in Martin, Czechoslovakia. He graduated in mathematics from the Czech Technical University, Prague, 1965. He received the Ph.D. degree in probability and statistics from the Charles University, Prague, in 1968, and the Dr.Sci. degree in mathematical informatics from the same university in 1990.

From 1965 he served as a Research Assistant, from 1966 as a Researcher, and from 1990 as Director of Research at the Institute of Information Theory and Automation, Czech Academy of Sciences, Prague. During 1966-2003, he was for short periods of time a Visiting Scientist or a Visiting Professor at the Institute of Information Transmission Theory, Moscow, U.S.S.R.; University of Rostock, Germany; Complutense University of Madrid, Spain; University of Maryland, Baltimore; Catholic University of Leuven, Belgium; Technical University of Budapest, Hungary; University of Montpellier II, France; and M. Hernandez University of Alicante, Spain. Recently, he also served as a Vice-Chairman of the Board for doctoral studies in mathematical engineering at the Czech Technical University in Prague. His research interests include information measures and statistical distances, information-theoretic statistical inference, statistical inference about random processes, and statistical decisions.

Dr. Vajda is an Associate Editor of *Kybernetika*, *Revista Matematica Complutense*, *Test*, and *Applications of Mathematics*. In 1988, he was awarded the Prize of the Czechoslovak Academy of Sciences for his part in a research on speech coding. *The American Journal of Mathematical and Management Sciences* awarded him in the 1997 Jacob Wolfowitz Prize for a joint paper with L. Györfi and E. C. van der Meulen. Since 1990, he has co-organized the Prague Conferences on Information Theory, Statistical Decision Functions, and Random Processes and in 2002, he co-organized the European Meeting of Statisticians in Prague.