

## THE DOUBLE COVERING OF THE QUANTUM GROUP $SO_q(3)$ <sup>†</sup>

Mathijs S. Dijkhuizen

A quantum analogue of the double covering of  $SO(3)$  by  $SU(2)$  is formulated and proved. Here the quantum group  $SO_q(3)$  is defined by means of the  $R$ -matrix given by FRT for root systems of type  $B$ . An explicit basis for the deformed function algebra of  $SO_q(3)$  is constructed as well as an algorithm to reduce any expression in the generators to a linear combination of basis elements.

### 1. The adjoint group of $SU_q(2)$

We shall make use of both the language of Hopf algebras and that of quantum groups. We view a Hopf algebra  $A = \mathcal{O}(G)$  as the algebra of polynomial functions on an (algebraic) quantum group  $G = \text{Spec}(A)$ . Hopf  $*$ -algebras then correspond to real forms of quantum groups. A morphism  $\phi: G \rightarrow G'$  of quantum groups resp. real quantum groups is by definition a morphism  $\phi: A' \rightarrow A$  of Hopf algebras resp. Hopf  $*$ -algebras. In order to be able to distinguish formally between these two kinds of morphisms, we shall usually write  $\phi^\#$  for the Hopf algebra morphism dual to the quantum group morphism  $\phi$ .

We recall the definition of the quantum group  $SU_q(2)$ . Let  $q \in \mathbb{R}$ ,  $q \neq 0$ . The algebra  $A_q = \mathcal{O}(SU_q(2))$  is the complex unital associative algebra generated by  $\alpha, \beta, \gamma, \delta$  subject to the following relations:

$$\begin{aligned} \alpha\beta &= q\beta\alpha, & \alpha\gamma &= q\gamma\alpha, & \beta\gamma &= \gamma\beta, & \beta\delta &= q\delta\beta, & \gamma\delta &= q\delta\gamma, \\ \delta\alpha - q^{-1}\beta\gamma &= 1, & \alpha\delta - q\beta\gamma &= 1. \end{aligned} \tag{1.1}$$

By using the diamond lemma one can prove that a linear basis of  $A_q$  is formed by the elements  $\alpha^k\beta^l\gamma^m$  ( $k, l, m \geq 0$ ) and  $\delta^k\beta^l\gamma^m$  ( $k \geq 1, l, m \geq 0$ ). See [B], [Kl].

---

<sup>†</sup> This paper is in final form and no version of it will be submitted for publication elsewhere.

The comultiplication  $\Delta$  and counit  $\varepsilon$  are defined by:

$$\Delta \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \otimes \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}, \quad \varepsilon \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (1.2)$$

this being shorthand notation for  $\Delta(\alpha) = \alpha \otimes \alpha + \beta \otimes \gamma$  etc.

The antipode  $S$  and the Hopf algebra involution  $*$  are defined by

$$S \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} \delta & -q^{-1}\beta \\ -q\gamma & \alpha \end{pmatrix}, \quad \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}^* = \begin{pmatrix} \delta & -q\gamma \\ -q^{-1}\beta & \alpha \end{pmatrix}. \quad (1.3)$$

Let us recall that a mapping  $*$ :  $A_q \rightarrow A_q$  is called a Hopf algebra involution if  $(A_q, *)$  is a unital  $*$ -algebra such that  $\Delta$  and  $\varepsilon$  are  $*$ -homomorphisms.

We recall that a (matrix) corepresentation of  $A_q$  is a matrix  $(t_{ij})$  with coefficients in  $A_q$  such that

$$\Delta(t_{ij}) = \sum_k t_{ik} \otimes t_{kj}, \quad \varepsilon(t_{ij}) = \delta_{ij}. \quad (1.4)$$

A corepresentation  $(t_{ij})$  is called *unitary* if  $t_{ij}^* = S(t_{ji})$ . Corepresentations of  $A_q$  are also called *representations* of the quantum group  $SU_q(2)$ . The finite-dimensional representation theory of  $SU_q(2)$  is known to be exactly analogous to the classical theory. With respect to a suitable basis the corepresentation of  $A_q$  corresponding to the adjoint representation of  $SU(2)$  is given by (see [Ko]):

$$\text{Ad}_q = \begin{pmatrix} \alpha^2 & (1+q^{-2})^{\frac{1}{2}}\alpha\beta & -\beta^2 \\ (1+q^{-2})^{\frac{1}{2}}\alpha\gamma & 1+(q+q^{-1})\beta\gamma & -(1+q^2)^{\frac{1}{2}}\delta\beta \\ -\gamma^2 & -(1+q^2)^{\frac{1}{2}}\delta\gamma & \delta^2 \end{pmatrix}. \quad (1.5)$$

It is easily checked that the subalgebra  $B_q$  of  $A_q$  generated by the matrix coefficients  $t_{ij}$  of  $\text{Ad}_q$  is spanned by all  $\alpha^k \beta^l \gamma^m$ ,  $\delta^k \beta^l \gamma^m$  such that  $k+l+m$  is even.  $B_q$  is obviously invariant under  $S$  and  $*$ . Moreover,  $\Delta(B_q) \subset B_q \otimes B_q$ , since  $B_q$  is generated by the matrix coefficients of a corepresentation. We conclude that  $B_q$  is a Hopf  $*$ -subalgebra of  $A_q$ . The quantum group corresponding to  $B_q$  is called the adjoint group of  $SU_q(2)$  and denoted by  $\text{Ad}(SU_q(2))$ .

We define an algebra anti-automorphism  $\sigma$  of  $A_q$  and an algebra isomorphism  $\tau: A_q \rightarrow A_{q^{-1}}$  by putting

$$\sigma \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} \delta & -\gamma \\ -\beta & \alpha \end{pmatrix}, \quad \tau \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} \delta & -\beta \\ -\gamma & \alpha \end{pmatrix}. \quad (1.6)$$

It is obvious that  $\sigma$  leaves  $B_q$  invariant and that  $\tau$  maps  $B_q$  onto  $B_{q^{-1}}$ . On the generators  $t_{ij}$  the mappings  $\sigma$  and  $\tau$  are given by

$$\sigma(T) = \begin{pmatrix} t_{33} & t_{32} & t_{31} \\ t_{23} & t_{22} & t_{21} \\ t_{13} & t_{12} & t_{11} \end{pmatrix}, \quad \tau(T) = \begin{pmatrix} t_{33} & t_{23} & t_{13} \\ t_{32} & t_{22} & t_{12} \\ t_{31} & t_{21} & t_{11} \end{pmatrix}, \quad (1.7)$$

$T$  being shorthand notation for  $(t_{ij})$ . Note that  $\sigma^2 = \tau^2 = \text{id}$  and  $\tau\sigma\tau\sigma = \text{id}$ .

Table 1

1.  $t_{11}t_{12} = s^2t_{12}t_{11}$   
 $t_{32}t_{33} = s^2t_{33}t_{32}$   
 $t_{21}t_{11} = s^{-2}t_{11}t_{21}$   
 $t_{33}t_{23} = s^{-2}t_{23}t_{33}$
2.  $t_{12}t_{13} = s^2t_{13}t_{12}$   
 $t_{31}t_{32} = s^2t_{32}t_{31}$   
 $t_{23}t_{13} = s^{-2}t_{13}t_{23}$   
 $t_{31}t_{21} = s^{-2}t_{21}t_{31}$
3.  $t_{11}t_{13} = s^4t_{13}t_{11}$   
 $t_{31}t_{33} = s^4t_{33}t_{31}$   
 $t_{33}t_{13} = s^{-4}t_{13}t_{33}$   
 $t_{31}t_{11} = s^{-4}t_{11}t_{31}$
4.  $t_{22}t_{11} = t_{11}t_{22} + (s^{-2} - s^2)t_{12}t_{21}$   
 $t_{33}t_{22} = t_{22}t_{33} + (s^{-2} - s^2)t_{23}t_{32}$
5.  $t_{23}t_{12} = t_{12}t_{23} + (s^{-2} - s^2)t_{13}t_{22}$   
 $t_{32}t_{21} = t_{21}t_{32} + (s^{-2} - s^2)t_{22}t_{31}$
6.  $t_{21}t_{12} = t_{12}t_{21}$   
 $t_{32}t_{23} = t_{23}t_{32}$
7.  $t_{22}t_{13} = t_{13}t_{22}$   
 $t_{31}t_{22} = t_{22}t_{31}$
8.  $t_{23}t_{11} = s^{-2}t_{11}t_{23} + (s^{-2} - s^2)t_{13}t_{21}$   
 $t_{33}t_{21} = s^{-2}t_{21}t_{33} + (s^{-2} - s^2)t_{23}t_{31}$   
 $t_{32}t_{11} = s^{-2}t_{11}t_{32} + s^{-2}(s^{-2} - s^2)t_{12}t_{31}$   
 $t_{33}t_{12} = s^{-2}t_{12}t_{33} + s^{-2}(s^{-2} - s^2)t_{13}t_{32}$
9.  $t_{22}t_{12} = t_{12}t_{22} + s(s^2 - s^{-2})t_{13}t_{21}$   
 $t_{32}t_{22} = t_{22}t_{32} + s(s^2 - s^{-2})t_{23}t_{31}$   
 $t_{21}t_{22} = t_{22}t_{21} + s^{-1}(s^{-2} - s^2)t_{12}t_{31}$   
 $t_{22}t_{23} = t_{23}t_{22} + s^{-1}(s^{-2} - s^2)t_{13}t_{32}$
10.  $t_{21}t_{13} = s^2t_{13}t_{21}$   
 $t_{31}t_{23} = s^2t_{23}t_{31}$   
 $t_{31}t_{12} = s^{-2}t_{12}t_{31}$   
 $t_{32}t_{13} = s^{-2}t_{13}t_{32}$
11.  $t_{33}t_{11} = t_{11}t_{33} + (s - s^{-1})t_{23}t_{21} + (s - s^{-1})t_{12}t_{32}$   
 $t_{31}t_{13} = t_{13}t_{31}$
12.  $t_{31}t_{13} = t_{13}t_{31}$
13.  $t_{21}t_{23} = s^2t_{23}t_{21} + s(s^{-2} - s^2)t_{13}t_{31}$   
 $t_{32}t_{12} = s^{-2}t_{12}t_{32} + s^{-1}(s^2 - s^{-2})t_{13}t_{31}$
14.  $t_{12}t_{12} = -s^2(s + s^{-1})t_{13}t_{11}$   
 $t_{21}t_{21} = -s^{-2}(s + s^{-1})t_{11}t_{31}$   
 $t_{32}t_{32} = -s^2(s + s^{-1})t_{33}t_{31}$   
 $t_{23}t_{23} = -s^{-2}(s + s^{-1})t_{13}t_{33}$
15.  $t_{11}t_{23} = -st_{12} + s^4t_{13}t_{21}$   
 $t_{21}t_{33} = -st_{32} + s^4t_{23}t_{31}$   
 $t_{11}t_{32} = -st_{21} + s^2t_{12}t_{31}$   
 $t_{12}t_{33} = -st_{23} + s^2t_{13}t_{32}$
16.  $t_{23}t_{22} = t_{23} - s^{-2}(s + s^{-1})t_{13}t_{32}$   
 $t_{22}t_{21} = t_{21} - s^{-2}(s + s^{-1})t_{12}t_{31}$   
 $t_{12}t_{22} = t_{12} - s^2(s + s^{-1})t_{13}t_{21}$   
 $t_{22}t_{32} = t_{32} - s^2(s + s^{-1})t_{23}t_{31}$
17.  $t_{21}t_{32} = t_{31} + s^2t_{22}t_{31}$   
 $t_{12}t_{23} = t_{13} + s^2t_{13}t_{22}$
18.  $t_{11}t_{22} = t_{11} + s^2t_{12}t_{21}$   
 $t_{22}t_{33} = t_{33} + s^2t_{23}t_{32}$
19.  $t_{12}t_{32} = s - st_{22} - s^2(s + s^{-1})t_{13}t_{31}$   
 $t_{23}t_{21} = s^{-1} - s^{-1}t_{22} - s^{-2}(s + s^{-1})t_{13}t_{31}$
20.  $t_{11}t_{33} = (1 - s^2) + s^2t_{22} + s^4t_{13}t_{31}$
21.  $t_{22}t_{22} = 2t_{22} - 1 + (s + s^{-1})^2t_{13}t_{31}$

**Lemma 1.1** — *The  $t_{ij} \in B_q$  satisfy all the relations listed in Table 1 with  $s = q$ .*

In order to minimize the amount of calculation we use the symmetry inherent in the relations of Table 1. In fact, in a given cluster  $i$ , one can obtain all the relations by repeatedly applying  $\sigma$  and  $\tau$  to the first relation of that cluster (in some cases one has to work modulo linear combinations of relations of clusters  $k < i$ ). It suffices therefore to check only the first relation of each cluster. This is straightforward using (1.5) and (1.1).

**Theorem 1.2** — *The  $t_{ij} \in B_q$  satisfy no other relations than those listed in Table 1 ( $s = q$ ). A linear basis of the vector space  $B_q$  is formed by the elements*

$$\begin{aligned} t_{13}^m t_{22} t_{31}^n & \quad (m, n \geq 0) \\ t_{13}^m t_{12}^i t_{11}^j t_{21}^k t_{31}^n & \quad (m, j, n \geq 0, 0 \leq i, k \leq 1) \\ t_{13}^m t_{23}^i t_{33}^j t_{32}^k t_{31}^n & \quad (m, j, n \geq 0, 0 \leq i, k \leq 1, i + j + k > 0) \end{aligned} \quad (1.8)$$

It is clear from (1.5) that the elements (1.8) span  $B_q$ . Their linear independence immediately follows from (1.5) and the fact that the elements  $\alpha^k \beta^l \gamma^m$  ( $k, l, m \geq 0$ ) and  $\delta^k \beta^l \gamma^m$  ( $k > 0, l, m \geq 0$ ) are linearly independent in  $A_q$ . Let now  $D$  be the abstract algebra generated by the  $t_{ij}$  subject to the relations in Table 1. It follows from [1.1]<sup>†</sup> that there is a unique algebra homomorphism  $\phi: D \rightarrow B_q$  sending  $t_{ij} \in D$  to  $t_{ij} \in B_q$ . This homomorphism is surjective, since the  $t_{ij}$  generate  $B_q$ . We prove that the elements (1.8) span  $D$ . To this end, we introduce a total ordering on the generators by putting  $t_{13} < t_{12} < t_{11} < t_{23} < t_{22} < t_{21} < t_{33} < t_{32} < t_{31}$ . We then order any two given monomials in the  $t_{ij}$  by length and, if they are of equal length, lexicographically with respect to the above ordering on the  $t_{ij}$ . Inspection of the relations in Table 1 shows that all of them express a monomial  $f$  in the  $t_{ij}$  as a linear combination of monomials strictly less than  $f$ . This implies that any monomial in the  $t_{ij}$  can be expressed as a linear combination of monomials  $t_{i_1} t_{i_2} \dots t_{i_n}$  such that for all  $1 \leq j \leq n - 1$  the monomial  $t_{i_j} t_{i_{j+1}}$  does not occur on the left-hand side of any equation in Table 1. It can be easily read off from the relations in Table 1 that the monomials satisfying this last condition are precisely the ones listed in (1.8). This proves that the elements (1.8) span  $D$ . So  $\phi$  maps a family of vectors that span  $D$  to a linearly independent family of vectors in  $B_q$ . This implies that  $\phi$  is injective, which concludes the proof of the theorem.

The relations in Table 1 thus form a presentation of the algebra  $B_q$ . One easily derives from (1.5) and (1.3) that the involution  $*$  is given on the generators  $t_{ij}$  by

$$T^* = \begin{pmatrix} t_{33} & qt_{32} & q^2 t_{31} \\ q^{-1} t_{23} & t_{22} & qt_{21} \\ q^{-2} t_{13} & q^{-1} t_{12} & t_{11} \end{pmatrix}. \quad (1.9)$$

**Remark 1.3** — We use the terminology of [B]. The semigroup ordering on the monomials in the  $t_{ij}$  defined in the proof of [1.2] clearly satisfies the descending chain condition and is compatible with the reduction system specified by Table 1. It follows from [1.2] (without actually resolving a single ambiguity!) that all the ambiguities are resolvable. Therefore, the diamond lemma applies and we get an algorithm to reduce any monomial in the  $t_{ij}$  to

<sup>†</sup> Numbers between square brackets  $[\ ]$  refer to lemmas, propositions, theorems etc.

its (unique) expression in terms of the basis elements (1.8). This algorithm can be easily implemented on a computer using a computer algebra package for symbolic manipulation. We found that the package Reduce (version 3.4) was best suited to our purposes. In this way, one can perform explicit computations in the algebra  $B_q$  that would have been tiresome to do by hand.

We shall now briefly deliberate upon notions such as quantum subgroup, kernel and short exact sequence (cf. [PW]).

Let  $A$  be a Hopf algebra. A subspace  $\mathfrak{a}$  is called a two-sided coideal if  $\Delta(\mathfrak{a}) \subset A \otimes \mathfrak{a} + \mathfrak{a} \otimes A$ . A subspace  $\mathfrak{a}$  is called a Hopf ideal if  $\mathfrak{a}$  is an ideal and an  $S$ -invariant two-sided coideal. If  $\mathfrak{a}$  is a Hopf ideal, then  $A/\mathfrak{a}$  naturally inherits a Hopf algebra structure from  $A$ . A quantum group  $H$  is said to be a quantum subgroup of  $G$  if  $\mathcal{O}(H)$  is the quotient of  $\mathcal{O}(G)$  by a Hopf ideal  $\mathfrak{a}$ , which is then called the defining ideal of  $H$ . So the quantum subgroups of a given quantum group  $G$  are in 1-1 correspondence with the Hopf ideals in  $\mathcal{O}(G)$ .

Suppose  $\phi: G \rightarrow G'$  is a morphism of quantum groups and let  $\phi^\#: \mathcal{O}(G') \rightarrow \mathcal{O}(G)$  be the corresponding morphism of Hopf algebras. Define  $\mathfrak{a}$  to be the ideal in  $\mathcal{O}(G)$  generated by the image under  $\phi^\#$  of  $\ker(\varepsilon') \subset \mathcal{O}(G')$ . It is trivial that  $\mathfrak{a}$  is a Hopf ideal. The quantum subgroup of  $G$  corresponding to  $\mathfrak{a}$  is called the kernel of the morphism  $\phi$  and denoted  $\ker(\phi)$ . A sequence of morphisms of quantum groups

$$1 \longrightarrow H \xrightarrow{i} G \xrightarrow{\phi} G' \longrightarrow 1 \quad (1.10)$$

is called exact if  $\phi^\#$  is injective and  $i^\#$  surjective, and if  $\ker(i^\#)$  is the defining ideal of  $\ker(\phi)$ .

We now apply the above terminology to the adjoint group of  $SU_q(2)$ . We have a morphism  $\phi: SU_q(2) \rightarrow \text{Ad}(SU_q(2))$  such that  $\phi^\#$  is the canonical injection.

**Proposition 1.4** — *The ideal in  $A_q$  generated by  $\ker(\varepsilon|_{B_q})$  is equal to the ideal generated by  $\beta, \gamma, \alpha^2 - 1, \alpha - \delta$ .*

Let us write  $\mathfrak{a}$  for the ideal generated by  $\beta, \gamma, \alpha^2 - 1, \alpha - \delta$ . It follows from (1.2) that  $\ker(\varepsilon)$  is spanned by the elements  $\alpha^k - 1, \delta^k - 1, \alpha^k \beta^l \gamma^m, \delta^k \beta^l \gamma^m$  ( $k \geq 0, l + m > 0$ ) and generated as an ideal by  $\beta, \gamma, \alpha - 1, \delta - 1$ . So  $\ker(\varepsilon|_{B_q})$  is generated as ideal in  $B_q$  by  $\alpha^2 - 1, \delta^2 - 1, \alpha\beta, \alpha\gamma, \beta\gamma, \delta\beta, \delta\gamma$ . This implies that  $\mathfrak{a}$  contains  $\ker(\varepsilon|_{B_q})$ . On the other hand, multiplying  $\alpha^2 - 1$  on the right by  $\delta$ , we get  $\alpha(\alpha\delta) - \delta = \alpha(1 + q\beta\gamma) - \delta = \alpha - \delta + q\alpha\beta\gamma$ . Hence  $\alpha - \delta$  lies in the ideal generated by  $\ker(\varepsilon|_{B_q})$ . Multiplying  $\alpha\beta$  on the left by  $\delta$ , we obtain  $\delta\alpha\beta = \beta + q^{-1}\beta^2\gamma$ . This implies that  $\beta$  lies in the ideal generated by  $\ker(\varepsilon|_{B_q})$ . One reasons similarly for  $\gamma$ . The assertion follows.

The quotient  $A/\mathfrak{a}$  is generated by  $\alpha$  subject to the relation  $\alpha^2 = 1$ . This implies that it is a two-dimensional Hopf  $*$ -algebra. In fact,  $A/\mathfrak{a}$  is the Hopf  $*$ -algebra of functions on the finite group  $\mathbb{Z}_2$ . We now have proved:

**Theorem 1.5** — *There is the following short exact sequence of quantum groups:*

$$1 \longrightarrow \mathbb{Z}_2 \longrightarrow SU_q(2) \xrightarrow{\phi} \text{Ad}(SU_q(2)) \longrightarrow 1.$$

This justifies our writing  $\text{Ad}(SU_q(2)) = SU_q(2)/\{1, -1\}$ .

## 2. The quantum group $SO_q(3)$

We adopt the definition of  $SO_q(3)$  given in [FRT] with the one exception that we add a determinant relation (see also [T2]). In fact, the quantum group defined in [FRT] is a quantization of  $O_q(3)$  and not of  $SO_q(3)$ .

Let  $(t_{ij})$  ( $1 \leq i, j \leq 3$ ) denote a family of formal indeterminates and let  $\mathbb{C}\langle t_{ij} \rangle$  denote the free associative unital complex algebra generated by the  $t_{ij}$ . It will be convenient to arrange the indeterminates  $t_{ij}$  in matrix form  $T = (t_{ij})$ . We can then define  $9 \times 9$  matrices  $T_1$  resp.  $T_2$  with coefficients in the algebra  $\mathbb{C}\langle t_{ij} \rangle$  by putting  $T_1 = T \otimes I$  resp.  $T_2 = I \otimes T$ . Here  $I$  denotes the  $3 \times 3$  identity matrix.

Let  $e_1, e_2, e_3$  be the canonical basis of  $V = \mathbb{C}^3$ . Then  $V \otimes V$  has the basis  $e_i \otimes e_j$  ( $1 \leq i, j \leq 3$ ). Let  $e_{ij}$  denote the linear endomorphism of  $V$  sending  $e_j$  to  $e_i$  and  $e_k$  ( $k \neq j$ ) to 0. Let  $q > 0$ ,  $q \neq 1$ . We define a linear endomorphism  $R_q$  of  $V \otimes V$  or, equivalently, a  $9 \times 9$  matrix  $(R_{ij,kl}^q)$  with complex coefficients by putting:

$$R_q = q \sum_{i \neq i'} e_{ii} \otimes e_{ii} + e_{22} \otimes e_{22} + \sum_{i \neq j, j'} e_{ji} \otimes e_{ij} + \\ + q^{-1} \sum_{i \neq i'} e_{ii'} \otimes e_{i'i} + (q - q^{-1}) \sum_{i > j} e_{jj} \otimes e_{ii} - (q - q^{-1}) \sum_{i > j} q^{\rho_i - \rho_j} e_{i'j} \otimes e_{ij'}.$$

Here  $i' = 4 - i$  and the sequence  $(\rho_i)$  is defined as  $(\rho_1, \rho_2, \rho_3) = (\frac{1}{2}, 0, -\frac{1}{2})$ . Note that the matrix  $(R_{ij,kl}^q)$  is symmetric and hence diagonalizable. Straightforward computation shows that the eigenvalues of  $R_q$  are  $q, -q^{-1}, q^{-2}$  and that a basis of eigenvectors for  $V \otimes V$  is given by:

$$W_q : \begin{aligned} & e_1 \otimes e_1, e_3 \otimes e_3 \\ & qe_1 \otimes e_2 + e_2 \otimes e_1, qe_2 \otimes e_3 + e_3 \otimes e_2 \\ & qe_1 \otimes e_3 - (q^{\frac{1}{2}} + q^{-\frac{1}{2}})e_2 \otimes e_2 + q^{-1}e_3 \otimes e_1 \end{aligned}$$

$$W_{-q^{-1}} : \begin{aligned} & e_1 \otimes e_2 - qe_2 \otimes e_1, e_2 \otimes e_3 - qe_3 \otimes e_2 \\ & e_1 \otimes e_3 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})e_2 \otimes e_2 - e_3 \otimes e_1 \end{aligned} \quad (2.1)$$

$$W_{q^{-2}} : q^{-\frac{1}{2}}e_1 \otimes e_3 + e_2 \otimes e_2 + q^{\frac{1}{2}}e_3 \otimes e_1$$

We consider the two-sided ideal  $I_q$  in  $\mathbb{C}\langle t_{ij} \rangle$  generated by the relations (called *commutation relations*) coming from the matrix equation

$$R_q T_1 T_2 = T_1 T_2 R_q. \quad (2.2)$$

Equating the  $9 \times 9$  coefficients on both sides yields the following explicit form for the defining relations of  $I_q$ :

$$\sum_{m,n} R_{ij,mn}^q t_{mk} t_{nl} = \sum_{m,n} t_{im} t_{jn} R_{mn,kl}^q \quad \text{for } 1 \leq i, j, k, l \leq 3.$$

Let  $V^*$  denote the linear dual of  $V$  with dual basis  $e_i^*$ . We define a linear mapping  $\theta: V^* \otimes V^* \otimes V \otimes V \rightarrow \mathbb{C}\langle t_{ij} \rangle$  by putting

$$\theta(e_i^* \otimes e_j^* \otimes e_k \otimes e_l) = t_{ik}t_{jl}. \quad (2.3)$$

Let  $R_q^*: V^* \otimes V^* \rightarrow V^* \otimes V^*$  denote the transpose of  $R_q$ . It is clear that it is diagonalizable with the same eigenvalues as  $R_q$ . Since the matrix  $(R_{ij,kl}^q)$  is symmetric, bases for the eigenspaces  $W_\lambda^*$  of  $R_q^*$  are obtained by replacing  $e_i$  by  $e_i^*$  in (2.1). We now have:

**Proposition 2.1** — *The ideal  $I_q$  is generated by the subspace  $\bigoplus_{\lambda \neq \mu} \theta(W_\lambda^* \otimes W_\mu)$  of  $\mathbb{C}\langle t_{ij} \rangle$ .*

Let us write  $\xi = \theta \circ (\text{id} \otimes R_q - R_q^* \otimes \text{id})$ . On the one hand,  $I_q$  is generated by the image of  $\xi$  since  $\xi(e_i^* \otimes e_j^* \otimes e_k \otimes e_l) = \sum_{mn} R_{mn,ki}^q t_{im}t_{jn} - \sum_{mn} R_{ij,mn}^q t_{mk}t_{nl}$ . On the other hand,  $\xi(W_\lambda^* \otimes W_\mu) = (\mu - \lambda)\theta(W_\lambda^* \otimes W_\mu)$ . We conclude that  $I_q$  is generated by  $\bigoplus_{\lambda \neq \mu} \theta(W_\lambda^* \otimes W_\mu)$ , since  $R_q^*$  is diagonalizable.

The quotient algebra  $M_q = \mathbb{C}\langle t_{ij} \rangle / I_q$  is called the *algebra of polynomial functions on the complex orthogonal quantum matrix space of rank 3*. We indifferently write  $t_{ij}$  for the generators in  $\mathbb{C}\langle t_{ij} \rangle$  or their canonical images in  $M_q$ .

There are unique algebra homomorphisms  $\Delta: M_q \rightarrow M_q \otimes M_q$  and  $\varepsilon: M_q \rightarrow \mathbb{C}$  such that

$$\Delta(t_{ij}) = \sum_k t_{ik} \otimes t_{kj}, \quad \varepsilon(t_{ij}) = \delta_{ij} \quad (i, j = 1, 2, 3).$$

With these mappings  $M_q$  becomes a bialgebra.

We now define the *quantized orthogonal exterior algebra*  $\Lambda_q V$  to be the tensor algebra over  $V$  divided out by the ideal generated by the subspace  $\overline{W}_q = W_q \oplus W_{q^{-2}} \subset V \otimes V$ . It follows from (2.1) that a basis of  $\overline{W}_q$  is formed by

$$\begin{aligned} &e_1 \otimes e_1, e_3 \otimes e_3, (q^{\frac{1}{2}} - q^{-\frac{1}{2}})e_1 \otimes e_3 - e_2 \otimes e_2 \\ &qe_1 \otimes e_2 + e_2 \otimes e_1, qe_2 \otimes e_3 + e_3 \otimes e_2, e_1 \otimes e_3 + e_3 \otimes e_1. \end{aligned} \quad (2.4)$$

So  $\Lambda_q V$  is the algebra generated by  $e_1, e_2, e_3$  subject to the relations

$$e_1^2 = 0, e_3^2 = 0, e_2e_1 = -qe_1e_2, e_3e_2 = -qe_2e_3, e_3e_1 = -e_1e_3, e_2^2 = (q^{\frac{1}{2}} - q^{-\frac{1}{2}})e_1e_3. \quad (2.5)$$

An application of the diamond lemma shows that  $\Lambda_q V$  is an 8-dimensional vector space with basis  $e_{i_1} \dots e_{i_n}$  ( $1 \leq i_1 < \dots < i_n \leq 3$ ). It can be shown that there is a unique algebra homomorphism  $\delta: \Lambda_q V \rightarrow M_q \otimes \Lambda_q V$  such that

$$\delta(e_i) = \sum_j t_{ij} \otimes e_j \quad (i = 1, 2, 3). \quad (2.6)$$

The mapping  $\delta$  defines a left coaction of  $M_q$  on  $\Lambda_q V$ , i.e. it satisfies the properties:

$$(\Delta \otimes \text{id}) \circ \delta = (\text{id} \otimes \delta) \circ \delta, \quad (\varepsilon \otimes \text{id}) \circ \delta = \text{id}. \quad (2.7)$$

It follows from (2.6) and (2.5) that there is a unique element  $\det_q T \in M_q$  such that

$$\delta(e_1e_2e_3) = \det_q T \otimes e_1e_2e_3. \quad (2.8)$$

The equalities (2.7) then imply that

$$\Delta(\det_q T) = \det_q T \otimes \det_q T, \quad \varepsilon(\det_q T) = 1.$$

An explicit expression for  $\det_q T$  is:

$$\begin{aligned} \det_q T = & t_{11}t_{22}t_{33} - qt_{12}t_{21}t_{33} - qt_{11}t_{23}t_{32} + qt_{12}t_{23}t_{31} \\ & + qt_{13}t_{21}t_{32} - q^2t_{13}t_{22}t_{31} - q(q^{\frac{1}{2}} - q^{-\frac{1}{2}})t_{12}t_{22}t_{32}. \end{aligned} \quad (2.9)$$

We introduce a  $3 \times 3$  matrix  $C$  given by

$$\begin{pmatrix} 0 & 0 & q^{-\frac{1}{2}} \\ 0 & 1 & 0 \\ q^{\frac{1}{2}} & 0 & 0 \end{pmatrix}.$$

It satisfies  $C^2 = I$ . The so-called *orthogonality relations* are:

$$TC^tTC = C^tTCT = I. \quad (2.10)$$

It can be proved that the two-sided ideal  $J_q$  in  $M_q$  generated by the relations  $\det_q T = 1$  and (2.10) is a biideal. We can now define the algebra  $C_q = \mathcal{O}(SO_q(3))$  of polynomial functions on the quantum group  $SO_q(3)$  as the quotient of  $M_q$  by the biideal  $J_q$ .  $C_q$  is a bialgebra by definition. It becomes a Hopf  $*$ -algebra by putting

$$S(T) = C^tTC = \begin{pmatrix} t_{33} & q^{-\frac{1}{2}}t_{23} & q^{-1}t_{13} \\ q^{\frac{1}{2}}t_{32} & t_{22} & q^{-\frac{1}{2}}t_{12} \\ qt_{31} & q^{\frac{1}{2}}t_{21} & t_{11} \end{pmatrix} \quad \text{and} \quad T^* = {}^tS(T). \quad (2.11)$$

**Theorem 2.2** — *The algebra  $C_q$  is generated by the  $t_{ij}$  subject to the relations listed in Table 1 with  $s = q^{\frac{1}{2}}$ .*

The proof is completely elementary, although not entirely trivial, since one easily gets bogged down in a quagmire of relations. For this reason, we shall carefully indicate the line of reasoning to be followed, but not explicitly perform all the calculations. The proof consists of two parts.

We first prove that the linear span  $Z_q$  of the relations (2.2) in the free algebra  $\mathbb{C}\langle t_{ij} \rangle$  is equal to the linear span of the relations (1) till (14) in Table 1 plus some extra relations (see below). To this end, we apply [2.1]. Let us identify  $V$  and  $V^*$  via the bases  $(e_i)$  and  $(e_i^*)$ . Under this identification  $W_\lambda$  coincides with  $W_\lambda^*$ . It now follows from [2.1] that

$$Z_q = \theta(\overline{W}_q \otimes W_{-q^{-1}}) \oplus \theta(W_{-q^{-1}} \otimes \overline{W}_q) \oplus \theta(W_q \otimes W_{q^{-2}}) \oplus \theta(W_{q^{-2}} \otimes W_q).$$

The following remark may be of use. If  $\omega: \mathbb{C}\langle t_{ij} \rangle \rightarrow \mathbb{C}\langle t_{ij} \rangle$  denotes the unique algebra homomorphism such that  $\omega(t_{ij}) = t_{ji}$  then  $\omega \circ \theta(W_\lambda \otimes W_\mu) = \theta(W_\mu \otimes W_\lambda)$ . We use the bases of  $\overline{W}_q$  resp.  $W_q, W_{-q^{-1}}, W_{q^{-2}}$  given in (2.4) resp. (2.1). Let us call the basis vectors in (2.1)  $w_i$  ( $1 \leq i \leq 9$ ) and those in (2.4)  $\bar{w}_j$  ( $1 \leq j \leq 6$ ), in the order in which they are introduced. Thus,  $w_1 = e_1 \otimes e_1, w_2 = e_3 \otimes e_3$  etc. Straightforward calculation shows that the tensor products (in either order) of  $\bar{w}_1, \bar{w}_2$  and  $w_6, w_7$  (meaning  $\bar{w}_1 \otimes w_6, w_6 \otimes \bar{w}_1, \bar{w}_1 \otimes w_7,$



$w_7 \otimes \bar{w}_1, \bar{w}_2 \otimes w_6$  etc.) yield precisely the relations (1) and (2) in Table 1. Similarly, from the tensor products (in either order) of  $\bar{w}_4, \bar{w}_5$  and  $w_6, w_7$  one derives precisely the relations (4), (5), (6) and (7). The relations coming from the tensor products (in either order) of  $\bar{w}_1, \bar{w}_2$  and  $w_8$  together with the relations coming from the tensor products of  $w_1, w_2$  and  $w_9$  are easily seen to be equivalent to relations (3) and (14). The tensor products  $\bar{w}_4 \otimes w_8, w_6 \otimes \bar{w}_3, w_6 \otimes \bar{w}_6$  and  $w_3 \otimes w_9$  lead to the following four equations:

$$\begin{aligned} qt_{11}t_{23} + q(q^{\frac{1}{2}} - q^{-\frac{1}{2}})t_{12}t_{22} - qt_{13}t_{21} + t_{21}t_{13} + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})t_{22}t_{12} - t_{23}t_{11} &= 0 \\ (q^{\frac{1}{2}} - q^{-\frac{1}{2}})t_{11}t_{23} - q(q^{\frac{1}{2}} - q^{-\frac{1}{2}})t_{21}t_{13} - t_{12}t_{22} + qt_{22}t_{12} &= 0 \\ t_{11}t_{23} - qt_{21}t_{13} + t_{13}t_{21} - qt_{23}t_{11} & \\ q^{\frac{1}{2}}t_{11}t_{23} + qt_{12}t_{22} + q^{\frac{3}{2}}t_{13}t_{21} + q^{-\frac{1}{2}}t_{21}t_{13} + t_{22}t_{12} + q^{\frac{1}{2}}t_{23}t_{11} &= 0. \end{aligned} \quad (2.12)$$

We can eliminate  $t_{23}t_{11}$  from the first and fourth equations of (2.12) by using the third equation. From the resulting two equations one can eliminate  $t_{11}t_{23}$  by using the second equation of (2.12). One then obtains:

$$\begin{aligned} (q^{\frac{3}{2}} + q^{-\frac{1}{2}})t_{12}t_{22} - (q^{\frac{3}{2}} + q^{-\frac{1}{2}})t_{22}t_{12} - (q + q^{-1})t_{13}t_{21} + (q^2 + 1)t_{21}t_{13} &= 0 \\ (q^{\frac{3}{2}} + q^{-\frac{1}{2}})t_{12}t_{22} - (q^{\frac{3}{2}} + q^{-\frac{1}{2}})t_{22}t_{12} + (q + q^{-1})(q - 1)t_{13}t_{21} + (q^2 - q - q^{-1} + 1)t_{21}t_{13} &= 0, \end{aligned} \quad (2.13)$$

from which one deduces (10.a). Resubstituting (10.a) in the first equation of (2.13) resp. the third equation of (2.12) one obtains (9.a) resp. (8.a). Using (9.a) and (10.a) we can rewrite the second equation of (2.12) as

$$t_{11}t_{23} = -q^{\frac{1}{2}}t_{12}t_{22} - qt_{13}t_{21}, \quad (2.14)$$

and (8.a), (9.a), (10.a) and (2.14) are in fact equivalent to (2.12). In an exactly analogous way, one derives all the (other) relations of (8), (9) and (10) by using the tensor products  $\bar{w}_4 \otimes w_8, \bar{w}_5 \otimes w_8, \bar{w}_6 \otimes w_6, \bar{w}_6 \otimes w_7, w_3 \otimes w_9, w_4 \otimes w_9$  and their images under the flip  $v \otimes w \mapsto w \otimes v$ . The analogues of (2.14) are:

$$\begin{aligned} t_{21}t_{33} &= -q^{\frac{1}{2}}t_{22}t_{32} - qt_{23}t_{31} \\ t_{22}t_{21} &= (q^{\frac{1}{2}} - q^{-\frac{1}{2}} - q^{-\frac{3}{2}})t_{12}t_{31} - q^{-\frac{1}{2}}t_{11}t_{32} \\ t_{23}t_{22} &= (q^{\frac{1}{2}} - q^{-\frac{1}{2}} - q^{-\frac{3}{2}})t_{13}t_{32} - q^{-\frac{1}{2}}t_{12}t_{33}. \end{aligned} \quad (2.15)$$

Finally, the tensor products  $\bar{w}_3 \otimes w_8, \bar{w}_6 \otimes w_8, w_5 \otimes w_9$  and their images under the flip give rise to six equations which can be seen to be equivalent to the relations (11), (12), (13) plus the following relations:

$$\begin{aligned} t_{23}t_{21} &= q^{-1}t_{12}t_{32} + q^{-\frac{1}{2}}(q - q^{-1})t_{13}t_{31} \\ t_{22}t_{22} &= t_{11}t_{33} + (q^{\frac{1}{2}} - 2q^{-\frac{1}{2}})t_{12}t_{32} + q^{-1}t_{13}t_{31}. \end{aligned} \quad (2.16)$$

Summarizing, we have now proved that the relations (2.2) are equivalent to the relations (1) till (14) plus the relations in (2.14), (2.15) and (2.16).

In the second part of the proof we shall make use of the terminology laid down in [B]. As generators we take the  $t_{ij}$ . We order the monomials in the  $t_{ij}$  as in the proof of [1.2]. As

reduction system we take the relations (1) till (14) plus the relations in (2.14), (2.15) and (2.16). Note that this reduction system is compatible with the ordering of the monomials in the  $t_{ij}$ . Given any relation  $\sigma$  in the  $t_{ij}$  we can first reduce it to an irreducible expression and then rewrite it in the form  $f = \sum_i c_i f_i$  where the  $f_i$  are monomials strictly less than the monomial  $f$ . If  $\sigma$  is of degree at most two the result of these two operations on  $\sigma$  is uniquely determined. We call it the reduced form of  $\sigma$ . It clearly is compatible with the ordering of the monomials in the  $t_{ij}$ . If we apply this procedure to the orthogonality relations (2.10) we end up with a single relation:

$$t_{11}t_{33} = 1 - qt_{13}t_{31} - q^{\frac{1}{2}}t_{12}t_{32}. \quad (2.17)$$

We add (2.17) to our reduction system. Given any ambiguity, it either is resolvable or gives rise to a new relation which can be written in reduced form and added to the reduction system. Starting from the ambiguities  $t_{32}t_{23}t_{11}$ ,  $t_{33}t_{22}t_{12}$ ,  $t_{33}t_{21}t_{11}$ ,  $t_{32}t_{22}t_{11}$ ,  $t_{33}t_{23}t_{11}$  and  $t_{33}t_{22}t_{11}$  we get:

$$\begin{aligned} t_{12}t_{23}t_{31} &= t_{13}t_{21}t_{32} \\ t_{12}t_{23}t_{32} &= (q+1)t_{13}t_{22}t_{32} + q(q^{\frac{1}{2}} + q^{-\frac{1}{2}})t_{13}t_{23}t_{31} \\ t_{11}t_{22}t_{32} &= 2q^2(q^{\frac{1}{2}} + q^{-\frac{1}{2}})t_{13}t_{21}t_{31} + q^{\frac{3}{2}}(q^{\frac{1}{2}} + 2q^{-\frac{1}{2}})t_{12}t_{22}t_{31} - q^{\frac{1}{2}}t_{21} \\ t_{12}t_{21}t_{32} &= q(q^{\frac{1}{2}} + q^{-\frac{1}{2}})t_{13}t_{21}t_{31} + (q+1)t_{12}t_{22}t_{31} \\ t_{12}t_{22}t_{33} &= 2q^2(q^{\frac{1}{2}} + q^{-\frac{1}{2}})t_{13}t_{23}t_{31} + q^{\frac{3}{2}}(q^{\frac{1}{2}} + 2q^{-\frac{1}{2}})t_{13}t_{22}t_{32} - q^{\frac{1}{2}}t_{23} \\ t_{11}t_{22}t_{33} &= (q^{-\frac{1}{2}} - q^{\frac{1}{2}} - q^{\frac{3}{2}})t_{12}t_{22}t_{32} - qt_{13}t_{22}t_{31} - 2q(q - q^{-1})t_{13}t_{21}t_{32} + t_{22}. \end{aligned} \quad (2.18)$$

We add these relations to our reduction system. The reduced form of the relation  $\det_q T = 1$  (see (2.9)) then becomes:

$$t_{12}t_{22}t_{32} = q^{\frac{1}{2}} - q^{\frac{1}{2}}t_{22} + q^2(q^{\frac{1}{2}} + q^{-\frac{1}{2}})t_{13}t_{22}t_{31} - 2q(q^{\frac{1}{2}} + q^{-\frac{1}{2}})t_{13}t_{21}t_{32}. \quad (2.19)$$

We add (2.19) to our reduction system too. We then get new ambiguities  $t_{12}t_{12}t_{22}t_{32}$  and  $t_{12}t_{22}t_{32}t_{32}$  from which one derives (16.c) and (16.d). We add these last two equations to our reduction system. (From now on, this will be done automatically every time we derive a new relation in Table 1.) The new (inclusion) ambiguities  $t_{11}t_{22}t_{32}$  and  $t_{12}t_{22}t_{33}$  then lead to (15.c) and (15.d). Since (2.19) is not irreducible anymore, we should rewrite it as:

$$t_{13}t_{21}t_{32} = (q+1)^{-1} - q^{-1}(q^{\frac{1}{2}} + q^{-\frac{1}{2}})^{-1}t_{12}t_{32} + qt_{13}t_{22}t_{31} - (q+1)^{-1}t_{22}. \quad (2.20)$$

The ambiguities  $t_{22}t_{32}t_{21}$  and  $t_{21}t_{22}t_{32}$  lead to (17.a) and then the ambiguity  $t_{13}t_{21}t_{32}$  leads to (19.a). Using the new reduction rules (19.a), (16.c/d) and (15.c/d), one sees that the relations (2.17), (2.14), (2.15) and (2.16) can be rewritten as (20), (15.a), (15.b), (16.b), (16.a), (19.b) and (21) respectively. The ambiguities  $t_{23}t_{12}t_{22}$ ,  $t_{11}t_{12}t_{32}$  and  $t_{12}t_{32}t_{33}$  now lead to (17.b), (18.a) and (18.b) respectively. It is easy to see that the reduction rules (2.18) and (2.20) can now be discarded. This concludes the proof of the theorem.

**Corollary 2.3** — *There is a unique algebra homomorphism  $\chi^\#: C_{q^2} \rightarrow B_q$  such that*

$$\chi^\#(T) = \begin{pmatrix} \alpha^2 & (1+q^{-2})^{\frac{1}{2}}\alpha\beta & -\beta^2 \\ (1+q^{-2})^{\frac{1}{2}}\alpha\gamma & 1+(q+q^{-1})\beta\gamma & -(1+q^2)^{\frac{1}{2}}\delta\beta \\ -\gamma^2 & -(1+q^2)^{\frac{1}{2}}\delta\gamma & \delta^2 \end{pmatrix}.$$

The mapping  $\chi^\sharp$  is an isomorphism of Hopf  $*$ -algebras.

It follows from [2.2] and [1.2] that  $\chi^\sharp$  is well-defined and bijective. That  $\chi^\sharp$  is a Hopf algebra morphism follows from the fact that  $\text{Ad}_q$  (see (1.5)) is a corepresentation. Finally,  $\chi^\sharp$  respects the  $*$ -operations because of (1.9) and (2.11).

We identify  $SO_{q^2}(3)$  and  $\text{Ad}(SU_q(2))$  via  $\chi$ . We then have a morphism  $\phi: SU_q(2) \rightarrow SO_{q^2}(3)$  (see above [1.4]) and [1.5] can be restated as

**Corollary 2.4** — *There is the following exact sequence of quantum groups:*

$$1 \longrightarrow \mathbb{Z}_2 \longrightarrow SU_q(2) \xrightarrow{\phi} SO_{q^2}(3) \longrightarrow 1.$$

During the winter school in Zdikov O. Ogievetsky pointed out to me that the quantum analogue of the classical isomorphism  $A_1 \sim B_1$  had already been proved in [JO] in a more general context. In [T1] a result similar to [2.3] was announced, but to my knowledge a proof of this claim has never been published.

## REFERENCES

- [B] Bergman, G.M., “The diamond lemma for ring theory”, *Advances Math.*, 29 (1978), 178–218.
- [JO] Jain, V., Ogievetsky, O., “Classical isomorphisms for quantum groups”, *Mod. Phys. Lett. A* 7 (1992) 2199.
- [Kl] Koelink, H.T., “On  $*$ -representations of the Hopf  $*$ -algebra associated with the quantum group  $U_q(n)$ ”, *Compositio Math.*, 77 (1991), 199–231.
- [Ko] Koornwinder, T.H., “Askey-Wilson polynomials as zonal spherical functions on the  $SU(2)$  quantum group”, *CWI Report AM-R9013* (1990).
- [PW] Parshall, W., Wang J., “Quantum linear groups”, *Memoirs of the American Mathematical Society*, vol. 89–439 (1991).
- [FRT] Reshetikhin, N.YU., Takhtadzhyan, L.A., Faddeev, L.D., “Quantization of Lie groups and Lie algebras”, *Leningrad Math. J.*, 1 (1) (1990), 193–225.
- [T1] Takeuchi, M., “Quantum orthogonal and symplectic groups and their embedding into quantum  $GL$ ”, *Proc. Japan Acad.*, 65 Ser. A (1989), 55–58.
- [T2] Takeuchi, M., “Matric bialgebras and quantum groups”, *Israel J. Math.*, 72 (1–2) (1990), 232–251.
- [VS] Vaksman, L.L., Soibelman, Y.S., “Algebra of functions on the quantum group  $SU(2)$ ”, *Funct. Anal. Appl.*, 22 (1988), 170–181.

CWI (CENTRE FOR MATHEMATICS AND COMPUTER SCIENCE)  
P.O. BOX 4079  
NL-1009 AB AMSTERDAM  
EMAIL THIJS@CWI.NL