

MATRIX ELEMENTS OF IRREDUCIBLE REPRESENTATIONS OF
 $SU(2) \times SU(2)$ AND VECTOR-VALUED ORTHOGONAL
POLYNOMIALS*

TOM H. KOORNWINDER[†]

Abstract. The matrix elements of irreducible representations of $SU(2) \times SU(2)$ in a $\text{diag}(SU(2) \times SU(2))$ -basis are expressed in terms of vector-valued orthogonal polynomials, which generalize the Jacobi polynomials.

0. Introduction. It is well known (cf. Vilenkin [11, Chap. 3]) that the matrix elements of the irreducible representations of $SU(2)$ in $S(U(1) \times U(1))$ -basis can be expressed in terms of Jacobi polynomials, such that the orthogonality relations for these polynomials are equivalent to Schur's orthogonality relations for the matrix elements. More generally, let G be a compact Lie group with closed subgroup K such that each irreducible representation of G , restricted to K , is multiplicity free. Consider the matrix elements of the irreducible representations of G in a K -basis. Is it possible to express them in terms of some kind of orthogonal polynomials? For the case $G = SU(2) \times SU(2)$, $K = \text{diagonal in } G$, this paper will give a positive answer. (Note that this case is a covering of the pair $(G, K) = (SO(4), SO(3))$.) The resulting polynomials are vector-valued and orthogonal on $[-1, 1]$ with respect to a positive definite matrix-valued weight function. It would be of interest to generalize these results to the cases $(G, K) = (SO(n), SO(n-1))$ or $(U(n), U(n-1))$.

The topic of this paper originated from work on the global approach to the representation theory of a noncompact semisimple Lie group G (cf. [7]) for $SL(2, \mathbb{R})$, Kosters [8] for $SL(2, \mathbb{C})$). In this approach one needs some knowledge of the matrix elements of the principal series representations of G in a K -basis (K maximal compact subgroup of G). These matrix elements have integral representations in terms of the matrix elements of irreducible representations of K (cf. (4.1) in the case $G = SL(2, \mathbb{C})$). Manipulation of these integral representations will be simplified if one can express the matrix elements for K in terms of orthogonal polynomials. Thus the results of the present paper will be useful for the analysis on $SO_0(4, 1)$.

It is the author's feeling that the highly nontrivial example of vector-valued orthogonal polynomials presented here is interesting for its own sake. Hopefully this paper will also be useful for physicists, who have already studied the matrix elements for $SO(4)$ for a long time (cf. for instance Freedman and Wang [3], Smorodinskii and Shepelev [10], Basu and Srinivasan [1]). Many authors start with the matrix elements of the principal series representations of $SO_0(3, 1)$ (cf. [1], [10]) and then obtain the matrix elements for the compact case by analytic continuation. In the present paper, with its emphasis on orthogonal polynomials, it seemed more natural to start with the compact case, but in the final §4 the noncompact analogue is briefly discussed.

Sections 1 and 2 are of a preliminary nature. In §1 matrix elements for $SU(2)$ are reviewed, both as a tool needed later and as a motivating example. In §2 Schur's orthogonality relations for matrix elements for $SU(2) \times SU(2)$ are expressed as an

*Received by the editors June 7, 1983.

[†]Centre for Mathematics and Computer Science, P.O. Box 4079, 1009 AB Amsterdam, The Netherlands.

orthogonality for vector-valued functions on $[0, \pi]$ and good candidates are selected for the expected vector-valued orthogonal polynomials. In §3 these polynomials are really obtained together with an integral representation and a power series expansion. There are two further matters of particular interest in §3: First, a trick to deform the integral of an analytic function over $SU(2)$ into the complexification $SL(2, \mathbb{C})$ by multiplication on the right of the integration variable with a particular element of $SL(2, \mathbb{C})$ (cf. the transition (3.3)→(3.6)) and, second, an unexpected symmetry (3.11) for the vector-valued polynomials.

1. The matrix elements for $SU(2)$. Let $l \in \frac{1}{2}\mathbb{Z}_+ := \{0, \frac{1}{2}, 1, \frac{3}{2}, \dots\}$. Let H_l be the space of homogeneous polynomials of degree $2l$ in two complex variables, made into a Hilbert space by the choice of orthonormal basis $\{\psi'_n | n = -l, -l+1, \dots, l\}$

$$(1.1) \quad \psi'_n(x, y) := \binom{2l}{l-n}^{1/2} x^{l-n} y^{l+n}.$$

Define a representation T^l of $GL(2, \mathbb{C})$ on H_l by

$$(1.2) \quad \left(T^l \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} f\right)(x, y) := f(\alpha x + \gamma y, \beta x + \delta y).$$

The T^l 's form a complete system of representatives for $(SU(2))^\wedge$ (cf. Vilenkin [11, Chap. 3]).

Write $T^l(g) (g \in GL(2, \mathbb{C}))$ as a matrix $(t^l_{mn}(g))$ with respect to the basis functions ψ'_n

$$(1.3) \quad T^l(g) \psi'_n = \sum_{m=-l}^l t^l_{mn}(g) \psi'_m, \quad g \in GL(2, \mathbb{C}).$$

If g is a diagonal matrix then so is $(t^l_{mn}(g))$. It follows from (1.1), (1.2), (1.3) that

$$(1.4) \quad \binom{2l}{l-n}^{1/2} (\alpha x + \gamma y)^{l-n} (\beta x + \delta y)^{l+n} = \sum_{m=l}^l t^l_{mn} \binom{2l}{l-m}^{1/2} x^{l-m} y^{l+m}.$$

Expansion of the left-hand side of (1.4) yields

$$(1.5) \quad t^l_{mn} \binom{\alpha \quad \beta}{\gamma \quad \delta} = ((l-m)!(l+m)!(l-n)!(l+n)!)^{1/2} \cdot \sum_{r=0 \vee (-n-m)}^{(l-n) \wedge (l-m)} \frac{\alpha^l \beta^{l-m-r} \gamma^{l-n-r} \delta^{m+n+r}}{r!(l-m-r)!(l-n-r)!(m+n+r)!}.$$

This implies the symmetries

$$(1.6) \quad \beta^m \gamma^n t^l_{mn} \binom{\alpha \quad \beta}{\gamma \quad \delta} = \beta^n \gamma^m t^l_{nm} \binom{\alpha \quad \beta}{\gamma \quad \delta},$$

$$(1.7) \quad t^l_{mn} \binom{\alpha \quad \beta}{\gamma \quad \delta} = t^l_{nm} \binom{\alpha \quad \gamma}{\beta \quad \delta}.$$

From (1.4) and (1.7) we obtain the integral representation

$$(1.8) \quad t_{mn}^l \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \left(\frac{(l-n)!(l+n)!}{(l-m)!(l+m)!} \right)^{1/2} \cdot \frac{1}{2\pi} \int_0^{2\pi} (\alpha e^{i\phi} + \beta e^{-i\phi})^{l-m} (\gamma e^{i\phi} + \delta e^{-i\phi})^{l+m} e^{2in\phi} d\phi.$$

The following symmetry is apparent from (1.8).

$$(1.9) \quad t_{mn}^l \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = t_{-m, -n}^l \begin{pmatrix} \delta & \gamma \\ \beta & \alpha \end{pmatrix}.$$

Now specialize to $SU(2)$. We will use the notation

$$(1.10) \quad k(\alpha, \beta) := \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \quad \text{where } |\alpha|^2 + |\beta|^2 = 1,$$

$$(1.11) \quad b_\theta := k\left(\cos \frac{\theta}{2}, \sin \frac{\theta}{2}\right),$$

$$(1.12) \quad m_\phi := k(e^{i\phi/2}, 0).$$

Note that

$$(1.13) \quad t_{mn}^l(m_\phi) = e^{-in\phi} \delta_{mn}.$$

By the Cartan decomposition each element of $SU(2)$ can be written as $m_\phi b_\theta m_\psi$ and the corresponding integration formula reads

$$(1.14) \quad \int_{SU(2)} f(g) dg = \frac{1}{2} \int_0^\pi \int_0^{4\pi} \int_0^{4\pi} f(m_\phi b_\theta m_\psi) \sin \theta d\theta \frac{d\phi}{4\pi} \frac{d\psi}{4\pi}, \quad f \in C(SU(2)).$$

By Schur's orthogonality relations, (1.13) and (1.14) we obtain

$$\int_0^\pi t_{mn}^l(b_\theta) t_{m', n}^{l'}(b_\theta) \sin \theta d\theta = 0, \quad l \neq l'.$$

Suppose that $m+n \geq 0, m-n \geq 0$. Then the "lowest" element of the orthogonal system $\{t_{mn}^l | l = m, m+1, \dots\}$ is t_{mn}^m . From (1.5) we obtain

$$(1.15) \quad t_{mn}^m(b_\theta) = (-1)^{m-n} \binom{2m}{m-n}^{1/2} \left(\sin \frac{\theta}{2}\right)^{m-n} \left(\cos \frac{\theta}{2}\right)^{m+n}.$$

Hence, if $l \neq l'$, then

$$\int_0^\pi \frac{t_{mn}^l(b_\theta)}{t_{mn}^m(b_\theta)} \frac{t_{mn}^{l'}(b_\theta)}{t_{mn}^m(b_\theta)} \left(\sin \frac{\theta}{2}\right)^{2m-2n+1} \left(\cos \frac{\theta}{2}\right)^{2m+2n+1} d\theta = 0.$$

By (1.5) $t_{mn}^l(b_\theta)/t_{mn}^m(b_\theta)$ is a polynomial in $\cos \theta$ of degree $\leq l-m$. It follows that

$$t_{mn}^l(b_\theta)/t_{mn}^m(b_\theta) = \text{const. } P_{l-m}^{(m-n, m+n)}(\cos \theta),$$

where the *Jacobi polynomial* $P_{l-m}^{(m-n, m+n)}$ is an orthogonal polynomial of degree $l-m$ with respect to the weight function $(1-x)^{m-n}(1+x)^{m+n}$ on the interval $(-1, 1)$. Of course, this result has been derived in many other ways (cf. Vilenkin [11, Chap. 3]).

2. The matrix elements for $SU(2) \times SU(2)$. Let $K := SU(2)$, $G := K \times K$, $K^* := \text{diag}(K \times K)$, $A := \{a_\theta := (m_\theta, m_{-\theta})\}$ (m_θ is defined by (1.12)). Then $G = K^*AK^*$ is a Cartan decomposition. The corresponding integral formula is

$$(2.1) \quad \int_G f(g) dg = \frac{1}{2\pi} \int_0^\pi \int_{K^*} \int_{K^*} f(k_1 a_\theta k_2) \sin^2 \theta d\theta dk_1 dk_2, \quad f \in C(G),$$

which is a special case of Helgason [5, Prop. X.1.19].

A complete system of representatives for \hat{G} is given by the representations $T^{l_1, l_2}(l_1, l_2 \in \frac{1}{2}\mathbb{Z}_+)$:

$$(2.2) \quad T^{l_1, l_2}(k_1, k_2) := T^{l_1}(k_1) \otimes T^{l_2}(k_2), \quad k_1, k_2 \in K.$$

The representation space $H_{l_1} \otimes H_{l_2}$ of T^{l_1, l_2} can be identified with the space of polynomials in four complex variables x, y, u, v , homogeneous of degree $2l_1$ in x, y and homogeneous of degree $2l_2$ in u, v . An orthonormal basis of $H_{l_1} \otimes H_{l_2}$ is given by the polynomials

$$(x, y, u, v) \mapsto \psi_{j_1}^{l_1}(x, y) \psi_{j_2}^{l_2}(u, v).$$

PROPOSITION 2.1 (cf. [6, Thms. 3.1, 3.2]). *The functions $\phi_{l, j}^{l_1, l_2} (|l_1 - l_2| \leq l \leq l_1 + l_2, |j| \leq l)$ defined by*

$$(2.3) \quad \phi_{l, j}^{l_1, l_2}(x, y, u, v) := (-1)^{l_1 + l_2 - l} \left(\frac{(2l+1)(2l_1)!(2l_2)!}{(l_1+l_2-l)!(l_1+l_2+l+1)!} \right)^{1/2} \cdot (xv - yu)^{l_1 + l_2 - l} t_{l_2 - l_1, j}^l \begin{pmatrix} x & y \\ u & v \end{pmatrix}$$

form an orthonormal basis of $H_{l_1} \otimes H_{l_2}$ such that

$$(2.4) \quad T^{l_1, l_2}(k, k) \phi_{l, j}^{l_1, l_2} = \sum_{j' = -l}^l t_{j, j'}^l(k) \phi_{l, j'}^{l_1, l_2}, \quad k \in K.$$

Define the matrix elements of T^{l_1, l_2} with respect to this K^* -basis $\{\phi_{l, j}^{l_1, l_2}\}$ by

$$(2.5) \quad T^{l_1, l_2}(g) \phi_{l, j}^{l_1, l_2} = \sum_{l' = |l_1 - l_2|}^{l_1 + l_2} \sum_{j' = -l'}^l t_{l, j; l', j'}^{l_1, l_2}(g) \phi_{l', j'}^{l_1, l_2}, \quad g \in G.$$

Since the elements of A commute with the elements (m_θ, m_θ) in K^* and since

$$T^{l_1, l_2}(m_\theta, m_\theta) \phi_{l, j}^{l_1, l_2} = e^{-ij\theta} \phi_{l, j}^{l_1, l_2}$$

by (2.4) and (1.12), we conclude that

$$(2.6) \quad t_{l, j; l', j'}^{l_1, l_2}(a_\theta) = 0 \quad \text{if } j \neq j'.$$

By (2.4), (2.6) and the decomposition $G = K^*AK^*$ the matrix elements $t_{l, j; l', j'}^{l_1, l_2}$ will be known if we know the functions $t_{l, j; l', j}^{l_1, l_2}|_A$.

PROPOSITION 2.2 (cf. [3]). *There are the orthogonality relations*

$$(2.7) \quad \frac{1}{2\pi} \sum_{j = -(l \wedge m)}^{l \wedge m} \int_0^\pi t_{l, j; m, j}^{l_1, l_2}(a_\theta) t_{l, j; m, j}^{l_1, l_2}(a_\theta) \sin^2 \theta d\theta = \frac{(2l+1)(2m+1)}{(2l_1+1)(2l_2+1)} \delta_{l_1, l_1'} \delta_{l_2, l_2'}.$$

Proof. It follows from Schur's orthogonality relations, (2.1), (2.4) and (2.6) that

$$\begin{aligned} & \frac{\delta_{l_1, l_1'} \delta_{l_2, l_2'}}{(2l_1+1)(2l_2+1)} \\ &= \frac{1}{2\pi} \int_0^\pi \int_{K^*} \int_{K^*} t_{l_1, p; m, p}^{l_1, l_2}(k_1 a_\theta k_2) \overline{t_{l_1, p; m, p}^{l_1', l_2'}(k_1 a_\theta k_2)} \sin^2 \theta \, d\theta \, dk_1 \, dk_2 \\ &= \sum_{j=-(l \wedge m)}^{l \wedge m} \sum_{j'=- (l \wedge m)}^{l \wedge m} \frac{1}{2\pi} \int_0^\pi \int_{K^*} \int_{K^*} t_{p, j}^l(k_1) t_{l_1, j; m, j}^{l_1, l_2}(a_\theta) t_{j, p}^m(k_2) \\ & \quad \cdot \overline{t_{p, j'}^{l_1}(k_1) t_{l_1, j'; m, j'}^{l_1', l_2'}(a_\theta) t_{j', p}^m(k_2)} \sin^2 \theta \, d\theta \, dk_1 \, dk_2 \\ &= \frac{1}{(2l+1)(2m+1)} \sum_{j=-(l \wedge m)}^{l \wedge m} \frac{1}{2\pi} \int_0^\pi t_{l_1, j; m, j}^{l_1, l_2}(a_\theta) \overline{t_{l_1, j; m, j}^{l_1', l_2'}(a_\theta)} \sin^2 \theta \, d\theta. \end{aligned}$$

It follows from (2.5) and (2.3) that $t_{l_1, j; m, j}^{l_1, l_2}(a_\theta)$ is real. \square




FIG. 1.

From now on fix l and m ($l, m \in \frac{1}{2}\mathbb{Z}_+, l-m \in \mathbb{Z}$) such that $l \leq m$. (Because of unitariness of T^{l_1, l_2} this last condition is not an essential restriction). Then the indices l_1, l_2 in $t_{l_1, j; m, j}^{l_1, l_2}(a_\theta)$ can assume all values in $\frac{1}{2}\mathbb{Z}_+$ such that (cf. Fig. 1)

$$(2.8) \quad l_1 + l_2 \geq m, \quad |l_1 - l_2| \leq l, \quad l_1 + l_2 - l \in \mathbb{Z}$$

and $j \in \{-l, -l+1, \dots, l\}$. Thus, (2.7) can be viewed as the orthogonality relations for the vector-valued functions

$$(2.9) \quad \theta \mapsto \left(t_{l_1, -l; m, -l}^{l_1, l_2}(a_\theta), t_{l_1, -l+1; m, -l+1}^{l_1, l_2}(a_\theta), \dots, t_{l_1, l; m, l}^{l_1, l_2}(a_\theta) \right),$$

where (l_1, l_2) run through all values satisfying (2.8). Like at the end of §1 we pick the "lowest" elements of this orthogonal family. Candidates for these elements are all

functions of the form (2.9) with $l_1 + l_2 = m$. Suppose that we can prove that for all θ in $(0, \pi)$ the matrix

$$(2.10) \quad \left(t_{l,j;m,j}^{(m+p)/2(m-p)/2}(a_\theta) \right)_{j,p=-l,-l+1,\dots,l}$$

is nonsingular. Then, for $n = 0, 1, 2, \dots$ and $k = -l, -l + 1, \dots, l$ we can define the real vector-valued functions

$$(2.11) \quad x \mapsto P_{n,k}^{l,m}(x) = (P_{n,k,-l}^{l,m}(x), P_{n,k,-l+1}^{l,m}(x), \dots, P_{n,k,l}^{l,m}(x))$$

on $(-1, 1)$ by

$$(2.12) \quad t_{l,j;m,j}^{l_1,l_2}(a_\theta) = \sum_{p=-l}^l t_{l,j;m,j}^{(m+p)/2(m-p)/2}(a_\theta) P_{l_1+l_2-m,l_2-l_1,p}^{l,m}(\cos \theta).$$

Also define

$$(2.13) \quad W_{p,q}^{l,m}(\cos \theta) := \sin \theta \sum_{j=-l}^l t_{l,j;m,j}^{(m+p)/2(m-p)/2}(a_\theta) t_{l,j;m,j}^{(m+q)/2(m-q)/2}(a_\theta).$$

Then

$$(2.14) \quad W^{l,m}(\cos \theta) := \left(W_{p,q}^{l,m}(\cos \theta) \right)_{p,q=-l,\dots,l}$$

is a positive definite real symmetric matrix for all θ in $(0, \pi)$ and it follows from (2.7), (2.12), (2.13) that the vector-valued functions $P_{n,k}^{l,m}$ satisfy the orthogonality relations

$$(2.15) \quad \frac{1}{2\pi} \sum_{p,q=-l}^l \int_{-1}^1 P_{n,k,p}^{l,m}(x) P_{n',k',q}^{l,m}(x) W_{p,q}^{l,m}(x) dx = \frac{(2l+1)(2m+1)}{(n+m+1)^2 - k^2} \delta_{n,n'} \delta_{k,k'}.$$

In this paper we will show that the matrix (2.10) is indeed nonsingular for θ in $(0, \pi)$ and that $P_{n,k,p}^{l,m}$ is a polynomial of degree $n - |p + k|$. Hence the orthogonality relations (2.15) will characterize the vector-valued functions $P_{n,k}^{l,m}$ up to constant factors.

3. The vector-valued orthogonal polynomials. First we derive an integral representation for the canonical matrix elements. Consider (2.5) with $g = a_\theta$ and evaluate both sides for $(x, y, u, v) = (\alpha, \beta, -\bar{\beta}, \bar{\alpha})$, where $|\alpha|^2 + |\beta|^2 = 1$. In view of (2.3) and (2.6) we obtain

$$\begin{aligned} & (-1)^{l_1+l_2-m} \left(\frac{(2m+1)(2l_1)!(2l_2)!}{(l_1+l_2-m)!(l_1+l_2+m+1)!} \right)^{1/2} \\ & \cdot t_{l_2-l_1,j}^m \begin{pmatrix} e^{i\theta/2}\alpha & e^{-i\theta/2}\beta \\ -e^{-i\theta/2}\bar{\beta} & e^{i\theta/2}\bar{\alpha} \end{pmatrix} (e^{i\theta}|\alpha|^2 + e^{-i\theta}|\beta|^2)^{l_1+l_2-m} \\ & = \sum_{l=|l_1-l_2|}^{l_1+l_2} (-1)^{l_1+l_2-l} \left(\frac{(2l+1)(2l_1)!(2l_2)!}{(l_1+l_2-l)!(l_1+l_2+l+1)!} \right)^{1/2} \\ & \cdot t_{l,j;m,j}^{l_1,l_2}(a_\theta) t_{l_2-l_1,j}^l(k(\alpha, \beta)). \end{aligned}$$

Hence, by Schur's orthogonality relations:

$$(3.1) \quad t_{l_1, j; m, j}^{l_1, l_2}(a_\theta) = (-1)^{l-m} \left(\frac{(2l+1)(2m+1)(l_1+l_2-l)!(l_1+l_2+l+1)!}{(l_1+l_2-m)!(l_1+l_2+m+1)!} \right)^{1/2} \\ \cdot \int_K (e^{i\theta}|\alpha|^2 + e^{-i\theta}|\beta|^2)^{l_1+l_2-m} \\ \cdot t_{l_2-l_1, j}^m \left(\begin{matrix} e^{i\theta/2}\alpha & e^{-i\theta/2}\beta \\ -e^{-i\theta/2}\bar{\beta} & e^{i\theta/2}\bar{\alpha} \end{matrix} \right) t_{l_2-l_1, j}^l(k(\bar{\alpha}, \bar{\beta})) dk(\alpha, \beta).$$

Next, by some manipulations we will modify this integral representation into a form which is more suitable for our purpose. Substitution of (1.7) into (3.1) yields

$$t_{l_1, j; m, j}^{l_1, l_2}(a_\theta) = c_{l_1, j; m, j}^{l_1, l_2} \frac{1}{2\pi} \int_K \int_0^{2\pi} (e^{i\theta}|\alpha|^2 + e^{-i\theta}|\beta|^2)^{l_1+l_2-m} \\ \cdot (\alpha e^{i(\phi+\theta/2)} + \beta e^{-i(\phi+\theta/2)})^{m+l_1-l_2} \\ \cdot (-\bar{\beta} e^{i(\phi-\theta/2)} + \bar{\alpha} e^{-i(\phi-\theta/2)})^{m-l_1+l_2} \\ \cdot e^{2ij\phi} t_{l_2-l_1, j}^l(k(\bar{\alpha}, \bar{\beta})) dk(\alpha, \beta) d\phi,$$

where

$$(3.2) \quad c_{l_1, j; m, j}^{l_1, l_2} = (-1)^{l-m} \left(\frac{(2l+1)(2m+1)(l_1+l_2-l)!(l_1+l_2+l+1)!(m-j)!(m+j)!}{(l_1+l_2-m)!(l_1+l_2+m+1)!(m+l_1-l_2)!(m-l_1+l_2)!} \right)^{1/2}.$$

In this last integral representation consider the K -integral as the inner integral and make the transformation of integration variable $k(\alpha, \beta) \mapsto k(\bar{\alpha}, \bar{\beta})m_{-2\phi}$. Then the integrand no longer depends on ϕ and we obtain

$$(3.3) \quad t_{l_1, j; m, j}^{l_1, l_2}(a_\theta) = c_{l_1, j; m, j}^{l_1, l_2} \int_K (e^{i\theta}|\alpha|^2 + e^{-i\theta}|\beta|^2)^{l_1+l_2-m} \\ \cdot (\alpha e^{i\theta/2} - \beta e^{-i\theta/2})^{m-l_1+l_2} (\bar{\alpha} e^{i\theta/2} + \bar{\beta} e^{-i\theta/2})^{m+l_1-l_2} \\ \cdot t_{l_2-l_1, j}^l(k(\alpha, \beta)) dk(\alpha, \beta).$$

LEMMA 3.1. *Let K be a connected compact Lie group which has a complexification K_c . Let f be a complex analytic function on an open connected left- K -invariant subset V of K_c containing K . Then*

$$(3.4) \quad \int_K f(k) dk = \int_K f(kk') dk, \quad k' \in V.$$

Proof. The right-hand side is a complex analytic function of k' on V which is constant on K . \square

Now observe that the integrand in (3.3) is the restriction to $SU(2)$ of the complex analytic function

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \mapsto (e^{i\theta}\alpha\delta - e^{-i\theta}\beta\gamma)^{l_1+l_2-m} (\alpha e^{i\theta/2} - \beta e^{-i\theta/2})^{m-l_1+l_2} \\ \cdot (-\gamma e^{-i\theta/2} + \delta e^{i\theta/2})^{m+l_1-l_2} t_{l_1-l_2, j}^l \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \text{ on } SL(2, \mathbb{C}).$$

For $0 < \theta < \pi$ apply Lemma 3.1 to this function with K' chosen as

$$(3.5) \quad g_\theta := e^{i\pi/4} (2 \sin \theta)^{-1/2} \begin{pmatrix} e^{-i\theta/2} & e^{i\theta/2} \\ e^{i\theta/2} & e^{-i\theta/2} \end{pmatrix}.$$

We obtain

$$(3.6) \quad t_{l, j; m, j}^{l_1, l_2}(a_\theta) = c_{l, j; m, j}^{l_1, l_2} e^{3\pi i m/2} (2 \sin \theta)^m \\ \cdot \sum_{p=-l}^l t_{p, j}^l(g_\theta) \int_K (2|\beta|^2 \cos \theta + \alpha \bar{\beta} - \bar{\alpha} \beta)^{l_1+l_2-m} \\ \cdot \beta^{m-l_1+l_2} (-\bar{\beta})^{m+l_1-l_2} t_{l_2-l_1, p}^l(k(\alpha, \beta)) dk(\alpha, \beta).$$

PROPOSITION 3.2. *We have*

$$(3.7) \quad t_{l, j; m, j}^{(m+p)/2, (m-p)/2}(a_\theta) = \left(\frac{(2l+1)(m-j)!(m+j)!(m-p)!(m+p)!}{(2m)!(m-l)!(m+l+1)!} \right)^{1/2} \\ \cdot (-1)^{l+m} e^{3\pi i m/2} (2 \sin \theta)^m t_{p, j}^l(g_\theta).$$

For $0 < \theta < \pi$ the matrix $(t_{l, j; m, j}^{(m+p)/2, (m-p)/2}(a_\theta))_{j, p=-l, \dots, l}$ is nonsingular.

Proof. Formula (3.6), together with (1.13) and the invariance of the integral in (3.6) under right multiplication by m_ϕ yields

$$t_{l, j; m, j}^{(m+p)/2, (m-p)/2}(a_\theta) = c_{l, j; m, j}^{(m+p)/2, (m-p)/2} e^{3\pi i m/2} (2 \sin \theta)^m \\ \cdot t_{p, j}^l(g_\theta) \int_K \beta^{m-p} (-\bar{\beta})^{m+p} t_{-p, p}^l(k(\alpha, \beta)) dk(\alpha, \beta).$$

The integral can be evaluated by using (1.5), (1.14), the beta integral and the *Chu–Vandermonde sum*

$$(3.8) \quad {}_2F_1(-n, b; c; 1) = \frac{(c-b)_n}{(c)_n}, \quad n=0, 1, \dots; \quad c-b, c \neq 0, -1, \dots, -n+1.$$

Finally use (3.2). \square

THEOREM 3.3. *Formula (2.12) holds with*

$$(3.9) \quad P_{n, k, p}^{l, m}(x) = A_{n, k, p}^{l, m} \int_K (2|\beta|^2 x + \alpha \bar{\beta} - \bar{\alpha} \beta)^n \beta^{m+k} (-\bar{\beta})^{m-k} t_{k, p}^l(k(\alpha, \beta)) dk(\alpha, \beta),$$

where

(3.10)

$$A_{n,k,p}^{l,m} := (-1)^{2l} \left(\frac{(2m+1)!(n+m-l)!(n+m+l+1)!(m-l)!(m+l+1)!}{n!(n+2m+1)!(m-k)!(m+k)!(m-p)!(m+p)!} \right)^{1/2}.$$

There are the symmetries

$$(3.11) \quad P_{n,k,p}^{l,m} = P_{n,p,k}^{l,m} = P_{n,-k,-p}^{l,m} = P_{n,-p,-k}^{l,m},$$

$$(3.12) \quad P_{n,k,p}^{l,m}(-x) = (-1)^{n+k+p} P_{n,k,p}^{l,m}(x).$$

Proof. Formula (3.9) follows from (3.7), (3.6) and (3.2). The symmetries are derived from (3.9) by the use of (1.6) and (1.9) in the case of (3.11) and by (1.13) in the case of (3.12). \square

Of course, by the use of (2.12) and (3.7), the symmetries (3.11) imply certain symmetries for the matrix elements $t_{l,j;m,j}^{l_1,l_2} |_{\mathcal{A}}$. It would be interesting to get a deeper understanding of the first of these symmetries.

Now expand the integrand in (3.9) with respect to x and use the invariance of the integral under right multiplication with m_ϕ and (1.13). We obtain

$$(3.13) \quad P_{n,k,p}^{l,m}(x) = A_{n,k,p}^{l,m} \sum_{\substack{q=|p+k| \\ q+k+k \text{ even}}}^n d_{n,k,p,q}^{l,m} x^{n-q},$$

where

$$(3.14) \quad d_{n,k,p,q}^{l,m} = \frac{(-1)^{m-k+(q-k-p)/2} 2^{n-q} n!}{((q-k-p)/2)!((q+k+p)/2)!(n-q)!} \\ \cdot \int_K \alpha^{(q+k+p)/2} \bar{\alpha}^{(q-k-p)/2} \beta^{m+n+(k-p-q)/2} \\ \cdot \bar{\beta}^{m+n+(-k+p-q)/2} t_{kp}^l(k(\alpha, \beta)) dk(\alpha, \beta).$$

By using (1.5), (1.14) and the beta integral we obtain, for $k+p \geq 0$

$$(3.15) \quad d_{n,k,p,q}^{l,m} = d_{n,-k,-p,q}^{l,m} = \frac{(-1)^{l+m+(q+k+p)/2} 2^{n-q} n! (l+m+n-(q+k+p)/2)!}{((q-k-p)/2)!(n-q)!(k+p)!(l+m+n+1)!} \\ \cdot \sqrt{\frac{(l+k)!(l+p)}{(l-k)!(l-p)}} {}_3F_2 \left(\begin{matrix} -l+k, -l+p, (q+k+p)/2+1 \\ k+p+1, -l-m-n+(q+k+p)/2 \end{matrix} \middle| 1 \right).$$

For $q=p+k$ use (3.8). Then, for $k+p \geq 0$

(3.16)

$$d_{n,k,p,k+p}^{l,m} = d_{n,-k,-p,k+p}^{l,m} = \frac{(-1)^{l+m+p+k} 2^{n-p-k} n! (m+n-k)!(m+n-p)!}{(m-l+n)!(m+l+n+1)!(p+k)!(n-p-k)!} \\ \cdot \left(\frac{(l+k)!(l+p)!}{(l-k)!(l-p)!} \right)^{1/2} \neq 0.$$

Hence $P_{n,k,p}^{l,m}$ is a polynomial of degree $n - |p + k|$.

THEOREM 3.4. *The vector-valued polynomial $P_{n,k}^{l,m}$ satisfies the conditions*

$$(3.17) \quad P_{n,k,p}^{l,m}(x) = \frac{(-1)^{l-m} 2^n (m-k+1)_n (m+k+1)_n \delta_{k,-p} x^n}{(n!(2m+2)_n (m-l+1)_n (m+l+2)_n)^{1/2}}$$

+ polynomial of degree less than n ,

$$(3.18) \quad \sum_{p=-l}^l \int_{-1}^1 P_{n,k,p}^{l,m}(x) x^{n'} W_{p,q}^{l,m}(x) dx = 0$$

for all q in $\{-l, \dots, l\}$ and all n' in $\{0, \dots, n-1\}$.

Proof. Use (3.13), (3.16) and (3.10) for (3.17), and (2.15) together with (3.17) for (3.18). \square

Note that (3.17) and (3.18) completely determine $P_{n,k}^{l,m}$. They also imply (2.15) for $n \neq n'$. However, from the point of view of Theorem 3.4, the orthogonality relations (2.15) for $n = n', k \neq k'$ are rather unexpected.

Remark 3.5. Lemma 3.1 can also be applied in order to extract the factor $t_{mn}^m(b_\theta)$ from the integral representation (1.8) for $t_{mn}^l(b_\theta)$. Substitute $\alpha := \cos(\theta/2)$, $\beta := \sin(\theta/2)$ in (1.8) and make the successive transformations of integration variable $\phi \mapsto z \mapsto \psi \mapsto \chi$, where $e^{2i\phi} = z = e^{i\psi} \cot(\theta/2)$, $\chi = 2\psi$:

$$\begin{aligned} & \left(\frac{(l-m)!(l+m)!}{(l-n)!(l+n)!} \right)^{1/2} t_{mn}^l(b_\theta) \\ &= \frac{1}{2\pi i} \oint_{(0)} (z \cos(\theta/2) + \sin(\theta/2))^{l-m} (-z \sin(\theta/2)) + \cos(\theta/2)^{l+m} z^{n-l-1} dz \\ &= (\sin(\theta/2))^{m-n} (\cos(\theta/2))^{m+n} \\ & \quad \cdot \frac{1}{2\pi} \int_0^{2\pi} \left(e^{i\psi} \cos^2 \frac{\theta}{2} + \sin^2 \frac{\theta}{2} \right)^{l-m} e^{i\psi(n-l)} (1 - e^{i\psi})^{l+m} d\psi \\ &= (-2i)^{l+m} \left(\sin \frac{\theta}{2} \right)^{m-n} \left(\cos \frac{\theta}{2} \right)^{m+n} \\ & \quad \cdot \frac{1}{\pi} \int_0^\pi (\cos \chi + i \sin \chi \cos \theta)^{l-m} e^{2ni\chi} (\sin \chi)^{l+m} d\chi. \end{aligned}$$

Now assume $m \geq n$ and use [2, 1.5(29)]. Then

$$(3.19) \quad t_{mn}^l(b_\theta) / t_{mn}^m(b_\theta) = \text{const.} \int_0^\pi (\cos \chi + i \sin \chi \cos \theta)^{l-m} e^{2ni\chi} (\sin \chi)^{l+m} d\chi$$

with nonzero constant. Again by [2, 1.5 (29)], the right-hand side of (3.19) is a polynomial of degree $l-m$ in $\cos \theta$ which takes a nonzero value if $\cos \theta = 1$. In Greiner and Koornwinder [4, §1.3] the integral representation for Jacobi polynomials resulting from (3.19) is obtained in a quite different context.

4. The noncompact analogue. Let now $G := SL(2, C)$ with Iwasawa decomposition $G = KAN$ such that

$$K = SU(2), \quad A = \left\{ a_t := \begin{pmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{pmatrix} \middle| t \in \mathbb{R} \right\}, \quad N := \left\{ \begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix} \middle| z \in \mathbb{C} \right\}.$$

Let $k(\alpha, \beta)$ in K be defined by (1.10) and m_ϕ by (1.12). $M := \{m_\phi | 0 \leq \phi < 4\pi\}$ is the centralizer of A in K .

Let $\pi^{\lambda, k}(\lambda \in \mathbb{C}, k \in \frac{1}{2}\mathbb{Z})$ be the representation of G which is induced by the representation $m_\phi a_t n \mapsto e^{-ik\phi} e^{\lambda t}$ of MAN : a principal series representation. Then $\pi^{\lambda, k}|_K$ is unitary and decomposes as $\bigoplus_{l=k, k+1, \dots} T^l$. Choose a K -basis for which $\pi^{\lambda, k}$ has matrix elements $\pi_{l, p; m, q}^{\lambda, k}(l, m = k, k+1, \dots; p = -l, \dots, l; q = -m, \dots, m)$ such that

$$\pi_{l, p; m, q}^{\lambda, k}(k) = \delta_{l, m} t_{p, q}^l(k), \quad k \in K.$$

Then

$$(4.1) \quad \pi_{l, j; m, j}^{\lambda, k}(a_t) = (2l+1)^{1/2} \int_K \left(e^{-t} |\alpha|^2 + e^t |\beta|^2 \right)^{-\lambda-m-1} \\ \cdot t_{kj}^m \begin{pmatrix} e^{-t/2} \alpha & e^{t/2} \beta \\ -e^{t/2} \bar{\beta} & e^{-t/2} \bar{\alpha} \end{pmatrix} t_{kj}^l(k(\bar{\alpha}, \bar{\beta})) dk(\alpha, \beta),$$

cf. Rühl [9, §3–5], Kosters [8, §3.1].

Similarly to (3.3) we derive from (4.1) that

$$(4.2) \quad \pi_{l, j; m, j}^{\lambda, k}(a_t) = c_{k, l, m, j} \int_K \left(e^{-t} |\alpha|^2 + e^t |\beta|^2 \right)^{-\lambda-m-1} (e^{-t/2} \alpha - e^{t/2} \beta)^{m+k} \\ \cdot (e^{-t/2} \bar{\alpha} + e^{t/2} \bar{\beta})^{m-k} t_{kj}^l(k(\alpha, \beta)) dk(\alpha, \beta),$$

where

$$(4.3) \quad c_{k, l, m, j} := \left(\frac{(2l+1)(2m+1)(m-j)!(m+j)!}{(m-k)!(m+k)!} \right)^{1/2}.$$

For $s > 0$ let

$$(4.4) \quad h_s := (2 \operatorname{sh} s)^{-1/2} \begin{pmatrix} e^{s/2} & e^{-s/2} \\ e^{-s/2} & e^{s/2} \end{pmatrix}.$$

Then we can apply Lemma 3.1 to (4.2) with $k' := h_s$ for $0 < t < s$. We obtain

$$(4.5) \quad \pi_{l, j; m, j}^{\lambda, k}(a_t) = c_{k, l, m, j} 2^m (\operatorname{sh} s)^{-m} \\ \cdot \sum_{p=-l}^l t_{pj}^l(h_s) \int_K \left(\operatorname{ch} t - \operatorname{coth} s \operatorname{sh} t (|\alpha|^2 - |\beta|^2) + (\alpha \bar{\beta} - \beta \bar{\alpha}) \frac{\operatorname{sh} t}{\operatorname{sh} s} \right)^{-\lambda-m-1} \\ \cdot (\alpha \operatorname{sh} \frac{1}{2}(s-t) - \beta \operatorname{sh} \frac{1}{2}(s+t))^{m+k} \\ \cdot (\bar{\alpha} \operatorname{sh} \frac{1}{2}(s-t) + \bar{\beta} \operatorname{sh} \frac{1}{2}(s+t))^{m-k} \\ \cdot t_{kp}^l(k(\alpha, \beta)) dk(\alpha, \beta), \quad 0 < t < s.$$

If $\operatorname{Re} \lambda \leq m - 1$ then the limit passage $s \downarrow t$ is certainly allowed in (4.5).

(4.6)

$$\begin{aligned} \pi_{l,j;m,j}^{\lambda,k}(a_t) &= c_{k,l,m,j} (-1)^{2m} (2 \operatorname{sh} t)^m \\ &\cdot \sum_{p=-l}^l t_{pj}^l(h_t) \int_K (2|\beta|^2 \operatorname{ch} t + \alpha \bar{\beta} - \beta \bar{\alpha})^{-\lambda-m-1} \\ &\cdot \beta^{m+k} (-\bar{\beta})^{m-k} t_{kp}^l(k(\alpha, \beta)) dk(\alpha, \beta). \end{aligned}$$

Closer examination of the integral, using (1.14), shows that (4.6) holds with convergent integral if $\operatorname{Re} \lambda < 0$. Thus it is meaningful to study the vector-valued function $x \mapsto (P_{n,k,p}^{l,m}(x))_{p=-l,\dots,l}$, defined by (3.9), for complex n , $\operatorname{Re} n > 0$, and for $x > 1$. In particular, this function has a nice asymptotics as $x \rightarrow \infty$.

REFERENCES

- [1] D. BASU AND S. SRINIVASAN, *A unified treatment of the groups $SO(4)$ and $SO(3,1)$* , Czech. J. Phys., B 27 (1977), pp. 629–635.
- [2] A. ERDELYI, et al., *Higher Transcendental Functions, Vol. I*, McGraw-Hill, New York, 1953.
- [3] D. Z. FREEDMAN AND J.-M. WANG, *$O(4)$ symmetry and Regge-pole theory*, Phys. Rev., 160 (1967), pp. 1560–1571.
- [4] P. C. GREINER AND T. H. KOORNWINDER, *Variations on the Heisenberg spherical harmonics*, Math. Centrum Report, ZW 186, Mathematisch Centrum, Amsterdam, 1983.
- [5] S. HELGASON, *Differential Geometry and Symmetric Spaces*, Academic Press, New York, 1962.
- [6] T. H. KOORNWINDER, *Clebsch–Gordan coefficients for $SU(2)$ and Hahn polynomials*, Nieuw Arch. Wisk., (3) 29 (1981), pp. 140–155.
- [7] ———, *The representation theory of $SL(2, \mathbb{R})$, a noninfinitesimal approach*, Enseignement Math., 28 (1982), pp. 53–90.
- [8] M. T. KOSTERS, *A study of the representations of $SL(2, \mathbb{C})$ using noninfinitesimal methods*, Math. Centrum Report TW 190, Mathematisch Centrum, Amsterdam, 1979.
- [9] W. RÜHL, *The Lorentz Group and Harmonic Analysis*, Benjamin, Menlo Park, CA, 1970.
- [10] YA. A. SMORODINSKIĬ AND G. I. SHEPELEV, *Boost matrix elements in $O(3,1)$ and continuation to $O(4)$* , Soviet J. Nuclear Phys., 13 (1971), pp. 248–253.
- [11] N. J. VILENKIN, *Special Functions and the Theory of Group Representations*, AMS Transl. Math. Monographs, 22, American Mathematical Society, Providence, RI, 1968.