

Space Cold Atom Clock with Counter-Propagating Atoms

Desheng Lü^{1,2}, Bin Wang^{1,2}, Tang Li¹, and Liang Liu^{1,3,*}

¹Key Laboratory of Quantum Optics,

Shanghai Institute of Optics and Fine Mechanics,

Chinese Academy of Sciences, Shanghai 201800, China

²Graduate University of the Chinese Academy of Sciences,

Beijing 100039, China

³State Key Laboratory of Precision Spectroscopy,

East China Normal University, Shanghai 200062, China

*Corresponding author: liang.liu@siom.ac.cn

We discuss the feasibility to realize a space cold atom clock with counter-propagating cold atoms in microgravity. The design of the space clock is based on atomic beam clock with a Ramsey cavity, except a magneto-optical trap (MOT) is placed at each side. Cold atoms are launched from MOTs at both side of the clock simultaneously and move at counter-direction towards each other. The velocity of launched atoms is precisely controlled to the Ramsauer-Townsend resonance so that no additional collision frequency shift is taken place. Such a configuration can efficiently cancel the frequency shift led from cavity phase shift and increase the signal to noise.

In a classical thermal cesium clock, a beam of atoms effuses from an oven and passes through a state-selecting magnet, and subsequently passes through a Ramsey microwave cavity, and then detected. The velocity of thermal atoms can be as slow as 95 m/s and line width of the clock transition is typically around 60 Hz, as PTB Cs frequency standards apply [1]. In order to test the cavity phase frequency shift, in PTB's primary clock CS2, an oven and detector is placed at each end so that alternate operation of atomic beam in opposite directions can be performed [2].

In cold atom space clock, the PHARAO [3] for example, cold atoms are launched from a MOT at velocity as low as 5 cm/s, and 0.1 Hz width of central Ramsey fringes is predicted. The expected stability is $10^{-13}/\sqrt{\tau}$, where τ is the integration time, and the accuracy is up to 10^{-16} . For such a high stability and accuracy, the phase shift of the space clock's Ramsey cavity becomes more important. Certainly, it is possible to apply PTB's CS2 clock design in the space clock, and alternate operation of cold atoms in opposite direction gives the information of the cavity phase shift, but such a design wastes precious space resources.

In this paper, we propose a new type of cold atom space clock, whose design is similar to the PTB's CS2 [2], but with completely new operation mode. The new type of space clock aims at cancelling the frequency shift due to the phase difference of the Ramsey cavity and increasing the signal to noise ratio, and thus reducing the technical difficulties of the design and improving the performance of the space clock. As shown in Fig. 1, a MOT and a detection region is placed at each end. Cold atoms are launched from both MOTs simultaneously in opposite direction towards each other. Assuming Cloud A denotes the cold atoms launched from the left MOT, and Cloud B from the right MOT, Cloud A collides with Cloud B at the center of the Ramsey cavity after they pass through the first interaction region of the cavity.




FIG. 1: Schematic diagram of cold atom space clock with counter-propagating atoms

In atom-atom collisions, if the atoms are treated classically as hard balls, the calculated cross section is independent of the atomic energy. In quantum mechanics, however, the atoms are considered to present a dipole-dipole interaction of typical atomic dimensions for the scattering between atoms. The solution of Schrödinger Equation for two dipole-dipole potentials shows that the cross-section in atom-atom collisions will have a minimum at some specific energies. This is a simple illustration of the Ramsauer-Townsend resonance [4]. At the Ramsauer-Townsend resonance, the atoms are transparent with each other when they collide, and frequency shift lead from collisions becomes null. Gibble *et.al.* measured the *s*-wave frequency shift with juggling ^{133}Cs and ^{87}Rb fountain clock [5–7]. The first Ramsauer-Townsend shift null for ^{87}Rb happens at alternate launch time delays of $\Delta t = 22$ ms between two cold atom balls, corresponding to the velocity $v_{RT} = g\Delta t/2 = 10.78$ cm/s of each ball propagating at opposite direction. At such a velocity, the collision between the Cloud A and B does not contribute additional frequency shift. For the scattering of identical particles, the *p*-wave scattering does not contribute additional frequency shift.

We assume that the interrogation length of the Ramsey cavity is $L = 52.5$ cm, which gives the interrogation

time $T_{RT} = L/v_{RT} = 4.9$ s when Cloud A or Cloud B moves at the Ramsauer-Townsend velocity v_{RT} . Such an interrogation time corresponds to the linewidth of central Ramsey fringe at $\Delta\nu = 0.1$ Hz. If the number of detected atoms is $N = 10^6$, we have the Allan variance

$$\sigma_y(\tau) = \frac{\Delta\nu}{\pi\nu_0\sqrt{N}}\sqrt{\frac{T}{\tau}} = 1.5 \times 10^{-14}/\sqrt{\tau} \quad (1)$$

where, $T = 10$ s which includes time for the preparation of cold atoms, state selection, interrogation time and detection, and $\nu_0 = 6.835$ GHz is the frequency of clock transition of ^{87}Rb , and τ is the integration time. If we do not consider the phase shift of the Ramsey cavity, we can average the detected atoms from both Cloud A and B such that $N = 2 \times 10^6$, which leads to the Allan variance to be reduced by a factor of $\sqrt{2}$ from Eq. (1), that is $\sigma_y(\tau) = 1.1 \times 10^{-14}/\sqrt{\tau}$.

On the other hand, with atoms' counter-propagating through a Ramsey cavity, the cavity phase shift can be accurately measured, and thus the frequency shift can be adjusted away from the error budgets of the clock. Or the frequency shift due to phase differences of the cavity can be cancelled if we average the signals from both Cloud A and Cloud B. Assuming a phase difference $\Delta\varphi$ between two zones of the Ramsey cavity, we can easily have the probability to find the two-level system in the excited state as [8]

$$p_A = \frac{1}{2} \sin^2 \Omega t \{1 + \cos[2\pi(\nu - \nu_0)T_{RT} + \Delta\varphi]\} \quad (2)$$

$$p_B = \frac{1}{2} \sin^2 \Omega t \{1 + \cos[2\pi(\nu - \nu_0)T_{RT} - \Delta\varphi]\} \quad (3)$$

Here, p_A and p_B are the probability for Cloud A and Cloud B, respectively. Ω is the Rabi frequency, t is the interaction time between atoms and microwave. Generally, the phase difference $\Delta\varphi$ shifts the center of Ramsey fringe by

$$\frac{\Delta\nu_\varphi}{\nu_0} = -\frac{\Delta\varphi}{2\pi\nu_0 T_{RT}} \quad (4)$$

in p_A and $-\Delta\nu_\varphi/\nu_0$ in p_B . Typically, the phase difference of a U-type Ramsey cavity can be controlled below a few hundred μrad . If we take the phase difference $\Delta\varphi = 500$ μrad for example, we have $\Delta\nu_\varphi/\nu_0 = 2.3 \times 10^{-15}$. Thus in order to get accuracy of a few 10^{-16} , the frequency shift due to the phase difference of the cavity must be carefully considered.

If we take an average over the probability of Cloud A and Cloud B, we have

$$\begin{aligned} p &= \frac{p_A + p_B}{2} \\ &= \frac{1}{2} \sin^2 \Omega t [1 + \cos 2\pi(\nu - \nu_0)T_{RT} \cdot \cos \Delta\varphi] \end{aligned} \quad (5)$$

Obviously, in p , the phase difference of the cavity does not contribute any frequency shift, but the width of the

central Ramsey fringe has been broadened to

$$\Delta\nu_p = \Delta\nu + 2\Delta\nu_\varphi \quad (6)$$

Typically phase difference of the Ramsey cavity is around a few hundred μrad [1]. From Eq. (4), we have $\Delta\nu_\varphi \approx 1.6 \times 10^{-5}$ Hz when $\Delta\varphi = 500$ μrad for example, which can be neglected in Eq. (6), and thus the width broadening due to the average of the signal from both counter-propagating atoms can be neglected in the Allan variance given in Eq. (1). Since the collision between Cloud A and Cloud B, each moves at the Ramsauer-Townsend velocity, does not contribute additional frequency shift, the frequency shift due to the cavity phase difference in our system can be cancelled without additional cost.

In addition, our design has a very important function for reducing the noise of the interrogation time due to the vibrations on the space craft and variations of residual microgravity in the atoms' propagating direction. This feature was first realized by Fertig *et al.* in the design of a microgravity atomic clock with the double Ramsey cavity [9]. Fertig's design has two Ramsey cavities located at both sides of a MOT, and cold atoms are launched alternatively to each Ramsey cavity. Since the alternatively launching of cold atoms in the opposite direction does not happen at the same time, the cold atoms do not sense the same vibration and residual microgravity. In our design, however, the counter-propagating atoms through same cavity at the same time, the detected signals of Cloud A and Cloud B behave oppositely and simultaneously and thus the effect of vibrations and variations of the microgravity can be removed by the averaged signal p .

Our design has some unique features especially suitable for some experiments. For example, if the velocity of launched atoms varies, the cross-section of atom-atom scattering can be measured by our space clock even more precisely than by the juggling fountain clock. Such a measurement can give detailed data for atom-atom scattering and test the fundamental principle of quantum mechanics.

In conclusions, we have proposed a new type of cold atom space clock with counter-propagating atoms. The cold atoms move at Ramsauer-Townsend velocity so that the collision between counter-propagating atoms becomes null. We pointed out that such a null collision cross section can cancel the frequency shift lead from the phase difference of the Ramsey cavity, and also increase the signal to noise ratio. We estimated the Allan variance of such a space clock at $1.5 \times 10^{-14}/\sqrt{\tau}$. Besides, our design of the cold atom space clock can efficiently remove the noise due to the vibration and residual microgravity of the space craft, and thus reduce the requirement of space environment. Such a kind of space clock will give a new way to test the fundamental physics.

This work is supported by the National High-Tech Programme under Grant No. 2006AA12Z311 and National Basic Research Programme of China under Grant No. 2005CB724506.

-
- [1] A. Bauch, "The PTB primary clocks CS1 and CS2," *Metrologia* **42**, S43 (2005)
- [2] A. Bauch, "Caesium atomic clocks: functions, performance and applications," *Meas. Sci. Technol.* **14**, 1159 (2003)
- [3] Ph. Laurent, M. Gbgrall, Ch. Jentsch, P. lemonde, G. Santarelli, A. Clairon, I. Maksimovic, S. Bize, Ch. Salomon, D. Blonde, J.F. Vega, O. Grosjean, F. Picard, M. Saccoccio, M. Chaubet, N. Ladiette, L. Guillet, I. Zenone, Ch. Delaroche, and Ch. Sirmain, "Design of the cold atom PHARAO space clock and initial test results," *Appl. Phys. B* **84**, 683 (2006)
- [4] N.F. Mott and H.S.W. Massey, *The Theory of Atomic Collisions*, (Oxford University Press, London, 1965), Chap. 18
- [5] R. Legere and K. Gibble, "Quantum scattering in a juggling atomic fountain," *Phys. Rev. Lett.* **81**, 5780 (1998)
- [6] C. Fertig, J. Bouttier, and K. Gibble, "Laser-cooled Rb clock," in *Proceedings of the 2000 IEEE/EIA International Frequency Control Symposium and Exhibition*, pp.680-686
- [7] C. Fertig, J.I. Rees, and K. Gibble, "A juggling Rb fountain clock and a direct measurement of population differences," in *Proceedings of the 2001 IEEE International Frequency Control Symposium and PDA Exhibition*, pp.18-21
- [8] F. Riehle, *Frequency Standards*, (Wiley-VCH Verlag, 2004), p.193
- [9] C. Fertig and K. Gibble, B. Klipstein, L. Maleki, D. Seidel, R. Thompson, "RACE: Laser-cooled Rb microgravity clock," in *Proceedings of the 2000 IEEE/EIA International Frequency Control Symposium and Exhibition*, pp.676-679