

Proof that Nonlinear Plane Waves cannot be destabilized
by Scalar Diffusion

John Pearson
Los Alamos Publication: LAUR 93-1758
Center for Nonlinear Studies, MS B258
Los Alamos National Laboratory
Los Alamos, NM 87545
505-667-9958
pearson@infidel.lanl.gov

We begin with the reaction-diffusion equation:

$$\frac{\partial u}{\partial t} = \mathbf{D} \frac{\partial^2}{\partial x^2} u + \mathbf{D} \nabla_{\perp}^2 u + F(u) \quad 1$$

where u is a *vector* of chemical concentrations and \mathbf{D} is the diffusion matrix. Here ∇_{\perp}^2 refers to directions orthogonal to x . Assume that there exists a plane wave solution to Eq. (1). Denote this solution by $U(x-ct)$. Define $\xi = x - ct$. Then we have

$$-cU_{\xi} = \mathbf{D}U_{\xi\xi} + F(U) \quad 2$$

Now we ask under what conditions is the plane wave solution stable in multiple dimensions. To do this we work in the travelling coordinate system and write:

$$u(\xi, y, t) = U(\xi) + \eta(\xi, y, t) \quad 3$$

where η is a small perturbation to the plane wave solution. We expand η in Fourier modes in y . (The dimensionality of the system is irrelevant. For simplicity I work only in 2 space dimensions here, ξ and y .) Thus we have:

$$\eta = \sum_k a_k(\xi) \phi_k(y) \exp(\lambda_k t) \quad 4$$

where the ϕ_k are eigenfunctions of ∇_{\perp}^2 . Inserting equations (3) and (4) into (1) we find the eigenvalue problem for λ_k :

$$\lambda_k a_k(\xi) = \mathbf{D} \frac{\partial^2 a_k}{\partial \xi^2} - \mathbf{D} k^2 a_k(\xi) + J(\xi) a_k(\xi) + c \frac{\partial a_k}{\partial \xi} \quad 5$$

In the case that $\mathbf{D} = D\mathbf{I}$ we have

$$\lambda_k = \lambda_0 - Dk^2 \quad 6$$

Equation (6) means that diffusion stabilizes small perturbations of the wave front in the case of scalar diffusion which is the desired result.



LOS ALAMOS NAT'L LAB LIBS



3 9338 00205 3998

LOS ALAMOS NAT'L LAB.
IS-4 REPORT SECTION
RECEIVED

'93 OCT 19 AM 9 41