

# **LEGIBILITY NOTICE**

**A major purpose of the Technical Information Center is to provide the broadest dissemination possible of information contained in DOE's Research and Development Reports to business, industry, the academic community, and federal, state and local governments.**

**Although a small portion of this report is not reproducible, it is being made available to expedite the availability of information on the research discussed herein.**

LA-UR--89-860

DE89 009391

APR 05 1989

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405 ENG 36

TITLE Nuclear Fission - An Inherently Non-Equilibrium Process?

AUTHOR(S) Victor (Avigdor) Gavron

MASTER

SUBMITTED TO Talk to be presented at "50 Years of Nuclear Fission" Conference, Berlin, April 3-7, 1989, and subsequently to be published in Nuclear Physics A.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the substance form of this contribution or to allow others to do so for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

Los Alamos Los Alamos National Laboratory Los Alamos, New Mexico 87545

## Heavy Ion Fission - An Inherently Non-Equilibrium Process?

A. Gavron

Los Alamos National Laboratory, Los Alamos, NM87545

Recent measurements of neutron emission in coincidence with fission fragments indicate a strong enhancement of the neutron multiplicity preceding fission compared with statistical model calculations. This enhancement has enabled the determination of the reduced nuclear dissipation coefficient  $\beta$  which, in turn, indicates that nuclear collective motion is overdamped. We examine some possible sources of error in this determination and speculate on the consequences of the obtained value of  $\beta$ .

### Introduction

Over the past decade, numerous measurements of charged-particle and neutron emission in coincidence with fission fragments in Heavy-Ion reactions have been made<sup>1-4</sup>. The dominant feature in all the measurements was the increase in the multiplicity of particles preceding fission compared to expectations based on statistical-model calculations. The implication of this enhanced emission is that the compound nucleus, formed by the colliding heavy-ions, moves relatively slowly towards scission, compared to the time scale of particle emission. The particles can be emitted as the collective deformation coordinate starts moving towards the saddle point and up to the point when the fragments have completely separated and obtained their final relative velocity. Details of the model used to analyze the data of ref. 1 were presented in ref. 5. In this report we will analyze several sources of errors which affect the value of the reduced dissipation coefficient  $\beta$ , inferred from these models. We find that despite a possible uncertainty in the magnitude of  $\beta$ , the conclusion that the motion of the fission coordinate is overdamped remains unchallenged. We examine possible consequences of this high dissipation, which include: 1) the inherent non equilibrium nature of the fission decay of  $^{158}\text{Er}$ , and 2) the effect of the reaction channel dependence of compound nucleus decay, which has been observed in the  $^{156}\text{Er}$  system<sup>b</sup>.

### Experimental

The experimental techniques that were used have been described in detail in previous publications: Neutrons are detected, in coincidence with fission fragments, using a combined time of flight and pulse shape discrimination technique. The angular distribution of the neutrons, with respect to the

fragments, can be fit using a model which contains three emission sources. 1) Non-equilibrium neutrons that can be described by a moving-source model, 2) Neutron emission from the composite system, which is approximately isotropic in the C.M. system, and 3) Neutron emission from the fission fragments during and after acceleration. The efficiency of the neutron detectors is calculated for the detector pulse-height threshold that we select and checked by neutron measurements with a  $^{252}\text{Cf}$  source mounted in  $2\pi$  geometry on a solid-stated surface-barrier detector. For the  $^{160,142}\text{Nd}$  system at 207 MeV beam energy, we obtained  $2.7 \pm 0.4$  neutrons preceding fission; in addition,  $0.9 \pm 0.1$  non-equilibrium neutrons are emitted. In a recent paper, Hinde *et al* obtained  $4.2 \pm 0.3$  neutrons preceding fission for the same system at a beam energy of 178 MeV<sup>4</sup>. Evaporation calculations using PACE2<sup>7</sup> (which reproduce xn results on rare-earth nuclei) predict 7.5 neutrons in coincidence with evaporation residues. We obtained  $5.7 \pm 0.2$  neutrons, a discrepancy we attribute to contamination of the residue singles spectrum, and which has no obvious bearing on the fission neutron results. If, however, we arbitrarily assume that we have a normalization error in the fission data, the maximum factor that should be applied to the data is  $7.5/5.7$ . The renormalized number of neutrons preceding fission would then be  $(2.7 \pm 0.4) \times (7.5/5.7) = 3.6 \pm 0.5$  which is still lower than Hinde's value extrapolated to 207 MeV<sup>4</sup>. We consider this discrepancy to be due, most probably, to angular-momentum effects. In our original analysis, we neglected it since we estimated it to be only a few-percent effect on the in-plane angular distribution of neutrons. This estimate is correct for neutrons in coincidence with evaporation residues. However, when considering neutron emission preceding fission, one is dealing with an angular momentum window spanning approximately 65 to 72M for the  $^{158}\text{Er}$  system. If the quantization axis is defined as the perpendicular to the plane containing the beam axis and the fission axis, evaporation calculations<sup>7</sup> show that the angular distribution of neutrons preceding fission is sharply peaked in-plane as shown in Fig. shown in Fig. 1a. If the quantization axis is defined as the beam axis, the neutron angular distribution is significantly forward peaked in this angular momentum window (Fig. 1b). This effect would lead us to underestimate the pre-fission neutron multiplicity when assuming an isotropic distribution. In fact, we do not know how well the quantization axis is defined with respect to the beam axis and the reaction plane: The fission process breaks the azimuthal symmetry of the reaction, but does not select a well defined axis perpendicular to the plane.




Fig. 1. Angular distribution of neutrons preceding fission. a) Quantization axis perpendicular to plane. (Angle is with respect to normal to plane). b) Quantization axis perpendicular to beam axis. (Angle is with respect to beam axis) Results are for  $l=65M$ . There is no significant difference for  $l=72M$ .

Other effects should be considered when comparing the two results. These are -

1) The shape of the neutron evaporation spectrum we used was  $E^\alpha e^{-E/T}$ ; The parameter  $\alpha$  was taken from PACE2 calculations<sup>7</sup> to be 0.6, which is close to the value determined by Madland and Nix<sup>8</sup>. In comparison, Hinde used  $E e^{-E/T}$ , where  $E$  is the neutron C.M. kinetic energy. There could also be a systematic error in our subtraction of the non equilibrium neutron component at low energies, due to our assumption of the spectrum shape being  $E e^{-E/T}$ . If we were to assume a  $E^\alpha e^{-E/T}$  shape with  $\alpha \neq 1$  we would obtain somewhat different multiplicities.

2) There is an angular correlation between the plane defined by the fission fragments and the beam axis, and the emission direction of non-equilibrium particles<sup>9</sup>. The angular distribution is not azimuthally symmetric around the beam axis and the effect of this distribution needs to be considered in both experimental configurations.

3) Our configuration employs two large solid-angle gas detectors to detect the fission fragments. The detectors were both position sensitive which enabled us to eliminate edge effects. The configuration of Hinde et al is more constrained - it is conceivable that the fragment coincidence selection requirement introduces a bias into the measured neutron distributions<sup>10</sup>.

4) An important parameter that these experiments should provide is the "little-a" parameter  $a_n$ . We find that  $a_n = A/f$ , where A is the atomic mass and  $f = 7.5 \pm 1.5$ ; Hinde et al use a value of  $f = 10$ . These values of  $a_n$  result in an error of approximately  $\pm 50\%$  in the calculated lifetimes of the emitted neutrons. In principal, this parameter can be determined for the compound nucleus by accurate measurements of neutron spectra in coincidence with evaporation residues.

### Theory

The theoretical apparatus we used to determine the reduced nuclear dissipation coefficient  $\beta$  also contains many assumptions and simplifications in order to achieve its goal. These need to be clarified, and, if possible, closely scrutinized.

An important assumption often made is that the non-equilibrium neutrons are emitted on a much shorter time scale than the equilibrium neutrons. This enables us to analyze the data in terms of two distinct neutron sources: The first - a non equilibrium source - completes its emission before the second - a compound-nucleus (equilibrium) source - commences neutron emission. In a recent paper, Blann<sup>11</sup> calculates the equilibration time for the  $^{160,60}\text{Ni}$  system and obtains  $(4-5)10^{-22}$  seconds at a C.M. energy comparable to that of our 207 MeV  $^{160}$  measurement. This is about 4 times faster than the emission time of the first neutron which would justify the consideration of the two distinct sources.

An additional assumption often made is that the various parameters, considered in the evaporation and diffusion problems, are temperature independent. The temperature dependence of the fission barrier of  $^{208}\text{Pb}$  has been calculated by Guet et al<sup>12</sup>. They find that at  $T = 2.6$  MeV (the temperature of  $^{158}\text{Fr}$  following non equilibrium neutron emission), the

fission barrier is reduced by 30% compared to the  $T=0$  value. At  $T=2.0$  MeV (after emission of the pre-fission neutrons) the barrier is still 20% lower than the  $T=0$  value. Incorporation of this temperature dependence would decrease the number of pre-fission neutrons in the framework of standard statistical model calculations. In the diffusion model, the motion over the saddle-point would proceed more rapidly, leading us to increase  $\beta$  to retain the agreement between the calculations and the measured values of the pre-fission neutron multiplicity.

$\beta$  is also assumed to be independent of temperature in the framework of the Wall-Window formalism<sup>13</sup>. At energies close to the fission barrier, we may expect  $\beta$  to decrease due to the lack of available states for quantum transitions. Such an effect is not presently considered. One may also need to consider quantum Brownian motion<sup>14</sup> at these energies.

### Discussion

The possible discrepancies between experiments and the existing uncertainties in the model applied to the data do not seem to be sufficient to challenge the major conclusion obtained by the various groups involved in neutron emission studies: The motion towards scission is overdamped! The debate is only over the question whether  $\beta \sim 6$  or whether  $\beta \geq 10$  in units of  $10^{21} \text{sec}^{-1}$ . In the following, we use the value of  $\beta = 6 \times 10^{21} \text{sec}^{-1}$  we have obtained using four reactions leading to the same composite system with a very similar angular-momentum window<sup>1</sup>. We can now follow the evolution of  $\Gamma_f$ , the fission decay width, and compare it to the total particle decay width  $\Gamma_p$  for different values of the angular-momentum  $l$ . The results are presented in Fig. 2. For  $l=65\hbar$ , asymptotically,  $\Gamma_f \sim \Gamma_p$ . This is the angular momentum, above which, fission exceeds charged-particle emission. For  $l=40\hbar$ ,  $\Gamma_f$  passes  $\Gamma_p$  after  $t = 1.8 \times 10^{-20}$  seconds. At this point, fission decay will dominate, even though  $\Gamma_f$  is still a factor of three below its equilibrium value. The situation is even more acute for  $l=25\hbar$ , where fission will occur, on average, when  $\Gamma_f$  reaches 1/10 of its asymptotic value. This implies that even though we have considered a reaction in which a "compound nucleus" is formed inside a well-defined saddle point, the nucleus may fission long before the fission degree of freedom is equilibrated with the other degrees of freedom. This seems somewhat

paradoxical since the "compound nucleus" concept is construed to imply complete equilibration of all degrees of freedom.

The saddle-point approach to fission width calculations is assumed to be valid as long as the fission barrier  $B_f(l)$  is greater than the temperature at the saddle-point<sup>15</sup>. Indeed, this was the rationale behind our selection of reactions and beam energies for this study. The selection resulted in partial waves between  $l=65\hbar$  and  $l=72\hbar$ , where  $B_f(l) \geq T$  - we had assumed that this partial wave window in  $^{158}\text{Er}$  will avoid the complications associated with "quasi-fission". Nevertheless, we find that at the highest excitation energies, the transition over the fission barrier occurs when the fission probability is far below its asymptotic (equilibrium) value!




Fig. 2. Calculated decay widths as a function of time. Horizontal lines are  $\Gamma(l)$ , the total particle decay width. Curved lines are  $\Gamma_f(l,t)$ . The dotted line is for  $65\hbar$ , the dashed line for  $70\hbar$  and the solid line for  $75\hbar$ .

Another possible implication of the large degree of dissipation pertains to the  $^{64}\text{Ni} + ^{92}\text{Zr}$  reaction<sup>9</sup>. Differences between evaporated neutron multiplicities in this reaction and in the  $^{120,144}\text{Sm}$  reaction have been considered evidence for non statistical behavior in the Ni-Zr system. The possibility of a super deformed minimum in the potential energy surface of the fusing nuclei has also been considered in this context<sup>10</sup>. We wish to



point out that the relative kinetic energy between the Ni and the Zr nuclei is comparable to that between outgoing fission fragments of the  $^{156}\text{Er}$  system. Thus, when the distance between the individual centers-of-mass of the Ni and Zr decrease and approach that of the saddle point, the motion should be described by the Fokker-Planck equation. This implies a very slow formation time for the compound nucleus; during this time, particle emission can take place from the deformed fusing system. Indeed, we calculate<sup>7</sup> the lifetime of the first neutron to be  $\sim 10^{-13}$  sec which is comparable to the transit time over the fission barrier for near-symmetric systems<sup>1</sup>. However, why this should result in a suppression of neutron emission<sup>6</sup> is still not clear.

The author would like to thank Dr. F. Plasil for reminding him of the importance of angular momentum, and Dr. J. R. Nix for other valuable comments.

#### References

1. A. Gavron et al., Phys. Rev. C35 (1987), 579.
2. D. Hilscher, D. J. Hinde and H. Rossner, Fission Dynamics of Hot Nuclei at Temperatures between 1 and 5 MeV, Proceedings of the Texas A&M Symposium on Hot Nuclei, Eds. S. Shlomo, R.P. Schmitt and J. B. Natowitz, World Scientific Publishers (1988), and references therein.
3. J. O. Newton, Nuclear Fission Induced by Heavy Ions, Submitted to Soviet Journal of particles and Nuclei, Australian National University preprint ANU-P/1024, September, 1988 and references therein.
4. D. J. Hinde et al., Phys. Rev. C37 (1988), 2932.
5. P. Grangé et al., Phys. Rev. C34 (1986), 209.
6. P. V. F. Janssens et al., Phys. Lett. B181 (1986), 16 and references therein.
7. A. Gavron, Phys. Rev. C21 (1980), 230.
8. D. G. Madland and J. R. Nix, Nucl. Sci. and Eng. 81 (1982), 213.
9. M. B. Tsang et al., Phys. Rev. Lett. 52 (1984), 1367.
10. A. Gavron, Nucl. Instr. and Meth. 115 (1974), 99.
11. M. Blann, Phys. Rev. C31 (1985), 1245.
12. C. J. J. Janssens et al., Phys. Lett. B205 (1988), 427.
13. See e.g. J. R. Nix and A. J. Sierk, New Picture of Dissipation in Heavy-Ion Reactions and other Collective Phenomena, International Symposium on Perspectives in Nuclear Physics, Madras, India, Jan. 19-23, 1987, Preprint LA UR 87-131.
14. Xizhen Wu et al., J. Phys. G14 (1988), 1043.
15. V. M. Strutinski, Sov. J. Nucl. Phys. 19 (1974), 107.