

LA-UR -81-1815

MASTER

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

TITLE Fertile-To-Fissile and Fission Measurements for Depleted Uranium Bombarded by 800-MeV Protons

AUTHOR(S) G. J. Russell, J. S. Gilmore, H. Robinson, and R. E. Praed

SUBMITTED TO Fifth Meeting of the International Collaboration on Advanced Neutron Sources (ICANS-V), Jülich, West Germany, 22-26 June, 1981

DISCLAIMER

By acceptance of this article the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution or to allow others to do so for U.S. Government purposes. The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

Fertile-To-Fissile and Fission Measurements
for Depleted Uranium Bombarded by 800-MeV Protons

G. J. Russell, J. S. Gilmore, H. Robinson, and R. E. Prael
Los Alamos National Laboratory
Los Alamos, New Mexico 87545, U.S.A.

ABSTRACT

Axial distributions of fertile-to-fissile conversions (^{238}U to ^{239}Pu) and fissions have been measured for a thick depleted uranium target bombarded by 800-MeV protons. We determine the ^{239}Pu production by measuring the amount of ^{239}Np produced. We integrate the axial distributions to get the total conversions and fissions occurring in the target. Our preliminary experimental results give 3.81 ± 0.19 ^{239}Np atoms produced per incident proton and 5.59 ± 0.56 fissions per incident proton. Corresponding calculated results are 3.44 ± 0.05 and 3.93 ± 0.06 . The computations did not include the effects of high-energy fission competition with evaporation. We also report measured axial distributions of ^{237}U and eleven fission products produced in the target. Our preliminary experimental data give 0.95 ± 0.05 ^{237}U atoms made per incident proton.

INTRODUCTION

As part of the Fertile-to-Fissile Conversion (FERFICON) program^{1,2} at the Los Alamos National Laboratory, we have measured (in a thick target of depleted uranium bombarded by 800-MeV protons): a) ^{239}Pu production, b) fission, and c) ^{237}U formation. We determine ^{239}Pu production by measuring the amount of ^{239}Np formed. Other laboratories^{3,4} have made similar measurements (at proton energies <800 MeV) by observing both the radial and axial distributions of the products of interest. Our experimental approach differs significantly (as suggested by one of the authors - J. S. Gilmore) in that, we combined depleted uranium foils to integrate the product formation radially, and only explicitly measured the axial distribution for the products of interest. This substantially reduces the number of samples to be counted, and the number of gamma-ray spectra to be analyzed. We integrate the measured axial distributions over the target to obtain the total number of each reaction.

The data described here are relevant to spallation neutron source development, accelerator breeder technology, and validating computer codes used in these applications (model evaluation).

We will compare our experimental data with calculated predictions using the following Monte Carlo codes: a) the Oak Ridge National Laboratory (ORNL) code HETC⁵ for particle transport $\gtrsim 20$ MeV, and b) the Los Alamos code MCNP⁶

for neutron transport ≤ 20 MeV. The present Los Alamos version of HETC does not include fission when predicting particle production from nucleon and pion collisions with a fissile nucleus. A version of HETC which accounts for this high-energy fission process in uranium will soon be released by ORNL⁷; Los Alamos has requested this version of HETC from ORNL.

The 800-MeV proton source is the Clinton P. Anderson Meson Physics Facility (LAMPF).⁸ We conduct the experiments at the Weapons Neutron Research facility (WNR)⁹--see Fig. 1. We describe here our experimental setup, show some preliminary results, and compare some of our data with calculated predictions.

EXPERIMENTAL SETUP AND PROCEDURES

The location of this 'conversion' experiment in the WNR beam channel is illustrated in Fig. 2. We used a 37-rod clustered target as shown in Fig. 3. The physical characteristics of the target are given in Table I. The axial distributions of ^{239}Np , 11 fission products, and ^{237}U (a spallation product) were determined from 171 (3.239-cm diam by 0.0062-cm thick) depleted uranium foils. Nineteen weighed and matched (~ 1 g) foils were placed in each of seven planes perpendicular to the target axis and on the front and back target faces (see Fig. 4). Each of the 9 planes contained one foil in the central rod and 18 foils loaded symmetrically in 3 of the 6 target sectors (see Fig. 3). We chose this loading for mechanical reasons and to minimize any effects from misaligning the proton beam, which was focused on the central rod.

After an irradiation of 4.3×10^{15} protons, we prepared nine solutions for counting by dissolving the foils in hydrochloric and nitric acids. For each plane, a representative sample was obtained by mixing one-half the solution of the central-rod foil with the solution of the remaining eighteen foils. Five percent (5.00 ml) of each combined solution was used for gamma-ray counting.

To determine the number of protons striking the target, we placed a 0.0254-cm-thick Al monitor foil (sandwiched between two 0.00254-cm-thick Al guard foils to compensate for recoil losses) ~ 65 cm in front of the target. We used the number of $^{27}\text{Al}(p,x)^7\text{Be}$, $^{27}\text{Al}(p,x)^{22}\text{Na}$, and $^{27}\text{Al}(p,3p)^{24}\text{Na}$ reactions occurring in the monitor foil to determine the incident proton dose. The guard foils were sufficiently thick (~ 7 mg/cm²) to compensate for ^7Be , ^{22}Na , and ^{24}Na recoil losses from the central monitor-foil.¹⁰ The number of protons determined from each of these reactions is given in Table II; we ultimately use the $^{27}\text{Al}(p,3p)^{24}\text{Na}$ reaction in our normalizations, because the cross section for this reaction is known best.¹⁰ We located the proton beam center from the discoloration of a cellophane foil that covered one of the Al guard foils, and measured the proton beam profile by counting concentric rings cut from one of the guard foils (see Fig. 5).

All samples were counted using a Ge(Li) detector and associated pulse height analyzer, which had been calibrated against a mixed radionuclide

gamma-ray reference standard.* The gamma-ray spectra were analyzed by the GAMANAL computer program.¹¹ After combining data from five or more counts by the CLSQ computer code,¹² we calculated the atoms of each nuclide formed using the specific gamma-rays, absolute intensities, and half-lives listed in Table III. Two of the isotopes (¹⁰³Rh and ¹⁴⁷Nd) had to be resolved from interfering activities by decay. We corrected all nuclide production for decay during irradiation and for gamma-ray attenuation in the sample.

EXPERIMENTAL RESULTS AND CONCLUSIONS

In Figs. 6-8, we show the measured number of atoms produced per proton per gram of uranium for eleven fission products. The shape of these (unnormalized) fission mass-yield curves do not change appreciably from front to back of the target. This latter point is further illustrated in Fig. 9, where we show the ratio of each fission product to ⁹⁹Mo as a function of axial position.

As shown in Fig. 10, the apparent fission-yield curve from this experiment resembles the known ~ 14.7-MeV-neutron fission of ²³⁸U.¹³ In Fig. 10, we compare this latter curve with our (average) measured value for each fission product. We obtained our yields by ratioing our measured values to that of ⁹⁹Mo; our 'best' estimate of the ⁹⁹Mo absolute yield is (5.7 ± 0.5)%. We selected our ⁹⁹Mo yield by envisioning curves whose summation would be noticeably greater or lower than 200% if the ⁹⁹Mo yield were as high as 6.2% or as low as 5.2%, respectively.

The axial distributions of the fissions (based on a ⁹⁹Mo yield of 5.7%), ²³⁹Np (²³⁹Pu precursor), and ²³⁷U are shown in Fig. 11 and tabulated in Table IV. The axial fission distribution is an important practical consideration (from an energy deposition viewpoint) when designing a uranium target for a spallation neutron source,¹⁴ while the peak in the axial distribution of ²³⁹Np is an aid in locating moderators to maximize low-energy (< 1 eV) neutron production from a uranium target.

The total number (per proton) for each reaction is

$$\text{Total}_i = \frac{M}{\ell \cdot p} \int N_i(z) dz \quad , \quad (1)$$

where M is the mass of the target in grams, ℓ is the target length in cm, p is the number of protons, and N_i is the number of atoms produced per gram of uranium. We evaluated the integral using Simpson's rule. Table V lists the measured ²³⁹Np, fission, and ²³⁷U production per incident proton from the 30.46-cm-long target. In Table V, we also show some calculated predictions for two target lengths; the longer target was used in the Los Alamos FERFICON water-bath measurements.¹⁶ When high-energy fission¹⁵ is neglected in the computations, our preliminary comparisons between experiment and calculations

*A Osham Corporation solution number R9/270/46.

indicate: a) that neutron production is underestimated by $\sim 10\%$, and b) that the number of fissions is underestimated by $\sim 42\%$. Note that the calculated quantities do not change appreciably for the 40.64-cm target compared to the 30.46-cm target. We are still evaluating our data and hope to report a few more fission and spallation product distributions in a final publication. For comparison with experimental data, we will calculate the axial distribution of various products, and the total product formation in the target. We will make a similar measurement using a thorium target in the near future.

ACKNOWLEDGMENTS

This work was performed under the auspices of the U. S. Department of Energy. We would like to acknowledge the support and encouragement of R. Woods, and useful discussions with I. M. Thorson, A. J. Miller, and J. H. Cook. We appreciate the help of J. R. Baldonado and K. J. Hughes in setting up the experiment and of R. D. Ryder and H. M. Howard for beam tuning and assistance during the irradiation. The target canister was remotely disassembled by A. G. Nicol and F. H. Newbury. We acknowledge the help of G. W. Knobeloch and J. Drake for preparing samples, G. W. Butler for advice and assistance in counting and the use of GAMANAL and CLSO computer programs, I. Binder for processing the Al monitor foils, and all the CNC-11 counting room personnel.

REFERENCES

1. G. J. Russell, et al., "Spallation Target-Moderator-Reflector Studies at the Weapons Neutron Research Facility," Symp. on Neutron Cross Sections from 10-50 MeV, Brookhaven National Laboratory, Upton, NY, May 12-14, 1980, BNL-NCS-51245, Vol. I, pp. 169-192.
2. G. J. Russell, et al., "Measurements of Spallation Target-Moderator-Reflector Neutronics at the Weapons Neutron Research Facility," Proc. of the 4th Meeting of the International Collaboration on Advanced Neutron Sources (ICANS-IV), National Laboratory for High Energy Physics (KEK), Tsukuba, Japan, October 20-24, 1980, KENS report II (March 1981).
3. R. G. Vasil'kov, et al., "Neutron Multiplication in Uranium Bombarded with 300-600-MeV Protons," Atomnaya Energiya, 44, No. 4, pp. 329-335 (1978).
4. I. M. Thorson, private communication.
5. K. C. Chandler and T. W. Armstrong, "Operating Instructions for the High Energy Nucleon Meson Transport Code HETC," Oak Ridge National Laboratory report ORNL-4744 (January 1972).
6. W. L. Thompson, ed., "MCNP - A General Monte Carlo Code for Neutron and Photon Transport," Los Alamos Scientific Laboratory report LA-7396-M (November 1979).
7. F. S. Alsmiller, et al., "A Phenomenological Model for Particle Production from the Collisions of Nucleons and Pions with Fissile Elements at Medium Energies," Oak Ridge National Laboratory report ORNL/TM-7528 (March 1981).
8. G. J. Russell, et al., "The WNR Facility -- A Pulsed Spallation Neutron Source at the Los Alamos Scientific Laboratory," Intl. Conf. on Neutron Physics and Nucl. Data for Reactors and Other Applied Purposes, Harwell, England (1978).
9. M. S. Livingston, "LAMPF A NUCLEAR RESEARCH FACILITY," Los Alamos Scientific Laboratory report LA-6878-MS, UC-28 and UC-34 (September 1977).
10. J. B. Cumming, et al., "Absolute Cross Sections for the $^{27}\text{Al}(p,3p)^{24}\text{Na}$ Reaction at 2.8 and 0.8 GeV," Nucl. Inst. and Meth., 180, pp. 37-44 (1981).
11. R. Gunnink and J. B. Niday, "Computerized Qualitative Analysis by Gamma-Ray Spectrometry," University of California NTIS, USAEC report UCRL-51061 (1972).
12. J. B. Cumming, "Applications of Computers to Nuclear and Radiochemistry," National Academy of Sciences National Research Council report NAS-NS-3107, pp. 25-33 (1962).

13. B. F. Rider and M. E. Meek, "Compilation of Fission Product Yields," Vallecitos Nuclear Center report NEDO-12154-2E (1978).
14. Y. Ishikawa, et al., "Proceedings of the 4th Meeting of the International Collaboration on Advanced Neutron Sources (ICANS-IV)," Tsukuba, Japan, 20-24 October 1980, National Laboratory for High Energy Physics report KENS-II (March 1981).
15. E. K. Hyde, The Nuclear Properties of the Heavy Elements, (Prentice-Hall, Inc., Englewood Cliffs, New Jersey, 196) Vol. III, Ch. 11.

TABLE I
PHYSICAL CHARACTERISTICS
OF DEPLETED URANIUM TARGET

DENSITY (g/cm ³)	DIAMETER (cm)	LENGTH (cm)	²³⁵ U CONTENT (wt %)
19.04	19.709 ^a	30.460	0.251

^aEffective diameter ($D = d \sqrt{n}$) for a 37-rod clustered target with an individual rod diameter of 3.2393 cm.

TABLE II
Al MONITOR FOIL DATA

REACTION	CROSS SECTION USED (mb)	MEASURED NUMBER OF PROTONS	STATISTICAL COUNTING ERROR (%)
²⁷ Al(p,x) ⁷ Be	5.7	4.25×10^{15}	0.9
²⁷ Al(p,x) ²² Ne	13.6	4.30×10^{15}	1.1
²⁷ Al(p,3pn) ²⁴ Ne	10.8	4.32×10^{15}	0.3

**TABLE II
NUCLEAR PARAMETERS USED TO CALCULATE
NUCLIDE YIELDS**

ISOTOPE	HALF LIFE (DAYS)	GAMMA ENERGY (keV)	INTENSITY (γ/d)
⁹⁵ Zr	64.05	756.71	0.546
⁹⁷ Zr	0.704	743.36	0.926
⁹⁹ Nb	2.767	140.51	0.8014
¹⁰³ Ru	39.45	497.08	0.8637
¹⁰⁵ Rh	1.473	319.4	0.1960
¹¹² Pd	0.838	617.4	0.4289
¹¹⁵ Cd	2.208	336.23	0.459
¹³² Te	3.25	954.56	0.1659
¹⁴⁰ Ba	12.8	487.03 1596.18	0.4463 0.9540
¹⁴³ Ce	1.375	293.26	0.4130
¹⁴⁷ Nd	11.04	531.10	0.1310
²³⁷ U	6.75	208.0	0.2170
²³⁹ Np	2.35	277.6	0.1387
⁷ Be	53.20	477.6	0.103
²² Na	949.6	1274.6	0.9995
²⁴ Na	0.625	1389.2	1.000

TABLE IV
MEASURED ^{239}Np AND ^{237}U PRODUCTION AND
NUMBER OF FISSIONS IN THE DEPLETED URANIUM TARGET

FOIL POSITION	DISTANCE FROM TARGET FRONT FACE (cm)	^{239}Np PRODUCTION (ATOMS/PROTON-GRAM)^a	^{237}U PRODUCTION (ATOMS/PROTON-GRAM)^a	NUMBER OF FISSIONS (FISSIONS/PROTON-GRAM)^b
1	0.00	1.69×10^{-5}	6.75×10^{-6}	3.65×10^{-5}
2	2.51	2.78×10^{-5}	1.07×10^{-5}	5.76×10^{-5}
3	5.01	3.40×10^{-5}	1.08×10^{-5}	6.04×10^{-5}
4	7.52	3.49×10^{-5}	9.62×10^{-6}	5.55×10^{-5}
5	10.02	3.19×10^{-5}	7.82×10^{-6}	4.63×10^{-5}
6	13.03	2.84×10^{-5}	6.08×10^{-6}	3.75×10^{-5}
7	17.04	2.16×10^{-5}	4.00×10^{-6}	2.59×10^{-5}
8	22.54	1.26×10^{-5}	2.04×10^{-6}	1.38×10^{-5}
9	30.46	4.23×10^{-6}	4.79×10^{-7}	3.24×10^{-6}

^aNominal estimated error is $\pm 5\%$.

^bNominal estimated error is $\pm 10\%$.

TABLE V
PRELIMINARY EXPERIMENTAL DATA
COMPARED WITH CALCULATED RESULTS

	EXPERIMENTAL TARGET LENGTH (30.46 cm)	CALCULATION ^a	
		TARGET LENGTH (30.46 cm)	TARGET LENGTH (40.64 cm)
²³⁹ Np PRODUCTION (atoms/protons)	3.18 · 0.19	3.46 · 0.05	3.71 · 0.05
NUMBER OF FISSIONS (fissions/proton)	5.59 · 0.56	3.93 · 0.06	4.09 · 0.06
²³⁷ U PRODUCTION (atoms/proton)	0.95 · 0.05	----	----

^aThe effects of fission competing with evaporation were not included in the calculations.




Fig. 1. General layout of the WNR showing the two target areas. The high-current target is located in a vertical proton beam and viewed by 11 horizontal flight paths. The low-current target is located in a horizontal proton beam and viewed by 11 horizontal flight paths and one vertical flight path.




Fig. 2. Section thru the WNR beam channel showing the location of the FERRICON conversion experiment.




Fig. 3. Illustration of the 37-rod clustered target and the location of the foils in the array.




Fig. 4. Illustration showing the foil positions within a rod.




Fig. 5. Measured spatial distribution of the proton beam.




Fig. 6. Measured production of various atoms as a function of target axial position. The position $z = 0.00$ cm is the front target face where the ^{235}U - ^{235}U protons were incident.




Fig. 7. Measured production of various atoms as a function of target thickness and position. The position at $z = 0.50$ cm is the front target face where the ^{235}U protons were incident.




Fig. 8. Measured production of fission products as a function of fission product mass number. The position $z = 30.46$ cm is the front target face when the ^{235}U products were analyzed.




Fig. 9. Ratio of measured fission product formation to that of ^{99}Mo as a function of target axial-position.




Fig. 10. Fission yield versus mass number for ^{235}U fission by 14.7 MeV neutrons.




Fig. 11. Axial distribution of fissions, ^{239}Np , and ^{237}U in the depleted uranium target.