

LA-UR-79-1511

MASTER

CONF-790660--3

DYNAMICS OF NUCLEAR FISSION AND HEAVY-ION REACTIONS

J. R. Nix and A. J. Sierk

For presentation at the
Topical Conference on Large Amplitude Collective Nuclear Motion
Keszthely, Lake Balaton, Hungary, June 10-16, 1979

By acceptance of this article for publication, the publisher recognizes the Government's (license) rights in any copyright and the Government and its authorized representatives have unrestricted right to reproduce in whole or in part said article under any copyright secured by the publisher.

The Los Alamos Scientific Laboratory requests that the publisher identify this article as work performed under the auspices of the USERDA.



los alamos
scientific laboratory
of the University of California
LOS ALAMOS, NEW MEXICO 87545

An Affirmative Action/Equal Opportunity Employer

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

NOTICE
This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Government, neither the contractor, subcontractors, or their employees, nor any of their liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

UNITED STATES
ENERGY RESEARCH AND
DEVELOPMENT ADMINISTRATION
CONTRACT W-7406-ENG. 36

DYNAMICS OF NUCLEAR FISSION AND HEAVY-ION REACTIONS

J. R. Nix and A. J. Sierk

Theoretical Division, Los Alamos Scientific Laboratory

University of California, Los Alamos, New Mexico 87545, USA

Abstract

We study large-amplitude collective motion in fission and heavy-ion reactions by solving classical equations of motion for the time evolution of the nuclear shape. In the nuclear potential energy of deformation, we calculate the generalized surface energy by means of a double volume integral of a Yukawa-plus-exponential function, which was obtained by requiring that two semi-infinite slabs of constant-density nuclear matter have minimum energy at zero separation. The collective kinetic energy is calculated for nuclear flow that is a superposition of incompressible, nearly irrotational collective-shape motion and rigid-body rotation. Nuclear dissipation is included by means of the Rayleigh dissipation function, which depends upon the physical mechanism that converts collective energy into internal energy. For both ordinary two-body viscosity and a combined wall and window one-body dissipation, we calculate fission-fragment kinetic energies for the fission of nuclei throughout the periodic table and compare with experimental results. Finally, we study explicitly the one-body dynamics of nucleons inside a cylinder colliding with a moving piston by solving exactly the collisionless Boltzmann equation for the distribution function. By examining the relative phases of the pressure at the piston and the piston's velocity, we are able to separately identify a dissipative force and an elastic restoring force.

DYNAMICS OF NUCLEAR FISSION AND HEAVY-ION REACTIONS

J. R. Nix and A. J. Sierk

Theoretical Division, Los Alamos Scientific Laboratory

University of California, Los Alamos, New Mexico 87545 USA

As illustrated in Fig. 1, we study large-amplitude collective motion in fission and heavy-ion reactions by solving classical equations of motion for the time evolution of the nuclear shape.^{1,2} In the nuclear potential energy of deformation, we calculate the generalized surface energy by means of a double volume integral of a Yukawa-plus-exponential function, which was obtained by requiring that two semi-infinite slabs of constant-density nuclear matter have minimum energy at zero separation.^{3,4} The resulting potential for heavy-ion reactions is compared with experimental results in Fig. 2 and with other heavy-ion potentials in Fig. 3. As shown by the solid curves in Fig. 4, our potential reproduces heavy-ion elastic scattering with approximately the same accuracy as does a Woods-Saxon potential (dashed curves).

The collective kinetic energy is calculated for nuclear flow that is a superposition of incompressible, nearly irrotational collective-shape motion and rigid-body rotation.^{1,2} Nuclear dissipation is included by means of the Rayleigh dissipation function, which depends upon the physical mechanism that converts collective energy into internal energy. Figure 5 compares results that have been calculated for various types of dissipation and an earlier version of the potential energy with the result of a time-dependent Hartree-Fock calculation.⁵ Figure 6 illustrates the type of potential-energy surface that is involved in the macroscopic calculations.⁶

Figure 7 shows dynamical trajectories calculated with the Yukawa-plus-exponential potential for both ordinary two-body viscosity¹ and a combined

wall and window one-body dissipation.⁷ In the latter case, the wall formula relative to the center of mass of the entire system is used until the neck radius reaches the indicated value, at which point a transition is made to a wall formula relative to the centers of mass of the two nascent fragments plus a window formula for the neck between them. As shown in Figs. 8 and 9, experimental fission-fragment kinetic energies for the fission of nuclei throughout the periodic table are reproduced by either ordinary two-body viscosity with a coefficient $\mu = 0.015 \text{ TP}$ or by a combined wall and window one-body dissipation with a transition neck radius $r_{\text{neck}} = 2.5 \text{ fm}$.

In Fig. 10 we examine the ability of the wall formula to predict the experimental widths of giant multipole resonances. By comparing the solid curve with the solid circles (giant quadrupole)⁸ and the dashed curve with the open circle (giant octupole),⁹ we see that for both resonances the wall-formula predictions are about three times as large as the experimental values.

Finally, as illustrated in Fig. 11, we study explicitly the one-body dynamics of nucleons inside a cylinder colliding with a moving piston by solving exactly the collisionless Boltzmann equation for the distribution function. By examining the relative phases of the pressure at the piston and the piston's velocity, we are able to separately identify a dissipative force and an elastic restoring force.


Acknowledgments

We are grateful to G. F. Bertsch, K. T. R. Davies, S. E. Koonin, H. J. Krappe, P. Möller, J. W. Negele, and W. J. Swiatecki for their collaboration on some of the studies reported here. This work was supported by the U. S. Department of Energy.

References

1. K. T. R. Davies, A. J. Sierk, and J. R. Nix, Phys. Rev. C 13, 2385 (1976).
2. J. R. Nix and A. J. Sierk, Phys. Rev. C 15, 2072 (1977).
3. H. J. Krappe, J. R. Nix, and A. J. Sierk, Phys. Rev. Lett. 42, 215 (1979).
4. H. J. Krappe, J. R. Nix, and A. J. Sierk, Phys. Rev. C, to be published.
5. J. W. Negele, S. E. Koonin, P. Möller, J. R. Nix, and A. J. Sierk, Phys. Rev. C 17, 1098 (1978).
6. P. Möller and J. R. Nix, Nucl. Phys. A272, 502 (1976).
7. J. Błocki, Y. Boneh, J. R. Nix, J. Randrup, M. Robel, A. J. Sierk, and W. J. Swiatecki, Ann. Phys. (N. Y.) 113, 330 (1978).
8. F. E. Bertrand, Ann. Rev. Nucl. Sci. 26, 457 (1976).
9. M. Sasao and Y. Torizuka, Phys. Rev. C 15, 217 (1977).

MACROSCOPIC APPROACH



Collective coordinates

$$q = q_1, \dots, q_n$$

Fundamental nuclear properties

1. Potential energy $V(q)$
2. Kinetic energy $T(q, \dot{q})$
3. Dissipation $F(q, \dot{q})$

Lagrangian $L = T - V$

Equations of motion

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_i} \right) + \frac{\partial F}{\partial \dot{q}_i} - \frac{\partial L}{\partial q_i} = 0, \quad i = 1, \dots, n$$

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6






Figure 8




Figure 9





Figure 10

Exact Solution of the Boltzmann Equation



$$\frac{\text{Oscillation period}}{\text{Transit time}} = 4,$$

$$\frac{v_0}{v_F} = 0.1$$

-14-




Figure 11